

TUBE REPRESENTATIONS AND TWISTING OF GRADED CATEGORIES

JYOTISHMAN BHOWMICK, SHAMINDRA GHOSH, NARAYAN RAKSHIT, AND
MAKOTO YAMASHITA

ABSTRACT. We study deformation of tube algebra under twisting of graded monoidal categories. When a tensor category \mathcal{C} is graded over a group Γ , a torus-valued 3-cocycle on Γ can be used to deform the associator of \mathcal{C} . We show that it induces a 2-cocycle on the groupoid of the adjoint action of Γ . Combined with the natural Fell bundle structure of tube algebra over this groupoid, we show that the tube algebra of the twisted category is a 2-cocycle twisting of the original one.

1. Introduction

In the theory of quantum symmetries, the concept of *quantum double* provides a powerful guiding principle to understand various aspects of quantum groups. There are several ways to precisely realize it mathematically: originally it appeared as the *Drinfeld double* of Hopf algebras which provides a uniform way to produce solutions of the Yang–Baxter equation, and can be regarded as a Hopf algebraic analogue of the complexification of compact semisimple Lie algebras. A closely related notion is the *Drinfeld center*, as formulated by Drinfeld, Majid, and Joyal and Street independently, in the more general context of tensor categories which generalizes the Drinfeld double. Besides producing braided tensor categories, it turned out to have far reaching roles in the theory of tensor categories, such as a natural framework to consider algebra objects encoding various categorical structures of interest.

While there are several other notable approaches, *tube algebra* is perhaps the most concrete (and combinatorial) formalism to define the quantum double for tensor categories, which was introduced by Ocneanu [Ocn94] in his pioneering study of subfactors and topological quantum field theory. Although at first sight this looks quite different from other definitions of the Drinfeld center, its precise correspondence was clarified through the subsequent work of Longo and Rehren [LR95], and Izumi [Izu00], to name a few.

The goal of this work is to understand the change of tube algebra induced by a change of associator on the tensor category. To be more specific, we consider a change of associator on a tensor category \mathcal{C} graded by a discrete group Γ , induced by group 3-cocycles on Γ ,

Received by the editors 2018-06-03 and, in final form, 2018-09-18.

Transmitted by Gabriella Böhm. Published on 2018-09-23.

2010 Mathematics Subject Classification: 18D10.

Key words and phrases: monoidal category, quantum double, tube algebra.

© Jyotishman Bhowmick, Shamindra Ghosh, Narayan Rakshit, and Makoto Yamashita, 2018. Permission to copy for private use granted.

inspired by the work of Kazhdan and Wenzl [KW93] on the classification of the tensor categories with the fusion rules of quantum $SL(n)$ groups.

A key insight is that the tube algebra $\mathcal{T}(\mathcal{C})$ has a structure of *Fell bundle* [Yam90, Kum98] over the action groupoid \mathcal{G} of Γ acting on itself by the adjoint action. Extending the analysis of pointed categories by Bisch, Das, and the second and the third named authors [BDGR17] (a scheme which was also known to Bántay and others in a slightly different framework, see [Bán90, ACM04, Wil08]), we show that $\mathcal{T}(\mathcal{C}^\omega)$, the tube algebra of \mathcal{C} with its associator twisted by ω , is isomorphic to the 2-cocycle twist of $\mathcal{T}(\mathcal{C})$ by a 2-cocycle on \mathcal{G} induced by ω .

When Γ is a cyclic group, this implies that the Drinfeld center $\mathcal{Z}(\mathcal{C}^\omega)$ of the twisted category is equivalent to $\mathcal{Z}(\mathcal{C})$ as a linear category. The monoidal structure still needs to be modified in the presence of ω , and we give an explicit formula for this twisting. For one thing, when \mathcal{C} is braided and $\mathcal{Z}(\mathcal{C})$ admits a good description as a tensor category, this allows us to determine whether \mathcal{C}^ω is braided or not which we will explain in detail for the Kazhdan–Wenzl categories (Example 5.4).

This paper is organized as follows: in Section 2 we recall basic concepts and fix notation, which can be easily skipped by an expert. In Section 3 we present the precise structure of tube algebras for nonstrict tensor categories, which serves as a basis for the twisting argument. Our main result is in Section 4. Finally, we present several examples in Section 5.

Acknowledgements It is our pleasure to thank Masaki Izumi, Corey Jones, Madhav Reddy, and Shigeru Yamagami for fruitful discussions at various stages of the project.

2. Preliminaries

In this paper, all categories are assumed to be small, and we mostly work with *C*-categories*, although most of our constructions can be carried out in the setting of semisimple tensor categories over more general coefficients. We denote the unit circle by $\mathbb{T} = \{z \in \mathbb{C} \mid |z| = 1\}$.

2.1. C*-TENSOR CATEGORIES. We mainly follow the convention of [NT13].

When \mathcal{C} is a category and X, Y are objects of \mathcal{C} , the morphism space from X to Y will be denoted by $\text{Mor}_{\mathcal{C}}(X, Y)$ or more simply by $\mathcal{C}(X, Y)$. The identity morphism of X is denoted by 1_X . For C*-categories, we always assume that direct sum of objects and images of projections exist in the category. We say that $X \in \text{Obj}(\mathcal{C})$ is *simple* if $\dim \mathcal{C}(X, X) = 1$, and that \mathcal{C} is *semisimple* if $\mathcal{C}(X, Y)$ is finite dimensional for any $X, Y \in \text{Obj}(\mathcal{C})$.

A *C*-tensor category* is given by a C*-category \mathcal{C} , a C*-bifunctor $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, a distinguished object $\mathbb{1} \in \text{Obj}(\mathcal{C})$, and natural unitary transformations $\lambda_X: \mathbb{1} \otimes X \rightarrow X$, $\rho_X: X \otimes \mathbb{1} \rightarrow X$, and

$$\alpha_{X,Y,Z}: (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$$

satisfying a standard set of axioms, most notably the *pentagon equation* saying that the diagram

$$\begin{array}{ccc}
 & ((U \otimes V) \otimes W) \otimes X & \\
 \alpha_{U,V,W} \otimes 1_X \swarrow & & \searrow \alpha_{U \otimes V, W, X} \\
 (U \otimes (V \otimes W)) \otimes X & & (U \otimes V) \otimes (W \otimes X) \\
 \alpha_{U, V \otimes W, X} \searrow & & \swarrow \alpha_{U, V, W \otimes X} \\
 U \otimes ((V \otimes W) \otimes X) & \xrightarrow{1_U \otimes \alpha_{V, W, X}} & U \otimes (V \otimes (W \otimes X))
 \end{array}$$

is commutative. In the following we always assume that $\mathbb{1}$ is simple.

When \mathcal{C} and \mathcal{C}' are such categories, a C^* -tensor functor from \mathcal{C} to \mathcal{C}' is given by (F, F_0, F_2) consisting of:

- a C^* -functor $F: \mathcal{C} \rightarrow \mathcal{C}'$,
- a unitary morphism $F_0: \mathbb{1}_{\mathcal{C}'} \rightarrow F(\mathbb{1}_{\mathcal{C}})$, and
- a natural unitary transformation $F_2: F(X) \otimes F(Y) \rightarrow F(X \otimes Y)$,

satisfying a standard set of compatibility conditions for λ , ρ , and α of \mathcal{C} and \mathcal{C}' . If F is an equivalence of categories, we say that the above C^* -tensor functor is a (unitary monoidal) equivalence of C^* -tensor categories.

A variant of Mac Lane’s coherence theorem implies that any C^* -tensor category is equivalent to a *strict* one, the latter being a C^* -tensor category where λ_X , ρ_X , and $\alpha_{X,Y,Z}$ are all given by identity morphisms. In this paper we carry out various constructions for categories with nontrivial associator α but still with λ_X and ρ_X given by identity morphisms. In practice, we still assume that the ‘starting point’ is given by a strict C^* -tensor category, and deform it to another one in which α is given by scalar multiples of identity morphisms.

In a *rigid* C^* -tensor category \mathcal{C} any object $X \in \text{Obj}(\mathcal{C})$ has a dual, that is, there exist $\bar{X} \in \text{Obj}(\mathcal{C})$ and $R \in \mathcal{C}(\mathbb{1}, \bar{X} \otimes X)$, $\bar{R} \in \mathcal{C}(\mathbb{1}, X \otimes \bar{X})$ satisfying the *conjugate equation* for X :

$$(1_{\bar{X}} \otimes \bar{R}^*) \alpha_{\bar{X}, X, \bar{X}}(R \otimes 1_{\bar{X}}) = 1_{\bar{X}}, \quad (1_X \otimes R^*) \alpha_{X, \bar{X}, X}(\bar{R} \otimes 1_X) = 1_X.$$

(Recall that we are assuming $\lambda_X = 1_X = \rho_X$ in \mathcal{C} .) The number $d(X) = \min_{(R, \bar{R})} \|R\| \|\bar{R}\|$ is called the *intrinsic dimension* of X , where the minimum is taken over all pairs (R, \bar{R}) as above. A solution (R, \bar{R}) satisfying $\|R\| = d(X)^{1/2} = \|\bar{R}\|$ is called a *standard* solution, which we denote by (R_X, \bar{R}_X) . Standard solutions are unique up to unitary morphisms. It implies that for any $X \in \text{Obj}(\mathcal{C})$, the functional Tr_X on $\mathcal{C}(X)$ defined by

$$R_X^* \circ (1_{\bar{X}} \otimes x) \circ R_X = \text{Tr}_X(x) 1_{\mathbb{1}} \quad (x \in \mathcal{C}(X))$$

is a tracial positive functional, called the *canonical picture trace* or *categorical trace*. Having fixed (R_X, \bar{R}_X) and (R_Y, \bar{R}_Y) , whenever $x \in \mathcal{C}(X, Y)$ we denote by x^\vee the morphism in $\mathcal{C}(\bar{Y}, \bar{X})$ satisfying $(1_{\bar{X}} \otimes x)R_X = (x^\vee \otimes 1_Y)R_Y$.

Rigid C*-tensor categories with simple units are automatically semisimple, and we work within this framework.

2.2. GRADED TENSOR CATEGORIES. In a C*-category \mathcal{C} , two full subcategories \mathcal{D} and \mathcal{E} are called *mutually orthogonal* if $\mathcal{C}(D, E) = \{0\}$ holds for all $D \in \text{Obj}(\mathcal{D})$ and $E \in \text{Obj}(\mathcal{E})$. We write $\mathcal{C} = \mathcal{D} \oplus \mathcal{E}$ if in addition any object in \mathcal{C} is isomorphic to a direct sum $D \oplus E$ for some $D \in \text{Obj}(\mathcal{D})$ and $E \in \text{Obj}(\mathcal{E})$, and say that \mathcal{C} is a direct sum of \mathcal{D} and \mathcal{E} . Of course, this has a straightforward extension to a family of subcategories $\{\mathcal{C}_j\}_{j \in J}$.

2.3. DEFINITION. [cf. [Tur10]] *Let Γ be a group. A C*-tensor category \mathcal{C} is called Γ -graded if there exists a collection of mutually orthogonal full subcategories $\{\mathcal{C}_\gamma\}_{\gamma \in \Gamma}$ such that*

- (i) $\mathcal{C} = \bigoplus_{\gamma \in \Gamma} \mathcal{C}_\gamma$, and
- (ii) $X \otimes Y$ is isomorphic to an object in $\mathcal{C}_{\gamma\eta}$ for all $\gamma, \eta \in \Gamma$, $X \in \text{Obj}(\mathcal{C}_\gamma)$, and $Y \in \text{Obj}(\mathcal{C}_\eta)$.

2.4. REMARK. The tensor unit $\mathbb{1}$, being simple, must belong to $\text{Obj}(\mathcal{C}_e)$ by conditions (i) and (ii) where e is the unit of Γ . It then follows that $\bar{X} \in \text{Obj}(\mathcal{C}_{\gamma^{-1}})$ (if it exists) for all $X \in \text{Obj}(\mathcal{C}_\gamma)$.

Note that if \mathcal{C} is graded over Γ , the ‘support’ of a grading defined as

$$\{\gamma \in \Gamma \mid \mathcal{C}_\gamma \text{ contains a nonzero object}\}$$

is a subgroup of Γ . Without losing generality we may assume that the support is always the entire group. Let us also note that when \mathcal{C} is Γ -graded and $\pi: \Gamma \rightarrow \Lambda$ is a (surjective) group homomorphism, we obtain a Λ -grading on \mathcal{C} by setting $\mathcal{C}_{\gamma'} = \bigoplus_{\pi(\gamma)=\gamma'} \mathcal{C}_\gamma$ for $\gamma' \in \Lambda$.

Any rigid semisimple C*-tensor category \mathcal{C} with simple unit admits a grading $\mathcal{C} = \bigoplus_{\gamma \in \text{Ch}(\mathcal{C})} \mathcal{C}_\gamma$ over the *universal grading group* $\text{Ch}(\mathcal{C})$ (also called the *chain group*) in the sense that any Γ -grading on \mathcal{C} is induced by a unique homomorphism $q: \text{Ch}(\mathcal{C}) \rightarrow \Gamma$ as above.

Concretely, $\text{Ch}(\mathcal{C})$ is constructed as follows. Fix a set $\mathcal{I} = \text{Irr}(\mathcal{C})$ of representatives from isomorphism classes of simple objects. The elements of $\text{Ch}(\mathcal{C})$ are represented by symbols $[X]$ for $X \in \mathcal{I}$, different symbols possibly representing the same element, subject to the rule $[Z] = [X][Y]$ whenever there is a nonzero morphism from Z to $X \otimes Y$, or equivalently, when Z is isomorphic to a subobject of $X \otimes Y$. From this description it immediately follows that $[\mathbb{1}]$ is the unit of $\text{Ch}(\mathcal{C})$, $[X]^{-1} = [\bar{X}]$. For $\gamma \in \text{Ch}(\mathcal{C})$, define \mathcal{C}_γ as the full subcategory of \mathcal{C} , whose objects decompose as direct sum of simple objects X satisfying $[X] = \gamma$. Clearly, $\{\mathcal{C}_\gamma\}_{\gamma \in \text{Ch}(\mathcal{C})}$ is a mutually orthogonal collection of subcategories satisfying conditions of Definition 2.3.

2.5. EXAMPLE. [Temperley–Lieb category] Consider the *Temperley–Lieb category* $\mathcal{C}_{\delta,\varepsilon}^{\text{TL}}$ with modulus $\delta \geq 2$ and the associativity sign $\varepsilon \in \{\pm 1\}$, which can also be seen as the category of finite dimensional representations of the compact quantum group $\text{SU}_q(2)$ [Wor87], with $q \in \mathbb{R}^\times$ satisfying $|q + q^{-1}| = \delta$ and $\varepsilon = -\text{sgn}(q)$. The set of isomorphism classes of simple objects of $\mathcal{C}_{\delta,\varepsilon}^{\text{TL}}$ is countably infinite. In fact, there is a set $\{X_n\}_{n \geq 0}$ of representatives of such classes indexed by nonnegative integers satisfying:

- (i) X_0 is the trivial object $\mathbb{1}$,
- (ii) $X_m \otimes X_n \cong X_{|m-n|} \oplus X_{|m-n|+2} \oplus \cdots \oplus X_{(m+n)-2} \oplus X_{m+n}$, and
- (iii) $\overline{X_m} \cong X_m$

for all $m, n \geq 0$. The parameters δ and ε (which do not affect $\text{Ch}(\mathcal{C}_{\delta,\varepsilon}^{\text{TL}})$) appear as follows: there is a morphism $R: X_0 \rightarrow X_1 \otimes X_1$, unique up to \mathbb{T} , such that

$$R^* R = \delta 1_{X_0}, \quad (1_{X_1} \otimes R^*) \alpha_{X_1, X_1, X_1}(R \otimes 1_{X_1}) = \varepsilon 1_{X_1}$$

Condition (ii) implies that every X_n is a subobject of $X_1^{\otimes n}$. It follows that the universal grading group $\text{Ch}(\mathcal{C}_{\delta,\varepsilon}^{\text{TL}})$ has to be cyclic generated by $[X_1]$. Again, using condition (ii), one can easily show that $[X_n]$ is either equal to $[\mathbb{1}]$ or to $[X_1]$ according as n is even or odd, which implies $\text{Ch}(\mathcal{C}_{\delta,\varepsilon}^{\text{TL}}) \cong \mathbb{Z}/2\mathbb{Z}$.

2.6. GROUP AND GROUPOID COHOMOLOGY. Let Γ be a (discrete) group, and M be a left Γ -module. The group cochain complex $C^*(\Gamma; M)$ is given by the mapping spaces $C^n(\Gamma; M) = \text{Map}(\Gamma^n, M)$ endowed with the coboundary maps $\delta^n: C^n(\Gamma; M) \rightarrow C^{n+1}(\Gamma; M)$ given by

$$\begin{aligned} \delta^n(\phi)(\gamma_1, \gamma_2, \dots, \gamma_{n+1}) &= \gamma_1 \phi(\gamma_2, \dots, \gamma_{n+1}) \\ &+ \sum_{i=1}^n (-1)^i \phi(\gamma_1, \gamma_2, \dots, \gamma_{i-1}, \gamma_i \gamma_{i+1}, \gamma_{i+2}, \dots, \gamma_{n+1}) + (-1)^{n+1} \phi(\gamma_1, \dots, \gamma_n). \end{aligned} \tag{2.1}$$

A cochain $\phi \in C^n(\Gamma; M)$ is *normalized* if $\phi(\gamma_1, \dots, \gamma_n) = 0$ whenever one of γ_i is e . The subcomplex $\bar{C}^*(\Gamma; M)$ of normalized cochains have the same cohomology as $C^*(\Gamma; M)$.

Next let \mathcal{G} be a groupoid. As usual let us denote by $\mathcal{G}^{(0)}$ its object set, and by $\mathcal{G}^{(n)}$ the set of n -tuples (g_1, \dots, g_n) of composable arrows. Thus, there are maps dom and codom from $\mathcal{G} = \mathcal{G}^{(1)}$ to $\mathcal{G}^{(0)}$ so that g_1 and g_2 are composable if and only if $\text{dom}(g_1) = \text{codom}(g_2)$. We also identify $\mathcal{G}^{(0)}$ as a subset of \mathcal{G} by the embedding $x \mapsto \text{id}_x$.

When M is a commutative group, the complex of normalized cochains on \mathcal{G} with coefficient M is given by [Wes69]

$$\bar{C}^n(\mathcal{G}; M) = \{\psi: \mathcal{G}^{(n)} \rightarrow M \mid \psi(g_1, \dots, g_n) = 0 \text{ if } \exists i: g_i \in \mathcal{G}^{(0)}\},$$

together with the differential $\delta^n: \bar{C}^n(\mathcal{G}; M) \rightarrow \bar{C}^{n+1}(\mathcal{G}; M)$ given by

$$\delta^n(\psi)(g_1, \dots, g_{n+1}) = \psi(g_2, \dots, g_{n+1}) + \sum_{i=1}^n (-1)^i \psi(g_1, \dots, g_i g_{i+1}, \dots, g_{n+1}) + (-1)^{n+1} \psi(g_1, \dots, g_n).$$

Its cohomology is denoted by $H^*(\mathcal{G}; M)$. Of course, we could have used the complex of non-normalized cochains, $C^n(\mathcal{G}; M) = \text{Map}(\mathcal{G}^{(n)}, M)$, to compute $H^*(\mathcal{G}; M)$.

We are particularly interested in the *action groupoid* $\Gamma \ltimes X$ arising from an action of a group Γ on a set X . This groupoid has the object set X , and $\Gamma \ltimes X = \Gamma \times X$ as a set, with domain and codomain maps defined by $\text{dom}(\gamma, x) = x$ and $\text{codom}(\gamma, x) = \gamma x$, and composition given by $(\gamma, x) \cdot (\eta, y) = (\gamma\eta, y)$ whenever $x = \eta y$. In this case $\bar{C}^n(\mathcal{G}; M)$ is nothing but the space $\bar{C}_\Gamma^n(X; M)$ of normalized equivariant cochains on the (discrete) set X with coefficient in M , or what amounts to the same thing, the space $\bar{C}^n(\Gamma; \text{Map}(X, M))$ of normalized cochains on Γ with coefficient $\text{Map}(X, M)$, where Γ acts on $\text{Map}(X, M)$ from left by $(\gamma f)(x) = f(\gamma^{-1}x)$. An explicit cochain isomorphism $\bar{C}^n(\mathcal{G}; M) \ni \psi \mapsto \tilde{\psi} \in \bar{C}^n(\Gamma; \text{Map}(X, M))$ is given by

$$\tilde{\psi}[\gamma_1, \dots, \gamma_n](x_0) = \psi(g_1, \dots, g_n), \tag{2.2}$$

where $g_i = (\gamma_i, x_i) \in \mathcal{G}$ and $x_i = \gamma_i^{-1}x_{i-1}$ for all $1 \leq i \leq n$.

2.7. TWISTING MONOIDAL CATEGORIES BY GROUP 3-COCYCLES. Again let Γ be a group, and let ω be a cocycle in $\bar{C}^3(\Gamma; \mathbb{T})$. So ω is a map from $\Gamma \times \Gamma \times \Gamma$ to \mathbb{T} satisfying

$$\omega(\gamma_1, \gamma_2, \gamma_3)\omega(\gamma_1, \gamma_2\gamma_3, \gamma_4)\omega(\gamma_2, \gamma_3, \gamma_4) = \omega(\gamma_1\gamma_2, \gamma_3, \gamma_4)\omega(\gamma_1, \gamma_2, \gamma_3\gamma_4) \quad (\gamma_i \in \Gamma),$$

and $\omega(\gamma_1, \gamma_2, \gamma_3) = 1$ whenever at least one of γ_i is the unit e of Γ .

When \mathcal{C} is a Γ -graded C^* -tensor category, we can consider a new category \mathcal{C}^ω which

- is same as \mathcal{C} as a C^* -category,
- has the same tensor bifunctor and left and right unit constraints, but
- has the associativity constraint α^ω given as a twist of that of \mathcal{C} by the cocycle ω :

$$\alpha_{X_1, X_2, X_3}^\omega = \omega(\gamma_1, \gamma_2, \gamma_3)\alpha_{X_1, X_2, X_3} \quad (X_i \in \text{Obj}(\mathcal{C}_{\gamma_i})).$$

The new associator α^ω still satisfies the same compatibility conditions as α because ω is a normalized 3-cocycle.

Note that \mathcal{C}^ω has the same fusion rules as \mathcal{C} , and in particular it inherits the Γ -grading of \mathcal{C} . However, typically they are monoidally equivalent (if and) only if ω is a coboundary.

2.8. PROPOSITION. *When \mathcal{C} is a rigid C^* -tensor category, \mathcal{C}^ω is also rigid.*

PROOF. It is enough to see that for any $\gamma \in \Gamma$ and $X \in \mathcal{C}_\gamma$, there is a dual object of X in \mathcal{C}^ω . By our standing assumption X admits a dual object in \mathcal{C} , say given by (Y, R, \bar{R}) . Then Y is an object of $\mathcal{C}_{\gamma^{-1}}$. Moreover we have $\omega(\gamma, \gamma^{-1}, \gamma) = \bar{\omega}(\gamma^{-1}, \gamma, \gamma^{-1})$, since by the 3-cocycle identity on $(\gamma, \gamma^{-1}, \gamma, \gamma^{-1})$ we have

$$\omega(\gamma^{-1}, \gamma, \gamma^{-1})\bar{\omega}(e, \gamma, \gamma^{-1})\omega(\gamma, e, \gamma^{-1})\bar{\omega}(\gamma, \gamma^{-1}, e)\omega(\gamma, \gamma^{-1}, \gamma) = 1$$

and the terms involving e are 1 by the normalization condition on ω . Consequently $R' = R$ and $\bar{R}' = \omega(\gamma^{-1}, \gamma, \gamma^{-1})\bar{R}$ satisfy

$$\begin{aligned} (1_Y \otimes \bar{R}'^*)\alpha_{Y,X,Y}^\omega(R' \otimes 1_Y) &= \bar{\omega}(\gamma^{-1}, \gamma, \gamma^{-1})\omega(\gamma^{-1}, \gamma, \gamma^{-1})1_Y = 1_Y, \\ (1_X \otimes R'^*)\alpha_{X,Y,X}^\omega(\bar{R}' \otimes 1_X) &= \omega(\gamma, \gamma^{-1}, \gamma)\omega(\gamma^{-1}, \gamma, \gamma^{-1})1_X = 1_X. \end{aligned}$$

Thus, the triple (Y, R', \bar{R}') gives a dual of X in \mathcal{C}^ω . ■

2.9. EXAMPLE. The first basic example comes from pointed categories, that is, when the tensor product of every simple object with its dual is isomorphic to the unit object. The category \mathcal{C}_Γ of finite dimensional Γ -graded Hilbert spaces is such an example. Generally, the isomorphism classes of simple objects in a pointed category automatically becomes a group with respect to tensor product where the inverse is given by taking dual; this is the universal grading group. In fact, any pointed C^* -tensor category is unitarily monoidally equivalent to $\mathcal{C}_\Gamma^\omega$ for some group Γ and a (normalized) \mathbb{T} -valued 3-cocycle ω on Γ .

3. Some formulas for nonstrict categories

3.1. ANNULAR ALGEBRAS. In [GJ16], annular algebras were defined assuming strictness of the tensor structure. While the C^* -variant of Mac Lane’s coherence theorem guarantees that there is no loss of generality in doing so, for our purposes it will be useful to work out concrete formulas for non-strict ones. In this section \mathcal{C} denotes a rigid C^* -tensor category with simple unit, and we denote a set of representative of isomorphism classes of simple objects in \mathcal{C} by $\mathcal{I} = \text{Irr}(\mathcal{C})$.

Suppose that $\mathcal{X} = \{X_j\}_{j \in \mathcal{J}}$ is a family of objects of \mathcal{C} which is *full* in the sense that any $S \in \mathcal{I}$ is isomorphic to a subobject of X_j for some j . In order to simplify the notation, let us use the index j in place of X_j when they appear in subscripts, so that α_{S,T,X_j} becomes $\alpha_{S,T,j}$ for example.

For $j, k \in \mathcal{J}$, let $\mathcal{A}_{k,j}$ denote the quotient of the vector space

$$\bigoplus_{S \in \text{Obj}(\mathcal{C})} \mathcal{C}(S \otimes X_j, X_k \otimes S)$$

over the subspace generated by elements of the form

$$f(f' \otimes 1_j) - (1_k \otimes f')f, \quad (f \in \mathcal{C}(T \otimes X_j, X_k \otimes S), f' \in \mathcal{C}(S, T)). \tag{3.1}$$

Further, we will write $\mathcal{A}_{k,j}^S$ for the image of $\mathcal{C}(S \otimes X_j, X_k \otimes S)$ under the quotient map, and the map $\psi_{k,j}^S: \mathcal{C}(S \otimes X_j, X_k \otimes S) \rightarrow \mathcal{A}_{k,j}^S$ will stand for the restriction of the quotient map.

The vector space $\mathcal{A}_{k,j}$ can be identified with the direct sum $\bigoplus_{S \in \mathcal{I}} \mathcal{C}(S \otimes X_j, X_k \otimes S)$, hence each $\psi_{k,j}^S$ is injective (thus bijective) for $S \in \mathcal{I}$. Indeed, for a general $S \in \text{Obj}(\mathcal{C})$ we can take an irreducible decomposition $S \cong \bigoplus_{\alpha} S_{\alpha}$ with $S_{\alpha} \in \mathcal{I}$. Let $v_{\alpha}: S_{\alpha} \rightarrow S$ be the corresponding isometry. When $f \in \mathcal{C}(S \otimes X_j, X_k \otimes S)$, we have $f = \sum_{\alpha} f(v_{\alpha} v_{\alpha}^* \otimes 1_j)$. From the formula (3.1) with $f(v_{\alpha} \otimes 1_j)$ in place of f and $v_{\alpha}^* \otimes 1_j$ in place of f' , we see that f and $\sum_{\alpha} (1_k \otimes v_{\alpha}^*) f(v_{\alpha} \otimes 1_j)$ represent the same element in $\mathcal{A}_{k,j}$. This already shows that $\mathcal{A}_{k,j}$ is a quotient of $\bigoplus_{S \in \mathcal{I}} \mathcal{C}(S \otimes X_j, X_k \otimes S)$. Since for different $S, T \in \mathcal{I}$ there is no nonzero morphism $f' \in \mathcal{C}(S, T)$, the images of $\mathcal{C}(S \otimes X_j, X_k \otimes S)$ for $S \in \mathcal{I}$ are linearly independent.

Let us put $\mathcal{A} = \mathcal{A}(\mathcal{X}) = \bigoplus_{j,k \in \mathcal{J}} \mathcal{A}_{k,j}$, and define a bilinear operation \bullet on \mathcal{A} characterized by

$$\psi_{m,k}^S(f) \bullet \psi_{k',j}^T(f') = \delta_{k,k'} \psi_{m,j}^{S \otimes T} (\alpha_{m,S,T}(f \otimes 1_T) \alpha_{S,k,T}^{-1}(1_S \otimes f') \alpha_{S,T,j}) \tag{3.2}$$

for $f \in \mathcal{C}(S \otimes X_k, X_m \otimes S)$ and $f' \in \mathcal{C}(T \otimes X_j, X_{k'} \otimes T)$. This is a straightforward adaptation of the product formula in [GJ16], by putting the associator α in appropriate places so that the overall formula makes sense in the current nonstrict setting. Independence of the definition of \bullet on the choice of f and f' easily follows from the naturality of α .

The space $\mathcal{A}_{k,j}$ can be presented in a more categorical way as follows. Let $\iota \otimes X_j$ be the endofunctor of \mathcal{C} given by $Y \mapsto Y \otimes X_j$, and similarly consider $X_k \otimes \iota$. Denote by $\text{Nat}_{00}(\iota \otimes X_j, X_k \otimes \iota)$ the space of natural transformations $(f_Y: Y \otimes X_j \rightarrow X_k \otimes Y)_{Y \in \text{Obj}(\mathcal{C})}$ from the functor from $\iota \otimes X_j$ to $X_k \otimes \iota$, such that $f_X = 0$ for $X \in \mathcal{I}$ except for finitely many of them. Then, using semisimplicity of \mathcal{C} , one can check that the map

$$\text{Nat}_{00}(\iota \otimes X_j, X_k \otimes \iota) \rightarrow \mathcal{A}_{k,j}, \quad f \mapsto \sum_{S \in \mathcal{I}} \psi_{k,j}^S(f_S)$$

is a vector space isomorphism. Moreover, $\bigoplus_{j,k \in \mathcal{J}} \text{Nat}_{00}(\iota \otimes X_j, X_k \otimes \iota)$ can be equipped with a canonical algebra structure \bullet for which the above mentioned map becomes an algebra isomorphism. Concretely, for $f \in \text{Nat}_{00}(\iota \otimes X_k, X_m \otimes \iota)$ and $f' \in \text{Nat}_{00}(\iota \otimes X_j, X_k \otimes \iota)$, $f \bullet f'$ is characterized by the following: given $X \in \mathcal{I}$, the morphism $(f \bullet f')_X$ can be expressed as

$$(f \bullet f')_X = \sum_{\substack{Y,Z \in \mathcal{I}, \\ v_a: X \rightarrow Y \otimes Z}} (1_m \otimes v_a^*) \alpha_{m,Y,Z}(f_Y \otimes 1_Z) \alpha_{Y,k,Z}^{-1}(1_Y \otimes g_Z) \alpha_{Y,Z,j}(v_a \otimes 1_j),$$

where $v_a: X \rightarrow Y \otimes Z$ runs through an orthonormal basis of $\mathcal{C}(X, Y \otimes Z)$ for each Y and Z . It follows that, for $f \in \mathcal{C}(S \otimes X_k, X_m \otimes S)$ and $f' \in \mathcal{C}(T \otimes X_j, X_k \otimes T)$, their product in terms of ψ can be written as

$$\psi_{m,k}^S(f) \bullet \psi_{k,j}^T(f') = \sum_{U,w} \psi_{m,j}^U ((1_m \otimes w^*) \alpha_{m,S,T}(f \otimes 1_T) \alpha_{S,k,T}^{-1}(1_S \otimes f') \alpha_{S,T,j}(w \otimes 1_j)),$$

where U runs through \mathcal{I} and w through an orthonormal basis of $\mathcal{C}(U, S \otimes T)$.

3.2. LEMMA. (\mathcal{A}, \bullet) is an associative algebra.

PROOF. Take indices $j, k, m, n \in \mathcal{J}$, objects $S, T, U \in \text{Obj}(\mathcal{C})$, and morphisms

$$f \in \mathcal{C}(S \otimes X_m, X_n \otimes S), \quad f' \in \mathcal{C}(T \otimes X_k, X_m \otimes T), \quad f'' \in \mathcal{C}(U \otimes X_j, X_k \otimes U).$$

On the one hand, $(\psi_{n,m}^S(f) \bullet \psi_{m,k}^T(f')) \bullet \psi_{k,j}^U(f'')$ is the image under $\psi_{n,j}^{(S \otimes T) \otimes U}$ of

$$\alpha_{n,S \otimes T,U} \left((\alpha_{n,S,T}(f \otimes 1_T) \alpha_{S,m,T}^{-1} (1_S \otimes f') \alpha_{S,T,k}) \otimes 1_U \right) \alpha_{S \otimes T,k,U} (1_{S \otimes T} \otimes f'') \alpha_{S \otimes T,U,j}. \quad (3.3)$$

On the other, $\psi_{n,m}^S(f) \bullet (\psi_{m,k}^T(f') \bullet \psi_{k,j}^U(f''))$ is the image under $\psi_{n,j}^{S \otimes (T \otimes U)}$ of

$$\alpha_{n,S,T \otimes U} (f \otimes 1_{T \otimes U}) \alpha_{S,m,T \otimes U}^{-1} \left(1_S \otimes (\alpha_{m,T,U}(f' \otimes 1_U) \alpha_{T,k,U}^{-1} (1_T \otimes f'')) \alpha_{T,U,j} \right) \alpha_{S,T \otimes U,j}. \quad (3.4)$$

Using $\psi_{n,j}^{S \otimes (T \otimes U)}((1_n \otimes \alpha) f' (\alpha^{-1} \otimes 1_j)) = \psi_{n,j}^{(S \otimes T) \otimes U}(f')$ together with the naturality and the pentagon equation for α , we indeed obtain that (3.3) and (3.4) are equal.

Let us indicate the first step. In the expression (3.3), by the naturality of α we can insert $\alpha_{S,m \otimes T,U}^{-1}$ and $\alpha_{S,T \otimes k,U}$ around $(1_S \otimes f') \otimes 1_U$, and $\alpha_{S,T,k \otimes U}^{-1}$ and $\alpha_{S,T,U \otimes j}$ around $1_{S \otimes T} \otimes f''$. Then between $(1_S \otimes (f' \otimes 1_U))$ and $(1_S \otimes (1_T \otimes f''))$ we have

$$\alpha_{S,T \otimes k,U} (\alpha_{S,T,k} \otimes 1_U) \alpha_{S \otimes T,k,U}^{-1} \alpha_{S,T,k \otimes U}^{-1}: S \otimes (T \otimes (X_k \otimes U)) \rightarrow S \otimes ((T \otimes X_k) \otimes U),$$

which is equal to $1_S \otimes \alpha_{T,k,U}^{-1}$ by the pentagon identity. This is exactly what we have between in f' and f'' in (3.4). The rest of the proof proceeds in a similar way. \blacksquare

Let us next describe the $*$ -structure on \mathcal{A} . We will denote this one by $\#$ in order to avoid confusion with the involution of morphisms of \mathcal{C} . Define a conjugate linear map $\#: \mathcal{A} \rightarrow \mathcal{A}$ by sending $\psi_{k,j}^S(f) \in \mathcal{A}_{k,j}^S$ to

$$\psi_{j,k}^{\bar{S}} \left(\left(\left((R_S^* \otimes 1_j) \alpha_{\bar{S},S,j}^{-1} \right) \otimes 1_{\bar{S}} \right) \left((1_{\bar{S}} \otimes f^*) \otimes 1_{\bar{S}} \right) \alpha_{\bar{S},k \otimes S,\bar{S}}^{-1} \left(1_{\bar{S}} \otimes (\alpha_{k,S,\bar{S}}^{-1} (1_k \otimes \bar{R}_S)) \right) \right) \quad (3.5)$$

in $\mathcal{A}_{j,k}^{\bar{S}}$, where (R_S, \bar{R}_S) is a standard solution to conjugate equations for S . Note that by the naturality of α and the quotient map ψ , this becomes well-defined as well as independent of the choice of \bar{S} and (R_S, \bar{R}_S) . Moreover, $\#$ is indeed an antimultiplicative involution; to see this, one argues along the same lines as in the proof of Lemma 3.2, using the fact that $\{w_i^{\vee*}\}_i$ is an orthonormal basis of $\mathcal{C}(\bar{U}, \bar{T} \otimes \bar{S})$ if $\{w_i\}_i$ is a one of $\mathcal{C}(U, S \otimes T)$ with respect to a standard solution (R_U, \bar{R}_U) for U , (R_T, \bar{R}_T) for T , and the associated one

$$\left(\alpha_{\bar{T} \otimes \bar{S},S,T} (\alpha_{\bar{T},\bar{S},S}^{-1} \otimes 1_T) \left((1_{\bar{T}} \otimes R_S) \otimes 1_T \right) R_T, \alpha_{S \otimes T,\bar{T},\bar{S}} (\alpha_{S,T,\bar{T}}^{-1} \otimes 1_{\bar{S}}) \left((1_S \otimes \bar{R}_T) \otimes 1_{\bar{S}} \right) \bar{R}_S \right) \quad (3.6)$$

for $S \otimes T$.

3.3. PROPOSITION. Let $F: \mathcal{C} \rightarrow \mathcal{C}'$ be a unitary monoidal equivalence of rigid C^* -tensor categories with simple units. Furthermore let $\mathcal{X} = \{X_j\}_{j \in \mathcal{J}}$ be a full family in \mathcal{C} , and put $\mathcal{X}' = \{F(X_j)\}_{j \in \mathcal{J}}$. Then $\mathcal{A}(\mathcal{X})$ is isomorphic to $\mathcal{A}(\mathcal{X}')$.

PROOF. By definition F comes with a C^* -functor $\mathcal{C} \rightarrow \mathcal{C}'$ (again denoted by the same symbol F) and a natural unitary transformation $F_2: F(X) \otimes F(Y) \rightarrow F(X \otimes Y)$ satisfying compatibility conditions for the monoidal structures. We then have a linear map

$$\mathcal{C}(S \otimes X_j, X_k \otimes S) \rightarrow \mathcal{C}'(F(S) \otimes F(X_j), F(X_k) \otimes F(S)), \quad f \mapsto F_2^{-1}F(f)F_2$$

which induces a linear map $\mathcal{A}(\mathcal{X})_{k,j} \rightarrow \mathcal{A}(\mathcal{X}')_{k,j}$. A tedious but straightforward computation yields that this is indeed a $*$ -isomorphism. ■

3.4. DEFINITION. [cf. [Ocn94, Jon01, GJ16]] *The $*$ -algebra $(\mathcal{A}(\mathcal{X}), \bullet, \#)$ is called the annular algebra of \mathcal{C} associated with the family $\mathcal{X} = \{X_j\}_{j \in \mathcal{J}}$. When $\mathcal{X} = \text{Irr } \mathcal{C}$, a set of representatives of the simple objects, we call $\mathcal{T} = \mathcal{T}(\mathcal{C}) = \mathcal{A}(\mathcal{X})$ the tube algebra of \mathcal{C} .*

The results of [GJ16] apply to our construction by Proposition 3.3. Let us summarize some important consequences:

- any element of $\mathcal{A}(\mathcal{X})$ has a uniform bound on the norm under the $*$ -representations of \mathcal{A} ; hence \mathcal{A} admits a universal C^* -envelope.
- there is a canonical faithful positive trace on $\tau: \mathcal{A} \rightarrow \mathbb{C}$ given by

$$\mathcal{A}_{k,j}^S \ni \psi_{k,j}^S(f) \mapsto \delta_{j,k} \sum_w \text{Tr}_j((1_j \otimes w^*) f (w \otimes 1_j)),$$

where w runs through an orthonormal basis of $\mathcal{C}(\mathbb{1}, S)$. (We suppressed the structure morphisms for $\mathbb{1}$ as before.) Note that simplicity of $\mathbb{1}$ is crucially used here, and the definition of τ is indeed independent of the choice of the orthonormal basis.

- the subspaces $\mathcal{A}_{k,j}^S$ of $\mathcal{A}_{k,j}$ for different $S \in \mathcal{I}$ are orthogonal with respect to the inner product defined by $\langle a_1, a_2 \rangle = \tau(a_1 \bullet a_2^\#)$.
- if $\mathcal{X} = \{X_j\}_{j \in \mathcal{J}}$ and $\mathcal{Y} = \{Y_{j'}\}_{j' \in \mathcal{J}'}$ are two full families, the algebras $\mathcal{A}(\mathcal{X})$ and $\mathcal{A}(\mathcal{Y})$ are strongly Morita equivalent.

3.5. HALF-BRAIDING AND MONAD. When X is an object of a C^* -tensor category \mathcal{C} , a unitary half-braiding on X is a natural family of unitary morphisms $c_Y: Y \otimes X \rightarrow X \otimes Y$ for $Y \in \text{Obj}(\mathcal{C})$ such that

$$\alpha_{X,Y,Z}(c_Y \otimes 1_Z) \alpha_{Y,X,Z}^{-1}(1_Y \otimes c_Z) \alpha_{Y,Z,X} = c_{Y \otimes Z}. \tag{3.7}$$

A morphism of half-braidings from (X, c) to (X', c') is a morphism x from X to X' (in \mathcal{C}) which satisfies $(x \otimes 1_Y)c_Y = c'_Y(1_Y \otimes x)$ for all Y . The category $\mathcal{Z}(\mathcal{C})$ of the pairs (X, c) as above is the (unitary) *Drinfeld center*.

Recall that we can enlarge \mathcal{C} to the ind-category $\text{ind-}\mathcal{C}$ whose objects are *direct limits* $\varinjlim X_i$ for some inductive system $(X_i, v_{ji})_{i,j \in \Lambda}$ of objects labeled over a directed set Λ , with connecting *isometries* $v_{ji}: X_i \rightarrow X_j$ for $i < j$ [NY16]. For semisimple \mathcal{C} , with a representative of irreducible classes \mathcal{I} as before, such a limit can always be represented

as ‘infinite direct sum’ $\bigoplus_{S \in \mathcal{I}} H_S \otimes S$. Here $H_S \otimes S$ is an amplification of S by a Hilbert spaces H_S , and the morphism space of two such direct sums is concretely given by

$$\text{Mor}_{\text{ind-}\mathcal{C}}\left(\bigoplus_{S \in \mathcal{I}} H_S \otimes S, \bigoplus_{S \in \mathcal{I}} H'_S \otimes S\right) = \ell_\infty\text{-}\prod_{S \in \mathcal{I}} B(H_S, H'_S),$$

where the right hand side denotes the space of uniformly bounded sequences of operators with respect the operator norm.

Then the category $\text{ind-}\mathcal{C}$ is again a (non-rigid) C^* -tensor category, and we can again form the Drinfeld center category $\mathcal{Z}(\text{ind-}\mathcal{C})$, which is often the ‘correct’ model of quantum double of \mathcal{C} when the number of simple classes are infinite. As explained in [Izu00, PSV18], the category of $*$ -representations of the tube algebra $\mathcal{T}(\mathcal{C})$ is equivalent to $\mathcal{Z}(\text{ind-}\mathcal{C})$ as a C^* -category.

The structure of the annular algebra can be understood using the *monadic* formulation of the quantum double [Müg03, BV12]. Since there is a slight issue of encoding the relevant structures only using uniformly bounded families of morphisms, let us consider purely algebraic direct sums, so that we just present our construction on each direct summand. Given $X \in \text{Obj}(\text{ind-}\mathcal{C})$, consider $\mathcal{Z}(X) = \bigoplus_{S \in \mathcal{I}}^{\text{alg}} \bar{S} \otimes (X \otimes S)$ as a direct sum ‘without completion over the S ’. Then the monoidal structure of \mathcal{C} induces natural transformations $\mu: \mathcal{Z}^2 \rightarrow \mathcal{Z}$ and $\eta: \text{Id}_{\mathcal{C}} \rightarrow \mathcal{Z}$ satisfying certain set of conditions analogous to monoids [ML98]. To be specific, let us fix a standard solution (R_S, \bar{R}_S) for each $S \in \mathcal{I}$, and also an orthonormal basis $(w_i)_i$ of $\mathcal{C}(U, S \otimes T)$ for each $S, T, U \in \mathcal{I}$. Then the part of μ from the summand $\bar{T} \otimes ((\bar{S} \otimes (X \otimes S)) \otimes T)$ to $\bar{U} \otimes (X \otimes U)$ is given by

$$\sum_i (w_i^\vee \otimes (1_X \otimes w_i^*)) (1_{\bar{T} \otimes \bar{S}} \otimes \alpha_{X, S, T}) \alpha_{\bar{T}, \bar{S}, (X \otimes S) \otimes T}^{-1} (1_{\bar{T}} \otimes \alpha_{\bar{S}, X \otimes S, T}), \tag{3.8}$$

where w_i^\vee is calculated using standard solutions (R_T, \bar{R}_T) , (R_U, \bar{R}_U) , and (3.6).

An ind- object X with a (non-unitary) half braiding c admits a structure of \mathcal{Z} -module, that is, a morphism $\mathcal{Z}(X) \rightarrow X$ which is compatible with $\mu_X: \mathcal{Z}^2(X) \rightarrow \mathcal{Z}(X)$. Concretely, the module structure is given by the collection of

$$(1_X \otimes R_S^*) \alpha_{X \bar{S}, S} (c_{\bar{S}} \otimes 1_S) \alpha_{\bar{S}, X, S}^{-1}: \bar{S} \otimes (X \otimes S) \rightarrow X.$$

The (non-unitary) Drinfeld center can be identified with the category $\mathcal{Z}\text{-mod}$ of \mathcal{Z} -modules, which contain objects of the form $\mathcal{Z}(X)$ as ‘free’ \mathcal{Z} -modules. We have natural isomorphisms $\mathcal{A}_{k,j} \cong \mathcal{C}(X_j, \mathcal{Z}(X_k))$, and the usual adjunction

$$\mathcal{C}(X_j, \mathcal{Z}(X_k)) \cong \text{Mor}_{\mathcal{Z}\text{-mod}}(\mathcal{Z}(X_j), \mathcal{Z}(X_k))$$

induces a $*$ -algebra structure on \mathcal{A} from that of $\text{End}_{\mathcal{Z}\text{-mod}}(\bigoplus_{j \in \mathcal{J}} \mathcal{Z}(X_j))$, which is precisely the structure of tube algebra.

4. Twisting and annular algebras

Let \mathcal{C} be a Γ -graded rigid C^* -tensor category with simple unit. For simplicity, we will assume that it is strict. Given a normalized 3-cocycle ω on Γ , our goal is to describe the tube (and more generally annular) algebra of \mathcal{C}^ω in terms of that of \mathcal{C} . Let \mathcal{X} be a family of objects in \mathcal{C} . Since both \mathcal{C} and \mathcal{C}^ω are identical as semisimple C^* -categories, $\mathcal{A} = \mathcal{A}(\mathcal{X})$ admit two different $*$ -algebra structures. Our goal is to extend the computation of [BDGR17] about $\mathcal{C}_\Gamma^\omega$ to \mathcal{C}^ω when \mathcal{X} is compatible with the grading as we specify below. In order to avoid confusion we denote by \bullet and $\#$ (resp. by $\bar{\bullet}$ and \star) the multiplication and the involution on \mathcal{A} induced by \mathcal{C} (resp. by \mathcal{C}^ω).

4.1. FELL BUNDLE STRUCTURE OF THE TUBE ALGEBRA. As before, let \mathcal{I} be a set of representatives of isomorphism classes of simple objects. Let us denote the map $\mathcal{I} \rightarrow \Gamma$ corresponding to the Γ -grading by $S \mapsto \gamma_S$.

4.2. DEFINITION. *Let \mathcal{C} be a Γ -graded category. A family of objects $\mathcal{X} = \{X_j\}_{j \in \mathcal{J}}$ of \mathcal{C} is adapted to the grading when each X_j is isomorphic to an object of \mathcal{C}_{γ_j} for some $\gamma_j \in \Gamma$.*

4.3. EXAMPLE. For example, \mathcal{I} is adapted for a trivial reason. Another important example comes from the representation of a compact quantum group G . Let $\{U_i\}_i$ be a set of representatives of irreducible representations, and H_i be the underlying Hilbert space of U_i . Then the family $\mathcal{X} = \{\bar{H}_i \otimes U_i\}_i$ in $\text{Rep } G$ is adapted to the grading. The associated annular algebra $\mathcal{A}(\mathcal{X})$ is isomorphic to the Drinfeld double of G [NY15b].

We start with giving several convenient direct sum decompositions of the annular algebra $\mathcal{A} = \mathcal{A}(\mathcal{X})$ for an adapted family \mathcal{X} . Recall that we have direct sum decompositions

$$\mathcal{A} = \bigoplus_{j,k \in \mathcal{J}} \mathcal{A}_{k,j}, \quad \mathcal{A}_{k,j} = \bigoplus_{S \in \mathcal{I}} \mathcal{C}(S \otimes X_j, X_k \otimes S)$$

as vector spaces. Note that the direct summand $\mathcal{A}_{k,j}^S = \mathcal{C}(S \otimes X_j, X_k \otimes S)$ is nonzero only if both $S \otimes X_j$ and $X_k \otimes S$ have subobjects isomorphic to a common element in \mathcal{I} , which will then imply $\gamma_S \gamma_j = \gamma_k \gamma_S$. Thus, if $\mathcal{A}_{k,j}^S$ is nonzero then γ_j and γ_k are conjugate to each other implemented by $\gamma_S^{\pm 1}$. This leads us to the following constructions. For $\gamma, \eta, s \in \Gamma$ such that $s\gamma = \eta s$, set

$$\mathcal{A}_{\eta,\gamma}^s = \bigoplus_{\substack{j,k \in \mathcal{J}, S \in \mathcal{I} \\ \gamma_j = \gamma, \gamma_k = \eta, \\ \gamma_S = s}} \mathcal{A}_{k,j}^S, \quad \mathcal{A}_{\eta,\gamma} = \bigoplus_{t: t\gamma = \eta t} \mathcal{A}_{\eta,\gamma}^t$$

Finally, let Σ be the set of conjugacy classes of Γ , and define $\mathcal{A}_\sigma = \bigoplus_{\gamma, \eta \in \sigma} \mathcal{A}_{\eta,\gamma}$ for $\sigma \in \Sigma$.

4.4. LEMMA. *The $*$ -algebra $(\mathcal{A}, \bullet, \#)$ decompose into the direct sum $\bigoplus_{\sigma \in \Sigma} \mathcal{A}_\sigma$ of $*$ -subalgebras. Thus, every representation of the tube algebra is an orthogonal direct sum of representations of \mathcal{A}_σ for $\sigma \in \Sigma$.*

PROOF. By the above discussion we have $\mathcal{A} = \bigoplus_{\sigma \in \Sigma} \mathcal{A}_\sigma$ as a vector space. The multiplication formula 3.2 tells us that \mathcal{A}_σ is closed under \bullet , and elements from different conjugacy classes are orthogonal. More precisely $\mathcal{A}_{t\eta t^{-1}, \eta}^t \mathcal{A}_{s\gamma s^{-1}, \gamma}^s$ contains a nonzero element (if and) only if $\eta = s\gamma s^{-1}$, in which case it is in $\mathcal{A}_{ts\gamma s^{-1}t^{-1}, \gamma}^{ts}$. From the explicit formula, we can see that $\#$ sends elements of $\mathcal{A}_{s\gamma s^{-1}, \gamma}^s$ to $\mathcal{A}_{\gamma, s\gamma s^{-1}}^{s^{-1}}$, so \mathcal{A}_σ is stable under involution. \blacksquare

4.5. REMARK. In terms of the Drinfeld center, the above direct sum decomposition corresponds to the fact that any object (X, c) of $\mathcal{Z}(\text{ind-}\mathcal{C})$ decomposes as a direct sum $\bigoplus_{\sigma \in \Sigma} (X_\sigma, c^\sigma)$ for some objects $X_\sigma \in \text{ind-}\bigoplus_{\gamma \in \sigma} \mathcal{C}_\gamma$ and unitary half braidings c^σ on the X_σ . Indeed, we can take a decomposition $X \cong \bigoplus_{\gamma} X_\gamma$ as an object in $\text{ind-}\mathcal{C}$ and put $X_\sigma = \bigoplus_{\gamma \in \sigma} X_\gamma$. If Y is an object in \mathcal{C}_η , the morphism c_Y from $\bigoplus_{\sigma} Y \otimes X_\sigma$ to $\bigoplus_{\sigma} X_\sigma \otimes Y$ has to be diagonally represented in σ , since there are no nontrivial morphisms between objects of $\text{ind-}\mathcal{C}_{\eta\gamma}$ and of $\text{ind-}\mathcal{C}_{\gamma'\eta}$ if γ and γ' are not conjugate in Γ . Thus, each X_σ inherits a half braiding c^σ by restricting c .

Let $\mathcal{G} = \Gamma \ltimes_{\text{Ad}} \Gamma$ be the action groupoid of Γ acting on itself by the adjoint action. The above argument shows that \mathcal{A} is a Fell bundle over \mathcal{G} [Yam90, Kum98]: to each $g = (s, \gamma)$ we have the direct summand $\mathcal{A}_g = \mathcal{A}_{s\gamma s^{-1}, \gamma}^s$ such that the $*$ -algebra structure of $\mathcal{A} = \bigoplus_{g \in \mathcal{G}} \mathcal{A}_g$ is compatible with the groupoid structure of \mathcal{G} , in the sense that $\mathcal{A}_g \mathcal{A}_{g'} \subset \mathcal{A}_{gg'}$ for $(g, g') \in \mathcal{G}^{(2)}$ and $(\mathcal{A}_g)^\# = \mathcal{A}_{g^{-1}}$.

From now on, we will assume that the adapted family \mathcal{X} is full.

4.6. PROPOSITION. Let σ be a conjugacy class of Γ . For any $a \in \sigma$, $\mathcal{A}_{a,a}$ is strongly Morita equivalent to \mathcal{A}_σ via the completion of $\bigoplus_{b \in \sigma} \mathcal{A}_{a,b}$ inside \mathcal{A}_σ .

PROOF. Because of Proposition 3.3 we may assume that \mathcal{C} is strict. Let b be an element of σ , and $j \in \mathcal{J}$ be any index satisfying $\gamma_j = b$. The assertion holds if we can show that $\mathcal{A}_{b,a} \mathcal{A}_{a,b}$ contains the unit of $\mathcal{A}_{j,j}$. In fact, we claim that if S is any irreducible object such that $a\gamma_S = \gamma_S b$, then the unit of $\mathcal{A}_{j,j}$ is contained in the span of $\mathcal{A}_{j,k}^{\bar{S}} \mathcal{A}_{k,j}^S$ where k runs through the indices of \mathcal{J} satisfying $\gamma_k = \gamma_S \gamma_j \gamma_S^{-1}$.

Let $v: T \rightarrow S \otimes X_j \otimes \bar{S}$ be an isometry giving a irreducible subobject of $S \otimes X_j \otimes \bar{S}$, and put

$$\begin{aligned} f^{(v)} &= d(S)^{-\frac{1}{2}} (R_S^* \otimes 1_{j \otimes \bar{S}}) (1_{\bar{S}} \otimes v) \in \mathcal{C}(\bar{S} \otimes T, X_j \otimes \bar{S}), \\ g^{(v)} &= d(S)^{-\frac{1}{2}} (v^* \otimes 1_S) (1_{S \otimes j} \otimes R_S) \in \mathcal{C}(S \otimes X_j, T \otimes S), \end{aligned}$$

where R_S is a part of a standard solution to the conjugate equation of the dual pair (S, \bar{S}) . Then, by the product formula 3.2, we get

$$\psi_{j,T}^{\bar{S}}(f^{(v)}) \bullet \psi_{T,j}^S(g^{(v)}) = d(S)^{-1} \psi_{j,j}^{\bar{S} \otimes S} ((R_S^* \otimes 1_{j \otimes \bar{S} \otimes S}) (1_{\bar{S}} \otimes v v^* \otimes 1_S) (1_{\bar{S} \otimes S \otimes j} \otimes R_S))$$

Taking the summation on v over a maximal system of orthogonal embeddings of irreducible subobjects of $S \otimes X_j \otimes \bar{S}$, we obtain

$$\sum_v \psi_{j,T}^{\bar{S}}(f^{(v)}) \bullet \psi_{T,j}^S(g^{(v)}) = d(S)^{-1} \psi_{j,j}^{\bar{S} \otimes S} (R_S^* \otimes 1_j \otimes R_S) = \psi_{j,j}^1(1_j).$$

Since \mathcal{X} is full, each T , being irreducible, must be a subobject of some $X_k \in \mathcal{X}$. Again, \mathcal{X} is adapted and v is nonzero; so, indeed $\gamma_k = \gamma_S \gamma_j \gamma_S^{-1}$. \blacksquare

4.7. REMARK. The above proposition shows that this Fell bundle satisfies the assumption of [MW08], namely, for each $g = (s, \gamma)$ in \mathcal{G} , \mathcal{A}_g is an imprimitivity bimodule between $\mathcal{A}_{\text{dom}(g)} = \mathcal{A}_{\gamma, \gamma}^e$ and $\mathcal{A}_{\text{codom}(g)} = \mathcal{A}_{s\gamma s^{-1}, s\gamma s^{-1}}^e$. See [IKSW18] for further implications on the structure of the C^* -envelope of \mathcal{A} .

4.8. TUBE ALGEBRA OF TWISTED CATEGORIES. Continuing to denote $\mathcal{A} = \mathcal{A}(\mathcal{X})$ for a full and adapted family $\mathcal{X} = \{X_j\}_{j \in \mathcal{J}}$, consider $\psi_{k,j'}^S(f_1) \in \mathcal{A}_{k,j'}^S$ and $\psi_{j,i}^T(f_2) \in \mathcal{A}_{j,i}^T$ for some indices $i, j, j', k \in \mathcal{J}$ and irreducible objects $S, T \in \mathcal{I}$. From the comparison of (3.2) for \mathcal{C} and \mathcal{C}^ω , we have

$$\psi_{k,j'}^S(f_1) \bar{\bullet} \psi_{j,i}^T(f_2) = \delta_{j,j'} \omega(\gamma_k, \gamma_S, \gamma_T) \bar{\omega}(\gamma_S, \gamma_j, \gamma_T) \omega(\gamma_S, \gamma_T, \gamma_i) \psi_{k,j}^S(f_1) \bullet \psi_{j,i}^T(f_2).$$

Note that this equation is meaningful only when γ_S (resp. γ_T) conjugates $\gamma_{j'}$ to γ_k (resp. γ_i to γ_j); otherwise, both sides will become zero by the discussion preceding Lemma 4.4. In terms of the Γ -grading, we may rewrite the above equation as

$$\chi_1 \bar{\bullet} \chi_2 = \delta_{\gamma_2, g_2'} \omega(\gamma_3, s, t) \bar{\omega}(s, \gamma_2, t) \omega(s, t, \gamma_1) \chi_1 \bullet \chi_2 \tag{4.1}$$

for all $\chi_1 \in \mathcal{A}_{\gamma_3, g_2'}^s$ and $\chi_2 \in \mathcal{A}_{\gamma_2, \gamma_1}^t$.

Similarly, combining (3.5) for the category \mathcal{C}^ω with Proposition 2.8, we see that the $*$ -structures are related by

$$\chi^* = \bar{\omega}(s^{-1}, s, \gamma_1) \bar{\omega}(s^{-1}, \gamma_2 s, s^{-1}) \bar{\omega}(\gamma_2, s, s^{-1}) \omega(s^{-1}, s, s^{-1}) \chi^\# \tag{4.2}$$

for $\chi \in \mathcal{A}_{\gamma_2, \gamma_1}^s$.

4.9. PROPOSITION. [cf. [Wil08, Theorem 3]] *Let us regard Γ as a Γ -set by the adjoint action. The normalized equivariant cochain $\Psi \in \bar{C}_\Gamma^2(\Gamma; \mathbb{T})$ defined by*

$$\Psi[s, t](\gamma) = \omega(\gamma, s, t) \bar{\omega}(s, s^{-1} \gamma s, t) \omega(s, t, t^{-1} s^{-1} \gamma s t),$$

is a cocycle.

PROOF. By adapting (2.1) in the context of the adjoint action of Γ on $X = \Gamma$, we see that a normalized equivariant 2-cocycle in $\bar{C}_\Gamma^2(\Gamma; \mathbb{T})$ is a map $\Xi: \Gamma \times \Gamma \rightarrow \text{Map}(\Gamma, \mathbb{T})$ such that

$$\Xi[t, u](s^{-1} \gamma s) \overline{\Xi[st, u](\gamma) \Xi[s, tu](\gamma) \Xi[s, t](\gamma)} = 1.$$

Expanding the left hand side for $\Xi = \Psi$, we obtain

$$\begin{aligned} & \underbrace{\omega(s^{-1} \gamma s, t, u) \bar{\omega}(t, t^{-1} s^{-1} \gamma s t, u) \omega(t, u, u^{-1} t^{-1} s^{-1} \gamma s t u)}_{\text{-----}} \\ & \times \underbrace{\bar{\omega}(\gamma, st, u) \omega(st, (st)^{-1} \gamma s t, u) \bar{\omega}(st, u, u^{-1} (st)^{-1} \gamma s t u)}_{\text{-----}} \\ & \times \underbrace{\omega(\gamma, s, tu) \bar{\omega}(s, s^{-1} \gamma s, tu) \omega(s, tu, (tu)^{-1} s^{-1} \gamma s t u)}_{\text{-----}} \\ & \times \underbrace{\bar{\omega}(\gamma, s, t) \omega(s, s^{-1} \gamma s, t) \bar{\omega}(s, t, t^{-1} s^{-1} \gamma s t)}_{\text{-----}}. \end{aligned}$$

By the 3-cocycle relation, the terms with:

- dotted underline give $\omega(s, t, u)\bar{\omega}(\gamma s, t, u)$,
- dashed underline give $\omega(s, t, t^{-1}s^{-1}\gamma stu)\bar{\omega}(s, t, u)$,
- wave underline give $\omega(\gamma s, t, u)\bar{\omega}(s, s^{-1}\gamma st, u)$, and those with
- double underline give $\omega(s, s^{-1}\gamma st, u)\bar{\omega}(s, t, t^{-1}s^{-1}\gamma stu)$.

Multiplying these terms we indeed obtain 1. ■

The corresponding groupoid 2-cocycle on $\mathcal{G} = \Gamma \ltimes_{\text{Ad}} \Gamma$ (via Equation 2.2) is given by

$$\psi(g_1, g_2) = \Psi[s_1, s_2](s_1 g_1 s_1^{-1}) = \omega(s_1 g_1 s_1^{-1}, s_1, s_2)\bar{\omega}(s_1, g_1, s_2)\omega(s_1, s_2, g_2) \tag{4.3}$$

for $(g_1, g_2) \in \mathcal{G}^{(2)}$ with $g_i = (s_i, g_i)$. Thus, we can twist any Fell bundle A over \mathcal{G} by ψ to another Fell bundle $A_\psi = \langle a^{(\psi)} \mid a \in A_\gamma, \gamma \in \mathcal{G} \rangle$ by

$$a^{(\psi)}b^{(\psi)} = \psi(g_1, g_2)(ab)^{(\psi)} \quad ((g_1, g_2) \in \mathcal{G}^{(2)}, a \in A_{g_1}, b \in A_{g_2}).$$

Note however that ψ is normalized only in the sense that $\psi(g_1, g_2) = 1$ when either g_1 or g_2 is in $\mathcal{G}^{(0)}$. Using this, one can easily check that $a^{(\psi)*} = \bar{\psi}(g^{-1}, g)a^{*(\psi)}$ for $a \in A_g$ gives a $*$ -structure.

4.10. THEOREM. *The $*$ -algebra $(\mathcal{A}, \bar{\bullet}, \star)$ can be regarded as the 2-cocycle twist of the Fell bundle $(\mathcal{A}, \bullet, \#)$ by the 2-cocycle ψ on \mathcal{G} .*

PROOF. Take $g_i = (s_i, \gamma_i)$ ($i = 1, 2$) in \mathcal{G} such that $(g_1, g_2) \in \mathcal{G}^{(2)}$. The product map $\mathcal{A}_{g_1} \times \mathcal{A}_{g_2} \rightarrow \mathcal{A}_{g_1 g_2}$ in $(\mathcal{A}, \bullet)_\psi$ is different from \bullet by the factor of (4.3). Since $\mathcal{A}_{g_1} = \mathcal{A}_{s_1 \gamma_1 s_1^{-1}, \gamma_1}^{s_1}$ and $\mathcal{A}_{g_2} = \mathcal{A}_{\gamma_1, \gamma_2}^{s_2}$, this is indeed the factor in (4.1).

It remains to compare the involutions. If $g = (s, \gamma_1) \in \mathcal{G}$, the involution of $(\mathcal{A}, \bullet, \#)_\psi$ on the summand \mathcal{A}_g is different from $\#$ by the factor of

$$\bar{\psi}(g^{-1}, g) = \bar{\Psi}[s^{-1}, s](\gamma_1) = \bar{\omega}(\gamma_1, s^{-1}, s)\omega(s^{-1}, s\gamma_1 s^{-1}, s)\bar{\omega}(s^{-1}, s, \gamma_1).$$

On the other hand, \star on \mathcal{A}_g is different from $\#$ by the factor of

$$\bar{\omega}(s^{-1}, s, \gamma_1)\bar{\omega}(s^{-1}, s\gamma_1, s^{-1})\bar{\omega}(s\gamma_1 s^{-1}, s, s^{-1})\omega(s^{-1}, s, s^{-1}).$$

By the 3-cocycle condition of $\bar{\omega}$ on quadruple $(s^{-1}, s\gamma_1, s^{-1}, s)$, we have

$$\begin{aligned} \bar{\omega}(s\gamma_1, s^{-1}, s)\bar{\omega}(s^{-1}, s\gamma_1, s^{-1}) &= \bar{\omega}(\gamma_1, s^{-1}, s)\omega(s^{-1}, s\gamma_1 s^{-1}, s)\bar{\omega}(s^{-1}, s\gamma_1, e) \\ &= \bar{\omega}(\gamma_1, s^{-1}, s)\omega(s^{-1}, s\gamma_1 s^{-1}, s). \end{aligned}$$

Eliminating these terms and the common factors $\bar{\omega}(s^{-1}, s, \gamma_1)$ and $\bar{\omega}(s^{-1}, s\gamma_1, s^{-1})$, we are reduced to checking

$$\bar{\omega}(s\gamma_1 s^{-1}, s, s^{-1})\omega(s^{-1}, s, s^{-1}) = \bar{\omega}(s\gamma_1, s^{-1}, s).$$

This equality follows from the 3-cocycle condition (and normalization condition) of $\bar{\omega}$ on the quadruple $(s\gamma_1, s^{-1}, s, s^{-1})$. ■

Let us also present a normalized version.

4.11. LEMMA. *When ψ is a 2-cocycle on \mathcal{G} satisfying $\psi(g, \text{dom}(g)) = 1 = \psi(\text{codom}(g), g)$, we have $\psi(g, g^{-1}) = \psi(g^{-1}, g)$ for any $g \in \mathcal{G}$.*

PROOF. Using the 2-cocycle condition on $(g, g^{-1}, g) \in \mathcal{G}^{(3)}$, we obtain

$$\psi(g^{-1}, g)\bar{\psi}(\text{codom}(g), g)\psi(g, \text{dom}(g))\bar{\psi}(g, g^{-1}) = 1.$$

Since ψ is normalized, the middle two terms are trivial. ■

It follows that we can choose a function $\xi: \mathcal{G} \rightarrow \mathbb{T}$ satisfying

$$\psi(g, g^{-1}) = \xi(g)^2, \quad \xi(g) = \xi(g^{-1}), \quad \xi(x) = 1 \quad (x \in \mathcal{G}^{(0)}).$$

Then the cohomologous cocycle $\psi' = \psi \cdot \delta^1 \bar{\xi}$ satisfies

$$\begin{aligned} \psi'(g, g^{-1}) &= \psi(g, g^{-1})\bar{\xi}(g^{-1})\bar{\xi}(g)\xi(\text{codom}(g)) = 1, \\ \psi'(g, \text{dom}(g)) &= \psi(g, \text{dom}(g)) = 1 = \psi'(\text{codom}(g), g) \end{aligned}$$

for any $g \in \mathcal{G}$. Thus, ψ' is a normalized 2-cocycle in the stronger sense, and $(\mathcal{A}, \bar{\bullet}, \star)$ is isomorphic to the twisting of $(\mathcal{A}, \bullet, \#)$ by ψ' by the map $\chi \mapsto \xi(g)\chi$ on \mathcal{A}_g .

Up to strong Morita equivalence, we can pick up one element from each orbit of \mathcal{G} (conjugacy class of Γ) and look at the stabilizers. In our setting, this means that choosing $a \in \sigma$ for each $\sigma \in \Sigma$, and considering the centralizer subgroup $C_\Gamma(a) = \{\gamma \in \Gamma \mid a\gamma = \gamma a\}$ in Γ . Then ψ induces a 2-cocycle on $C_\Gamma(a)$,

$$\varphi_a(s, t) = \omega(a, s, t)\bar{\omega}(s, a, t)\omega(s, t, a),$$

see [Wit96; BDGR17, Lemma 2.1].

The Fell bundle structure on \mathcal{A} implies that the \ast -algebras $(\mathcal{A}_{a,a}, \bullet, \#)$ and $(\mathcal{A}_{a,a}, \bar{\bullet}, \star)$ are graded over $C_\Gamma(a)$.

4.12. COROLLARY. *Let a be an element of Γ . Then $(\mathcal{A}_{a,a}, \bar{\bullet}, \star)$ is isomorphic to the twist of the $C_\Gamma(a)$ -graded \ast -algebra $(\mathcal{A}_{a,a}, \bullet, \#)$ by the normalized 2-cocycle $\varphi'_a = \varphi_a \cdot d\bar{\xi}$.*

4.13. REMARK. Since the 3-cocycle ω of Γ is normalized, the 2-cocycle φ_e on $C_\Gamma(e) = \Gamma$ turns out to be constant function 1. Consequently, if we consider the tube algebras, the two \ast -algebra structures are identical on $\mathcal{T}_{e,e} = \mathcal{T}_{\{e\}}$. Note that $\mathcal{T}_{e,e}$ contains the fusion algebra $\mathcal{T}_{1,1}$ as a full corner by Proposition 4.6. Recall that analytic properties \mathcal{C} , such as, amenability, Haagerup and property (T), are defined using admissible representations of the fusion algebra, that is, those representations of the fusion algebra which can be extended to the whole tube algebra (see [PV15, GJ16]). Since $\mathcal{T}_{e,e}$ is a \ast -ideal in \mathcal{T} , any admissible representation of $\mathcal{T}_{1,1}$ can extend up to $\mathcal{T}_{e,e}$. So, \mathcal{C} is amenable, has the Haagerup property, or has property (T) if and only if \mathcal{C}^ω exhibits the corresponding properties.

4.14. **REMARK.** Let us describe the corresponding deformation of monad as in Section 3.5. Let \mathcal{Z}^ω denote the corresponding monad of the tensor category \mathcal{C}^ω . Suppose that X has degree γ , and $S, T, U \in \mathcal{I}$ respectively have degree s, t, u . Then, from Proposition 2.8 and formulas (3.8), (3.6), we see that the morphism $\mu_X^\omega: (\mathcal{Z}^\omega)^2(X) \rightarrow \mathcal{Z}^\omega(X)$ is given by the collection of morphisms from $\bar{T} \otimes ((\bar{S} \otimes (X \otimes S)) \otimes T)$ to $\bar{U} \otimes (X \otimes U)$ given by

$$\omega(\gamma, s, t)\bar{\omega}(t^{-1}, s^{-1}, \gamma st)\omega(s^{-1}, \gamma s, t)\bar{\omega}(t^{-1}s^{-1}, s, t)\omega(t^{-1}, s^{-1}, s)\mu_X$$

which appears only when U is isomorphic to a subobject of $S \otimes T$, hence in particular $u = st$. This factor can be interpreted as a 2-cochain $\tilde{\psi}(g_1, g_2)$ on \mathcal{G} by setting $g_1 = (s, s^{-1}\gamma s)$ and $g_2 = (t, (st)^{-1}\gamma st)$. As expected, it is cohomologous to ψ via the coboundary of $\xi(g) = \omega(s^{-1}, s, \gamma)$ for $g = (s, \gamma)$, hence is a 2-cocycle in particular.

5. Examples

5.1. **GRADING BY CYCLIC GROUPS.** Let Γ be a cyclic group. Since $H^3(\mathbb{Z}; \mathbb{T})$ is trivial, we concentrate on the case of finite cyclic group $\Gamma = \mathbb{Z}/n\mathbb{Z}$. The considerations in the previous section become rather simple. As Γ is commutative, the direct sum decomposition of Lemma 4.4 becomes $\mathcal{T}(\mathcal{C}) = \bigoplus_{a=0}^{n-1} \mathcal{T}_{\{a\}}(\mathcal{C})$. Moreover, as $H^2(\Gamma; \mathbb{T})$ is trivial, Corollary 4.12 implies that $\mathcal{T}_{\{a\}}(\mathcal{C}^\omega)$ is isomorphic to $\mathcal{T}_{\{a\}}(\mathcal{C})$.

5.2. **EXAMPLE.** The group $\mathbb{Z}/2\mathbb{Z}$ has essentially just one nontrivial 3-cocycle, $\omega(a, b, c) = (-1)^{abc}$. For the categories of Example 2.5 we have (by [NY15a]) $(\mathcal{C}_{\delta,+1}^{\text{TL}})^\omega = \mathcal{C}_{\delta,-1}^{\text{TL}} \cong \text{Rep SU}_q(2)$ for positive q satisfying $q + q^{-1} = \delta$. This explains why the computation of the spectrum of annular algebra for the category $\mathcal{C}_{\delta,+1}^{\text{TL}}$ in [GJ16] gives exactly the same answer as the computation for the Drinfeld double of $\text{SU}_q(2)$ with positive q in [Pus93].

By the discussion made above, there is a ‘natural’ equivalence of the Drinfeld centers $\mathcal{Z}(\text{ind-}\mathcal{C})$ and $\mathcal{Z}(\text{ind-}\mathcal{C}^\omega)$ as C^* -categories, compatible with the associated gradings by $\mathbb{Z}/n\mathbb{Z}$. While a general description of the monoidal structure in terms of the tube algebra (cf. [DGG14]) will give a description of $\mathcal{Z}(\text{ind-}\mathcal{C}^\omega)$ as a monoidal category, there is a more concrete description available in this setting, as follows.

It is well known that the 3-cocycles on $\mathbb{Z}/n\mathbb{Z}$ can be represented (up to coboundary) by

$$\omega^k(a, b, c) = \exp\left(2\pi\sqrt{-1}k\left(\left\lfloor\frac{a+b}{n}\right\rfloor - \left\lfloor\frac{a}{n}\right\rfloor - \left\lfloor\frac{b}{n}\right\rfloor\right)\frac{c}{n}\right) \quad (k = 0, 1, \dots, n-1). \tag{5.1}$$

Note that $\omega^k(a, b, c)$ is invariant under congruence modulo n for $a, b, c \in \mathbb{Z}$. We will work with this concrete formula of ω^k . Thus, the associated 2-cocycle is

$$\varphi_a^k(s, t) = \exp\left(2\pi\sqrt{-1}k\left(\left\lfloor\frac{s+t}{n}\right\rfloor - \left\lfloor\frac{s}{n}\right\rfloor - \left\lfloor\frac{t}{n}\right\rfloor\right)\frac{a}{n}\right).$$

Let us introduce a new monoidal structure $\otimes^{(k)}$ on $\mathcal{Z}(\text{ind-}\mathcal{C})$. When (X, c) and (X', c') are objects of $\mathcal{Z}(\text{ind-}\mathcal{C})$ such that $X \in \text{Obj}(\text{ind-}\mathcal{C}_a)$ and $X' \in \text{Obj}(\text{ind-}\mathcal{C}_b)$, we set

$$(c \otimes^{(k)} c')_Y = \bar{\omega}^{2k}(a, b, s)(c \otimes c')_Y$$

when $Y \in \text{Obj}(\mathcal{C}_s)$ and extend by linearity to the general case, and put $(X, c) \otimes^{(k)} (X', c') = (X \otimes X', c \otimes^{(k)} c')$. This still defines a structure of monoidal category (with the same associator as $\mathcal{Z}(\text{ind-}\mathcal{C})$) because ω is a 2-cocycle in the first two variables.

5.3. PROPOSITION. *Let \mathcal{C} be a C^* -tensor category graded by $\mathbb{Z}/n\mathbb{Z}$, and ω^k be the 3-cocycle as above. Then $\mathcal{Z}(\text{ind-}\mathcal{C}^{\omega^k})$ is unitarily monoidally equivalent to $(\mathcal{Z}(\text{ind-}\mathcal{C}), \otimes^{(k)})^{\omega^k}$.*

PROOF. Let X be an object of $\text{ind-}\mathcal{C}$, and $(c_Y: Y \otimes X \rightarrow X \otimes Y)_Y$ be a unitary half braiding on X (with respect to the structure of \mathcal{C}). We first define a half braiding \tilde{c} on X as an object of \mathcal{C}^ω . By Remark 4.5, it is enough to consider $(X, c) \in \text{Obj}(\mathcal{Z}(\text{ind-}\mathcal{C}))$ with homogeneous object $X \in \text{Obj}(\text{ind-}\mathcal{C}_a)$.

For $m \in \mathbb{Z}$, let $r(m)$ denote the unique integer $0 \leq r(m) < n$ such that $m \equiv r(m) \pmod n$; in other words, $r(m) = m - n \lfloor \frac{m}{n} \rfloor$. When Y is an object in \mathcal{C}_s , put

$$\tilde{c}_Y = \exp\left(-2\pi\sqrt{-1}k \frac{r(s)r(a)}{n^2}\right) c_Y: Y \otimes X \rightarrow X \otimes Y,$$

and extend it to general Y using direct sum decomposition into homogeneous components. We claim that (X, \tilde{c}) is a half braiding in \mathcal{C}^{ω^k} . In view of (3.7), this amounts to verifying

$$\begin{aligned} \omega^k(a, s, t) \exp\left(-2\pi\sqrt{-1}k \frac{r(s)r(a)}{n^2}\right) \bar{\omega}^k(s, a, t) \exp\left(-\frac{2\pi\sqrt{-1}}{n^2}r(t)r(a)\right) \omega^k(s, t, a) \\ = \exp\left(-2\pi\sqrt{-1}k \frac{r(s+t)r(a)}{n^2}\right). \end{aligned}$$

This is indeed the case, since the terms involving ω give $\omega^k(s, t, a)$, and we also have

$$r(s) + r(t) - r(s+t) = n \left(\left\lfloor \frac{s+t}{n} \right\rfloor - \left\lfloor \frac{s}{n} \right\rfloor - \left\lfloor \frac{t}{n} \right\rfloor \right). \tag{5.2}$$

This way we obtain a C^* -functor $F: \mathcal{Z}(\text{ind-}\mathcal{C}) \rightarrow \mathcal{Z}(\text{ind-}\mathcal{C}^{\omega^k})$ given by $F(X, c) = (X, \tilde{c})$. It is an equivalence of categories, since we can ‘untwist’ \mathcal{C}^{ω^k} by $\bar{\omega}^k$ to recover \mathcal{C} , and perform the same construction to produce an inverse of F .

It remains to show that F can be enriched to a C^* -tensor functor if we first replace the monoidal structure of $\mathcal{Z}(\text{ind-}\mathcal{C})$ by $\otimes^{(k)}$, and then twist it by ω^k . We claim that the natural unitary transformation $F_2: (X, \tilde{c}) \otimes (X', \tilde{c}') \rightarrow (X \otimes X', c \otimes^{(k)} c')$ is simply represented by $1_{X \otimes X'}$. Now, $1_{X \otimes X'}$ is a morphism of half braiding if and only if the diagram

$$\begin{array}{ccc} Y \otimes (X \otimes X') & \xrightarrow{\tilde{c} \otimes \tilde{c}'} & (X \otimes X') \otimes Y \\ \downarrow 1 & & \downarrow 1 \\ Y \otimes (X \otimes X') & \xrightarrow{c \otimes^{(k)} c'} & (X \otimes X') \otimes Y \end{array}$$

commutes. To check this, as before we can also assume $X \in \text{Obj}(\text{ind-}\mathcal{C}_a)$, $X' \in \text{Obj}(\text{ind-}\mathcal{C}_b)$, and $Y \in \text{Obj}(\mathcal{C}_s)$. Then, the top row picks up the factor of

$$\exp\left(-2\pi\sqrt{-1}k \frac{r(s)(r(a) + r(b))}{n^2}\right) \exp\left(-2\pi\sqrt{-1}k \left(\left\lfloor \frac{a+b}{n} \right\rfloor - \left\lfloor \frac{a}{n} \right\rfloor - \left\lfloor \frac{b}{n} \right\rfloor\right) \frac{s}{n}\right),$$

while the bottom row picks up

$$\exp\left(-4\pi\sqrt{-1}k \left(\left\lfloor \frac{a+b}{n} \right\rfloor - \left\lfloor \frac{a}{n} \right\rfloor - \left\lfloor \frac{b}{n} \right\rfloor\right) \frac{s}{n}\right) \exp\left(-2\pi\sqrt{-1}k \frac{r(s)(r(a+b))}{n^2}\right).$$

These are indeed equal by (5.2).

It remains to check the compatibility of $(F, 1_{\mathbb{1}}, F_2)$ with the associators of categories $(\mathcal{Z}(\text{ind-}\mathcal{C}), \otimes^{(k)})^{\omega^k}$ and $\mathcal{Z}(\text{ind-}\mathcal{C}^{\omega^k})$, but it is obvious from the definitions. \blacksquare

The structure $\otimes^{(k)}$ agrees with the original one when $n = 2$, or more generally if n divides $2k$ in the parametrization of (5.1). Moreover, the above proposition has an obvious parallel for $\mathcal{Z}(\mathcal{C}^{\omega^k})$, and also for nonunitary variants.

Suppose that \mathcal{C} is (resp. unitarily) braided, or equivalently, that there is a (resp. \mathbb{C}^* -)tensor functor $F: \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C})$ of the form $X \mapsto (X, c)$. If n divides $2k$, the above result implies that $X \mapsto (X, \tilde{c})$ is a (resp. \mathbb{C}^* -)tensor functor from \mathcal{C}^{ω^k} to $\mathcal{Z}(\mathcal{C}^{\omega^k})$, hence \mathcal{C}^{ω^k} is again (resp. unitarily) braided. If n does not divide $2k$ the situation looks more complicated, but at least we can say that \mathcal{C}^{ω^k} is not braided when \mathcal{C} is either $\mathcal{C}_{\mathbb{Z}/n\mathbb{Z}}$ or $\text{RepSU}_q(n)$ as discussed below, cf. [NY15a, Remark 4.4].

5.4. EXAMPLE. Let us describe the case of $\text{RepSU}_q(n)$ for the root of unity $q = e^{\frac{\pi\sqrt{-1}}{m}}$, with $m \geq n - 1$. Then it makes sense as a rigid \mathbb{C}^* -tensor category \mathcal{C} (see, e.g., [Wen88]) which we assume to be strict. Its irreducible classes are parametrized by the dominant integral weights λ of \mathfrak{sl}_n such that $(\lambda + \rho, \alpha_{\max}) < m$, where ρ is the half sum of positive roots, α_{\max} is the highest root, and (λ, μ) is the invariant inner product on the weight space normalized so that each root α satisfies $(\alpha, \alpha) = 2$. If we denote the fundamental weights by $\omega_1, \dots, \omega_{n-1}$, the above condition on λ is equivalent to $\lambda = \sum_i \nu_i \omega_i$ with $\nu_i \geq 0$ and $\sum_i \nu_i \leq m - n$. Then \mathcal{C} is generated by the object X_1 (the ‘defining representation’) corresponding to $\lambda = \omega_1$, and $X_1^{\otimes n}$ contains $\mathbb{1}$ with multiplicity 1. It follows that the universal grading group of \mathcal{C} is $\mathbb{Z}/n\mathbb{Z}$ such that X_1 belongs to the homogeneous component \mathcal{C}_1 . Moreover, the classes U^i ($i = 1, 2, \dots, n - 1$) corresponding to $\lambda = (m - n)\omega_i$ exhaust the nontrivial invertible classes of \mathcal{C} , and the class of U^1 satisfies $(U^1)^{\otimes k} \cong U^k$, $(U^1)^{\otimes n} \cong \mathbb{1}$ [Bru00]. Now, since \mathcal{C} is a *modular tensor category*, its Drinfeld center $\mathcal{Z}(\mathcal{C})$ is equivalent to the Deligne product $\mathcal{C} \boxtimes \mathcal{C}^{\beta\text{op}}$ as a braided category, where βop denotes taking the opposite braiding [Müg03]. Indeed, the equivalence $F: \mathcal{C} \boxtimes \mathcal{C}^{\beta\text{op}} \rightarrow \mathcal{Z}(\mathcal{C})$ is given by $F(X \boxtimes X') = (X \otimes X', c)$, where $c = (c_Y)_Y$ is the half braiding on $X \otimes X'$ given by $c_Y = (1_X \otimes \beta_{X',Y}^{-1})(\beta_{Y,X} \otimes 1_{X'})$, with $\beta_{X,Y}: X \otimes Y \rightarrow Y \otimes X$ denoting the braiding of \mathcal{C} . In this picture the half braidings on the distinguished object X_1 are represented by $(X_1 \otimes U^i) \boxtimes U^{n-i}$ and $U^{n-i} \boxtimes (X_1 \otimes U^i)$ with the convention $U^0 = U^n = \mathbb{1}$. In the notation of Proposition 5.3, the n -th tensor power of $(X_1 \otimes U^i) \boxtimes U^{n-i}$ with respect to $\otimes^{(k)}$

is the product of $F(X_1^{\otimes n} \boxtimes \mathbb{1})$ with the half braiding on $\mathbb{1}$ given by $\exp(-4\pi\sqrt{-1}ks/n)\iota_Y$ for $Y \in \text{Obj}(\mathcal{C}_s)$ up to the natural identification of $\mathbb{1} \otimes Y$ and $Y \otimes \mathbb{1}$ with Y . Thus, it contains the trivial half braiding if and only if n divides $2k$, and the same can be said for $U^{n-i} \boxtimes (X_1 \otimes U^i)$. In view of Proposition 5.3, there is a (\mathbb{C}^*) -tensor functor from \mathcal{C}^{ω^k} to $\mathcal{Z}(\mathcal{C}^{\omega^k})$ which is identity on objects, or equivalently \mathcal{C}^{ω^k} is (unitarily) braided, if and only if n divides $2k$.

5.5. FREE PRODUCT. For $\varepsilon \in \{+, -\}$, let \mathcal{C}_ε be rigid semisimple \mathbb{C}^* -tensor categories with simple units $\mathbb{1}_\varepsilon$ respectively. Let \mathcal{I}_ε be a set of representatives of isomorphism classes of simple objects in \mathcal{C}_ε , and Γ_ε be the universal grading group. Consider the free product category $\mathcal{C} = \mathcal{C}_+ * \mathcal{C}_-$. Then the universal grading group Γ of \mathcal{C} is isomorphic to the group free product $\Gamma_+ * \Gamma_-$. Indeed, if \mathcal{W} is the set of words (including the empty one) whose letters alternately belong to $\mathcal{I}_+ \setminus \{\mathbb{1}_+\}$ and $\mathcal{I}_- \setminus \{\mathbb{1}_-\}$, then \mathcal{C} has a set of representatives of simple objects enumerated by \mathcal{W} , say $\mathcal{I} = \{X_w\}_{w \in \mathcal{W}}$ such that:

- (i) $X_\emptyset = \mathbb{1}$,
- (ii) $X_{w_1} \otimes X_{w_2} \cong X_{(w_1, w_2)}$ whenever the last letter of w_1 and the first of w_2 have opposite signs (that is, if one is in \mathcal{I}_+ , then other is in \mathcal{I}_-),
- (iii) if $w_1 = (w'_2, Y), w_2 = (Z, w'_2) \in \mathcal{W}$ such that Y and Z have the same sign $\varepsilon = \pm$, then the direct sum decomposition of $X_{w_1} \otimes X_{w_2}$ into simple objects can be obtained inductively on the lengths of w_1 and w_2 by the following rule:

$$(a) \quad X_{w_1} \otimes X_{w_2} \cong \bigoplus_{U \in \mathcal{I}_\varepsilon} (X_{(w'_1, U, w'_2)})^{\oplus \dim(\mathcal{C}_\varepsilon(U, Y \otimes Z))} \quad \text{when } Y \not\cong \bar{Z},$$

$$(b) \quad X_{w_1} \otimes X_{w_2} \cong (X_{w'_1} \otimes X_{w'_2}) \oplus \left(\bigoplus_{U \in \mathcal{I}_\varepsilon \setminus \{\mathbb{1}\}} (X_{(w'_1, U, w'_2)})^{\oplus \dim(\mathcal{C}_\varepsilon(U, Y \otimes Z))} \right) \quad \text{when } Y \cong \bar{Z}.$$

Using (i), (ii), and (iii), it is completely routine to check that the map

$$\Gamma \ni [X_w] \mapsto Y_1 \cdots Y_n \in \Gamma_+ * \Gamma_- \quad \text{where } w = (Y_1, \dots, Y_n)$$

is an isomorphism.

The cohomology of free product group is given by $H^3(\Gamma_+ * \Gamma_-; \mathbb{T}) \cong H^3(\Gamma_+; \mathbb{T}) \times H^3(\Gamma_-; \mathbb{T})$ (see, e.g., [Wei94, Section 6.2]). This allows us to decompose the twisting procedure to the cases when ω on $\Gamma_+ * \Gamma_-$ is induced from a 3-cocycle ω_0 on Γ_ε through the canonical quotient map $\Gamma_+ * \Gamma_- \rightarrow \Gamma_\varepsilon$. This induces a groupoid homomorphism (a functor) $\mathcal{G}_{\Gamma_+ * \Gamma_-} \rightarrow \mathcal{G}_{\Gamma_\varepsilon}$, and the associated map $H^2(\mathcal{G}_{\Gamma_\varepsilon}; \mathbb{T}) \rightarrow H^2(\mathcal{G}_{\Gamma_+ * \Gamma_-}; \mathbb{T})$. If Ψ_0 is the 2-cocycle on $\mathcal{G}_{\Gamma_\varepsilon}$ associated with ω_0 , its pullback Ψ on $\mathcal{G}_{\Gamma_+ * \Gamma_-}$ is the one associated with ω .

5.6. DIRECT PRODUCT. Next let us consider grading by direct product groups. One source of such a structure is the Deligne product $\mathcal{C}_1 \boxtimes \mathcal{C}_2$ of \mathcal{C}_1 and \mathcal{C}_2 . In this case we have $\text{Ch}(\mathcal{C}_1 \boxtimes \mathcal{C}_2) = \text{Ch}(\mathcal{C}_1) \times \text{Ch}(\mathcal{C}_2)$.

Suppose moreover that the grading is by a finite commutative group. The \mathbb{T} -valued 3-cocycles on such groups are concretely characterized in [NY15a, Proposition A.3], as follows. Write $\Gamma = (\mathbb{Z}/n_1\mathbb{Z}) \times \cdots \times (\mathbb{Z}/n_k\mathbb{Z})$ for $n_i \in \mathbb{N}$. Then a set of generators on $H^3(\Gamma; \mathbb{T})$ is given by:

- (i) those of the form (5.1) (with $k = 1$) for some factor $\mathbb{Z}/n_i\mathbb{Z}$,
- (ii) for distinct indices i, j :

$$\phi_{ij}(a, b, c) = \exp\left(2\pi\sqrt{-1} \left(\left\lfloor \frac{a_i + b_i}{n_i} \right\rfloor - \left\lfloor \frac{a_i}{n_i} \right\rfloor - \left\lfloor \frac{b_i}{n_i} \right\rfloor \right) \frac{c_j}{n_j}\right),$$

- (iii) for distinct indices i, j, k :

$$\phi_{ijk}(a, b, c) = \exp\left(\frac{2\pi\sqrt{-1}}{\text{gcd}(n_i, n_j, n_k)} a_i b_j c_k\right).$$

5.7. EXAMPLE. Consider the case $\Gamma = (\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})$, and let $a = (0, 0, 1)$. Then, the 3-cocycle ϕ_{123} gives $\varphi_a(s, t) = \exp(\pi\sqrt{-1}s_1t_2)$, or in a normalized form

$$\varphi'_a(s, t) = \exp\left(\frac{\pi\sqrt{-1}}{2}(s_1t_2 - s_2t_1)\right).$$

The tube algebra of \mathcal{C}_Γ is a commutative algebra of dimension 2^6 , while that of the twisted category $\mathcal{C}_\Gamma^{\phi_{123}}$ has a noncommutative direct summand $\mathcal{T}_{a,a}$. In particular, the number of irreducible classes in $\mathcal{Z}(\mathcal{C}_\Gamma^{\phi_{123}})$ is smaller than that of $\mathcal{Z}(\mathcal{C}_\Gamma)$.

References

- [ACM04] D. Altschuler, A. Coste, and J.-M. Maillard, *Representation theory of twisted group double*, Ann. Fond. Louis de Broglie **29** (2004), no. 4, 681–694, [arXiv:hep-th/0309257](#) [hep-th]. MR2146291
- [Bán90] P. Bántay, *Orbifolds and Hopf algebras*, Phys. Lett. B **245** (1990), no. 3-4, 477–479. MR1070067
- [BDGR17] D. Bisch, P. Das, S. K. Ghosh, and N. Rakshit, *Tube algebra of group-type subfactors*, Internat. J. Math. **28** (2017), no. 10, 1750069, 32, DOI:10.1142/S0129167X17500690, [arXiv:1701.00097](#) [math.OA]. MR3699167
- [Bru00] A. Bruguières, *Catégories prémodulaires, modularisations et invariants des variétés de dimension 3*, Math. Ann. **316** (2000), no. 2, 215–236, DOI:10.1007/s002080050011. MR1741269 (2001d:18009)

- [BV12] A. Bruguières and A. Virelizier, *Quantum double of Hopf monads and categorical centers*, Trans. Amer. Math. Soc. **364** (2012), no. 3, 1225–1279, DOI:10.1090/S0002-9947-2011-05342-0. MR2869176
- [DGG14] P. Das, S. K. Ghosh, and V. P. Gupta, *Drinfeld center of planar algebra*, Internat. J. Math. **25** (2014), no. 8, 1450076 (43 pages), DOI:10.1142/S0129167X14500761. MR3254423
- [GJ16] S. K. Ghosh and C. Jones, *Annular representation theory for rigid C^* -tensor categories*, J. Funct. Anal. **270** (2016), no. 4, 1537–1584, DOI:10.1016/j.jfa.2015.08.017, arXiv:1502.06543 [math.OA]. MR3447719
- [IKSW18] M. Ionescu, A. Kumjian, A. Sims, and D. P. Williams, *A stabilization theorem for Fell bundles over groupoids*, Proc. Roy. Soc. Edinburgh Sect. A **148** (2018), no. 1, 79–100, DOI:10.1017/S0308210517000129, MR3749336 [math.OA]. MR3749336
- [Izu00] M. Izumi, *The structure of sectors associated with Longo-Rehren inclusions. I. General theory*, Comm. Math. Phys. **213** (2000), no. 1, 127–179, DOI:10.1007/s002200000234. MR1782145 (2002k:46160)
- [Jon01] V. F. R. Jones, *The annular structure of subfactors*, Essays on geometry and related topics, Vol. 1, 2, 2001, pp. 401–463, Enseignement Math., Geneva, arXiv:math/0105071 [math.OA]. MR1929335 (2003j:46094)
- [KW93] D. Kazhdan and H. Wenzl, *Reconstructing monoidal categories*, I. M. Gel'fand Seminar, 1993, pp. 111–136, Amer. Math. Soc., Providence, RI. MR1237835 (95e:18007)
- [Kum98] A. Kumjian, *Fell bundles over groupoids*, Proc. Amer. Math. Soc. **126** (1998), no. 4, 1115–1125, DOI:10.1090/S0002-9939-98-04240-3, arXiv:math/9607230 [math.OA]. MR1443836 (98i:46055)
- [LR95] R. Longo and K.-H. Rehren, *Nets of subfactors*, Rev. Math. Phys. **7** (1995), no. 4, 567–597, DOI:10.1142/S0129055X95000232, arXiv:hep-th/9411077 [hep-th]. Workshop on Algebraic Quantum Field Theory and Jones Theory (Berlin, 1994). MR1332979 (96g:81151)
- [ML98] S. Mac Lane. (1998). *Categories for the working mathematician*, Second, Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, ISBN 0-387-98403-8. MR1712872 (2001j:18001)
- [Müg03] M. Müger, *From subfactors to categories and topology. II. The quantum double of tensor categories and subfactors*, J. Pure Appl. Algebra **180** (2003), no. 1-2, 159–219, DOI:10.1016/S0022-4049(02)00248-7, arXiv:math/0111205 [math.CT]. MR1966525 (2004f:18014)
- [MW08] P. S. Muhly and D. P. Williams, *Equivalence and disintegration theorems for Fell bundles and their C^* -algebras*, Dissertationes Math. (Rozprawy Mat.) **456** (2008), 1–57, DOI:10.4064/dm456-0-1, arXiv:0806.1022 [math.OA]. MR2446021 (2010b:46146)
- [NT13] S. Neshveyev and L. Tuset. (2013). *Compact quantum groups and their representation categories*, Cours Spécialisés [Specialized Courses], vol. 20, Société Mathématique de France, Paris, ISBN 978-2-85629-777-3. MR3204665
- [NY15a] S. Neshveyev and M. Yamashita, *Twisting the q -deformations of compact semisimple Lie groups*, J. Math. Soc. Japan **67** (2015), no. 2, 637–662, DOI:10.2969/jmsj/06720637, arXiv:1305.6949 [math.OA]. MR3340190
- [NY16] S. Neshveyev and M. Yamashita, *Drinfeld Center and Representation Theory for Monoidal Categories*, Comm. Math. Phys. **345** (2016), no. 1, 385–434, DOI:10.1007/s00220-016-2642-7, arXiv:1501.07390 [math.OA]. MR3509018

- [NY15b] S. Neshveyev and M. Yamashita, *A few remarks on the tube algebra of a monoidal category*, preprint (2015), available at [arXiv:1511.06332](https://arxiv.org/abs/1511.06332) [math.OA], to appear in Proc. Edinb. Math. Soc.
- [Ocn94] A. Ocneanu, *Chirality for operator algebras*, Subfactors (Kyuzeso, 1993), 1994, pp. 39–63, World Sci. Publ., River Edge, NJ. MR1317353 (96e:46082)
- [PSV18] S. Popa, D. Shlyakhtenko, and S. Vaes, *Cohomology and L^2 -Betti numbers for subfactors and quasi-regular inclusions*, Internat. Math. Res. Notices **2018** (2018), no. 8, 2241–2331, DOI:10.1093/imrn/rnw304, [arXiv:1511.07329](https://arxiv.org/abs/1511.07329) [math.OA].
- [PV15] S. Popa and S. Vaes, *Representation theory for subfactors, λ -lattices and C^* -tensor categories*, Comm. Math. Phys. **340** (2015), no. 3, 1239–1280, DOI:10.1007/s00220-015-2442-5, [arXiv:1412.2732](https://arxiv.org/abs/1412.2732) [math.OA]. MR3406647
- [Pus93] W. Pusz, *Irreducible unitary representations of quantum Lorentz group*, Comm. Math. Phys. **152** (1993), no. 3, 591–626, DOI:10.1007/BF02096620. MR1213303 (94c:81085)
- [Tur10] V. Turaev. (2010). *Homotopy quantum field theory*, EMS Tracts in Mathematics, vol. 10, European Mathematical Society (EMS), Zürich, DOI:10.4171/086, ISBN 978-3-03719-086-9. Appendix 5 by Michael Müger and Appendices 6 and 7 by Alexis Virelizier. MR2674592
- [Wei94] C. A. Weibel. (1994). *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, ISBN 0-521-43500-5; 0-521-55987-1. MR1269324
- [Wen88] H. Wenzl, *Hecke algebras of type A_n and subfactors*, Invent. Math. **92** (1988), no. 2, 349–383, DOI:10.1007/BF01404457. MR936086 (90b:46118)
- [Wes69] J. J. Westman, *Cohomology for ergodic groupoids*, Trans. Amer. Math. Soc. **146** (1969), 465–471. MR0255771
- [Wil08] S. Willerton, *The twisted Drinfeld double of a finite group via gerbes and finite groupoids*, Algebr. Geom. Topol. **8** (2008), no. 3, 1419–1457, [arXiv:math/0503266](https://arxiv.org/abs/math/0503266) [math.QA]. MR2443249
- [Wit96] S. J. Witherspoon, *The representation ring of the twisted quantum double of a finite group*, Canad. J. Math. **48** (1996), no. 6, 1324–1338. MR1426907
- [Wor87] S. L. Woronowicz, *Twisted $SU(2)$ group. An example of a noncommutative differential calculus*, Publ. Res. Inst. Math. Sci. **23** (1987), no. 1, 117–181, DOI:10.2977/prims/1195176848. MR890482 (88h:46130)
- [Yam90] S. Yamagami, *On primitive ideal spaces of C^* -algebras over certain locally compact groupoids*, Mappings of operator algebras (Philadelphia, PA, 1988), 1990, pp. 199–204, Birkhäuser Boston, Boston, MA. MR1103378 (92j:46110)

*DStat-Math Unit, Indian Statistical Institute
203 Barrackpore Trunk Road, Kolkata 700 108, India*

*Department of Mathematics, Ochanomizu University
Otsuka 2-1-1, Bunkyo, Tokyo 112-8610, Japan*

*current address: Department of Mathematics, University of Oslo
P.O box 1053, 0316 Oslo, Norway*

Email: jyotishmanbmath@gmail.com
shamindra.isi@gmail.com
narayan753@gmail.com
yamashita.makoto@ocha.ac.jp

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

INFORMATION FOR AUTHORS L^AT_EX₂ ϵ is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

T_EXNICAL EDITOR. Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT T_EX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Julie Bergner, University of Virginia: jeb2md@virginia.edu

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Gabriella Böhm, Wigner Research Centre for Physics: bohm.gabriella@wigner.mta.hu

Valeria de Paiva, Nuance Communications Inc: valeria.depaiva@gmail.com

Richard Garner, Macquarie University: richard.garner@mq.edu.au

Ezra Getzler, Northwestern University: getzler@northwestern.edu

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Dirk Hoffman, Universidade de Aveiro: dirk@ua.pt

Pieter Hofstra, Université d' Ottawa: phofstra@uottawa.ca

Anders Kock, University of Aarhus: kock@math.au.dk

Joachim Kock, Universitat Autònoma de Barcelona: kock@mat.uab.cat

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: matias.menni@gmail.com

Ieke Moerdijk, Utrecht University: i.moerdijk@uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Kate Ponto, University of Kentucky: kate.ponto@uky.edu

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: ross.street@mq.edu.au

Tim van der Linden, Université catholique de Louvain: tim.vanderlinden@uclouvain.be

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca