

$\overline{\text{Spec } \mathbb{Z}}$ AND THE GROMOV NORM

ALAIN CONNES AND CATERINA CONSANI

ABSTRACT. We define the homology of a simplicial set with coefficients in a Segal’s Γ -set (\mathfrak{s} -module). We show the relevance of this new homology with values in \mathfrak{s} -modules by proving that taking as coefficients the \mathfrak{s} -modules at the archimedean place over the structure sheaf on $\overline{\text{Spec } \mathbb{Z}}$ as in [2], one obtains on the singular homology with real coefficients of a topological space X , a norm equivalent to the Gromov norm. Moreover, we prove that the two norms agree when X is an oriented compact Riemann surface.

Contents

1	Introduction	155
2	Homology of a simplicial set with coefficients in an \mathfrak{s} -module	159
3	The archimedean place and the Gromov norm	170
4	Equality for Riemann surfaces of genus $g > 1$	174

1. Introduction

The notion of a Γ -set (due to G. Segal [12]) is a fundamental constituent in mathematics: it is the most embracing generalization of the datum given on a set by a commutative addition with a zero element and it provides a common framework for many of the present efforts to understand the “field with one element”. In [2] we defined on the Arakelov compactification $\overline{\text{Spec } \mathbb{Z}}$ of the algebraic spectrum of the integers a structure sheaf of Γ -rings which agrees with the classical structure sheaf when restricted to $\text{Spec } \mathbb{Z}$, but whose stalk at the archimedean place uses in a crucial way the new freedom of moving from the category of abelian groups to that of Γ -sets. To define Γ -sets one first introduces the small, full subcategory Γ^{op} of the category \mathfrak{Fin}_* of pointed finite sets, whose objects are pointed sets $k_+ := \{0, \dots, k\}$, for each integer $k \geq 0$ (0 is the base point) and with morphism the sets $\Gamma^{\text{op}}(k_+, m_+) = \{f : \{0, 1, \dots, k\} \rightarrow \{0, 1, \dots, m\} \mid f(0) = 0\}$. A Γ -set is then defined as a (covariant) functor $\Gamma^{\text{op}} \rightarrow \mathfrak{Sets}_*$ between pointed categories and the morphisms in

The authors would like to thank the referee for several useful remarks. The second author is partially supported by the Simons Foundation collaboration grant n. 353677.

Received by the editors 2019-10-02 and, in final form, 2020-02-11.

Transmitted by Ieke Moerdijk. Published on 2020-02-18.

2010 Mathematics Subject Classification: 16Y60; 20N20; 18G55; 18G30; 18G35; 18G55; 18G60; 14G40.

Key words and phrases: Gamma spaces, Gamma rings, Site, Gromov norm, Arakelov geometry, Homology theory.

© Alain Connes and Caterina Consani, 2020. Permission to copy for private use granted.

this category are natural transformations. The closed structure of the category $\Gamma\mathfrak{Sets}_*$ of Γ -sets is defined by setting

$$\underline{\Gamma\mathfrak{Sets}}_*(M, N) = \{k_+ \mapsto \Gamma\mathfrak{Sets}_*(M, N(k_+ \wedge -))\}, \quad (1)$$

where \wedge is the smash product of pointed sets. This formula uniquely defines the smash product of Γ -sets by applying the adjunction

$$\underline{\Gamma\mathfrak{Sets}}_*(M_1 \wedge M_2, N) = \underline{\Gamma\mathfrak{Sets}}_*(M_1, \underline{\Gamma\mathfrak{Sets}}_*(M_2, N)).$$

The notions of rings and modules then acquire a meaning in this symmetric monoidal closed category. In particular, Γ -sets can equivalently be viewed as modules over the simplest Γ -ring $\mathfrak{s} : \Gamma^{\text{op}} \rightarrow \mathfrak{Sets}_*$ whose underlying Γ -set is the identity functor, whence the name \mathfrak{s} -module to denote a Γ -set, and the more suggestive notation for morphisms in $\Gamma\mathfrak{Sets}_*$

$$\text{Hom}_{\mathfrak{s}}(M, N) := \Gamma\mathfrak{Sets}_*(M, N), \quad \underline{\text{Hom}}_{\mathfrak{s}}(M, N) := \underline{\Gamma\mathfrak{Sets}}_*(M, N).$$

Abelian groups form a full subcategory of the category $\mathfrak{s} - \text{Mod}$ of \mathfrak{s} -modules: the inclusion functor associates to an abelian group A the functor (Eilenberg-Mac Lane object) $HA : \Gamma^{\text{op}} \rightarrow \mathfrak{Sets}_*$ which assigns to a finite pointed set X the pointed set of A -valued maps on X vanishing at the base point of X ([3], 2.1.2). Note that the existence of an additive inverse is not used in the construction, and HM continues to make sense when M is a commutative monoid with a zero element.

At the conceptual level, it is important to make as explicit as possible the link between the category $\mathfrak{s} - \text{Mod}$ and the naive interpretation of vector spaces over \mathbb{F}_1 as pointed sets (see [7]). This link can be understood by viewing \mathfrak{s} -modules as *pointed objects* in the topos $\widehat{\Gamma}$ of covariant functors $\Gamma^{\text{op}} \rightarrow \mathfrak{Sets}$. Thus, provided one works in $\widehat{\Gamma}$, one may think of our basic objects as “pointed sets”. The reason for this choice of topos is to provide room for the identity functor $\text{Id} : \Gamma^{\text{op}} \rightarrow \Gamma^{\text{op}}$ which defines the simplest Γ -ring: \mathfrak{s} . In other words both Γ^{op} and objects in $\widehat{\Gamma}$ are based on the idea of pointed sets which underlies the naive interpretation of \mathbb{F}_1 . In this way one reaches a workable framework that strictly extends the category of \mathbb{Z} -modules.

To perform homological algebra one needs, guided by the Dold-Kan correspondence, to move from the basic category $\mathfrak{s} - \text{Mod}$ to its simplicial version, namely the category $\Gamma\mathcal{S}_*$ of Γ -spaces, where \mathcal{S}_* denote the category of pointed simplicial sets, *i.e.* contravariant functors $\Delta \rightarrow \mathfrak{Sets}_*$ where Δ is the ordinal number category and \mathfrak{Sets}_* is the category of pointed sets. The category $\Gamma\mathcal{S}_*$ plays a central role in [3]. We denote by $\underline{\text{Hom}}_{\mathcal{S}_*}$ the internal hom functor in \mathcal{S}_* . As explained in *op.cit.*, one can use the closed structure of \mathcal{S}_* to endow $\Gamma\mathcal{S}_*$ with the structure of a symmetric monoidal closed category. The closed structure is defined as follows

$$\underline{\text{Hom}}_{\Gamma\mathcal{S}_*}(M, N) := \{(k_+, [q]) \mapsto \text{Hom}_{\Gamma\mathcal{S}_*}(M \wedge \Delta[q]_+, N(k_+ \wedge -))\}. \quad (2)$$

The monoidal structure is given by the smash product where $M \wedge N$ is defined using the closed structure and can be described as a Day’s product (see *op.cit.* 2.1.2.1)

$$(M \wedge N)(Z) = \int^{(X,Y)} (M(X) \wedge N(Y)) \wedge \Gamma^{\text{op}}(X \wedge Y, Z).$$

The key result of Lydakis (*op.cit.* Theorem 2.1.2.4) states that there are choices of coherency isomorphisms so that the triple $(\Gamma\mathcal{S}_*, \wedge, \mathfrak{s})$ is a symmetric monoidal closed category.

Our goal is to use Γ -spaces to perform homological algebra in the category $\mathfrak{s} - \text{Mod}$ by applying an analogue of the Dold-Kan correspondence. For our arithmetic applications it is crucial to work with *non-fibrant* Γ -spaces and define a suitable substitute for the homotopy groups. In homotopy theory the Kan extension property is used in two ways:

- to show that the relation of homotopy is an equivalence relation,
- to define the group structure on π_n for $n \geq 1$.

To define the homology $H_n(X, F)$ of a pointed simplicial set X with coefficients in an \mathfrak{s} -module F , the problem to obtain the substitute of the group structure does not arise since, already in the classical case where $F = HA$ corresponds to an abelian group, the interchange law shows that the group structure in homology is the same as that inherited from the underlying Γ -set (see Remark 2.18). Thus the issue created by the lack of the Kan extension property occurs mainly at the level of pointed (non fibrant) simplicial sets X , and Γ^{op} is not involved there. One thus needs, as an intermediate step, to extend the combinatorial construction of the homotopy $\pi_n(X, \star)$ for a pointed simplicial set which is not fibrant. This step is described in Section 2.1 of the present paper. The main difficulty to obtain a meaningful combinatorial notion is that the relation of homotopy between n -simplices $x, y \in X_n$ as in [10] Definition 3.1 is no longer an equivalence relation. By definition (see *op.cit.*)

$$R = \{(x, y) \in X_n \times X_n \mid \partial_j x = \partial_j y \forall j \ \& \ \exists z \mid \partial_j z = s_{n-1} \partial_j x \forall j < n, \partial_n z = x, \partial_{n+1} z = y\} \quad (3)$$

The simplices involved in the definition of π_n correspond to the elements of $\text{Hom}_{\mathcal{S}_*}(S^n, X)$, *i.e.* by Yoneda's lemma to $x \in X_n$ with $\partial_j x = \star \forall j$. Here S^n is the combinatorial sphere, *i.e.* the pointed simplicial set $(\Delta[n], \partial\Delta[n])$ obtained by collapsing the boundary $\partial\Delta[n]$ of the standard simplex to a single base point. The relation R on $\text{Hom}_{\mathcal{S}_*}(S^n, X) \subset X_n$ coincides with the relation on the 0-skeleton Y_0 associated to the two boundary maps $\partial_j : Y_1 \rightarrow Y_0$, where $Y := \Omega^n(X)$ is obtained from X by iterating n -times the endofunctor $\Omega : \mathcal{S}_* \rightarrow \mathcal{S}_*$ of [11] (Definition 1.6). In this way one reduces the problem to the definition of $\pi_0 Y$ for $Y = \Omega^n(X)$. Then one can simply define $\pi_0 Y$ as the quotient $\pi_0^{\text{comb}} Y$ of Y_0 by the equivalence relation generated by R .

1.1. DEFINITION. *Let $n \geq 0$ be an integer and X a pointed simplicial set. Define*

$$\pi_n^{\text{comb}}(X) := \pi_0^{\text{comb}}(\Omega^n(X)) = \text{Hom}_{\mathcal{S}_*}(S^n, X) / \tilde{R} \quad (4)$$

where \tilde{R} is the equivalence relation generated by the restriction of the relation R of (3).

This notion developed in Section 2.1 suffices for the goals of the present paper, but for future applications we also wish to keep the finer information contained in the relation R . This is achieved by introducing the topos $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ in which the finer notion, denoted $\pi_n^{(2)}(X, \star)$, takes its value : *i.e.* $\pi_n^{(2)}(X, \star)$ is a 2-set, *i.e.* an object of $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$. This construction is described in Section 2.5 where we also show that the topos $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ is

related to the topos of quivers. In Section 2.10 we then obtain a general definition of homology of Γ -spaces, considered as simplicial Γ -sets. This homology is not, in general, a group but is a Γ -2-set *i.e.* a pointed covariant functor $\Gamma^{\text{op}} \rightarrow \mathfrak{Sets}_*^{(2)}$.

In Section 2.14 we construct, given an integer $n \geq 0$, an arbitrary pointed simplicial set X and an arbitrary \mathfrak{s} -module (Γ -set) F , the homology $H_n(X, F)$ as follows

1.2. DEFINITION. *Let $n \geq 0$ be an integer, X a pointed simplicial set, and F an \mathfrak{s} -module. Define*

$$H_n(X, F) := \{k \mapsto \pi_n^{\text{comb}}(F \circ (X \wedge k_+))\} \quad (5)$$

as an \mathfrak{s} -module.

As in [3], we extend the Γ -set F to an endofunctor of the category of pointed sets. When $F = HA$ for an abelian group A , $H_n(X, F)$ coincides with the standard definition of homology :

1.3. THEOREM. *Let A be an abelian group, and X a pointed simplicial set. For any integer $n \geq 0$ one has the equality of \mathfrak{s} -modules*

$$H_n(X, HA) = H(H_n(X, A)) \quad (6)$$

where $H_n(X, A)$ is the (reduced) abelian group homology of X with coefficients in A .

Again, we stress the fact that we apply Definition 1.2 in cases where the pointed simplicial set $F \circ (X \wedge k_+)$ is *not fibrant*. In particular, in our applications the \mathfrak{s} -modules $H_n(X, F)$ are rarely groups.

In Section 3 we apply Definition 1.2 to the \mathfrak{s} -modules we introduced in [2], at the archimedean place of $\overline{\text{Spec}} \mathbb{Z}$. We show that these coefficients yield a semi-norm on the ordinary singular homology $H_n(X, \mathbb{R})$ of a topological space X and our goal is to compare this semi-norm with the Gromov norm, whose definition is recalled in Section 3.3. In Section 3.1 we review our construction (see [2]) of the structure sheaf $\mathcal{O} \subset H\mathbb{Q}$ of \mathfrak{s} -algebras on $\overline{\text{Spec}} \mathbb{Z}$. The sheaves $\mathcal{O}(D)$ associated to Arakelov divisors $D = D_{\text{finite}} + D_{\infty}$, as in *op.cit.* provide a one parameter family of \mathfrak{s} -modules $\|H\mathbb{R}\|_{\lambda}$ ($\lambda \in \mathbb{R}_+$) which we can use as coefficients in formula (5). In Section 3.8, Proposition 3.10, we prove that for any topological space X the filtration of the singular homology group $H_n(X, \mathbb{R})$ by the $H_n(X, \|H\mathbb{R}\|_{\lambda})$ defines a semi-norm which is equivalent to the Gromov norm.

The final Section 4 is entirely devoted to show that the two norms on $H_n(X, \mathbb{R})$: the Gromov norm and our new norm, are in fact equal when $X = \Sigma$ is a compact Riemann surface. The difficulty in the proof of this result is due to the fact that in order to obtain elements of the homology $H_2(\Sigma, \|H\mathbb{R}\|_{\lambda})$ one needs to get singular chains which are not only cycles but are such that all their simplicial boundaries actually vanish. While one knows that this Moore normalization is possible the problem is to effect it without increasing the ℓ^1 -norm of the chain : this requires a delicate geometric work described in Sections 4.1 and 4.4. One then obtains the desired equality in the form of the following

1.4. THEOREM. *Let Σ be a compact Riemann surface and $[\Sigma]$ its fundamental class in homology. Then $[\Sigma]$ belongs to the range of the canonical map $H_2(\Sigma, \|H\mathbb{R}\|_\lambda) \rightarrow H_2(\Sigma, \mathbb{R})$ if and only if λ is larger than the Gromov norm of $[\Sigma]$.*

We expect that a similar statement holds in hyperbolic geometry in any dimension. The natural test ground for the homology $H_n(X, \|H\mathbb{R}\|_\lambda)$ is in hyperbolic spaces since the Gromov norm does not vanish there for $n > 1$ while it vanishes identically on all spheres. This is in contrast with the construction of the spectra associated to Γ -spaces M where the associated endofunctor $X \mapsto M \circ X$ is only tested on spheres.

2. Homology of a simplicial set with coefficients in an \mathfrak{s} -module

Our goal in this section is to reach a good definition of the homology of a pointed simplicial set with coefficients in an \mathfrak{s} -module and to show that it generalizes the standard notion in algebraic topology. This is achieved in Definition 2.15 and Theorem 2.17. As a preliminary step we need to refine the definition of the homotopy groups π_n by remaining at the combinatorial level and ignoring the group structure. Classically (see *e.g.* [3] Appendix A.2.3), the function space of maps between pointed simplicial sets X and Y is defined as the pointed simplicial set:

$$\text{Map}_*(X, Y) := \underline{\text{Hom}}_{\mathcal{S}_*}(X, \text{sin}|Y|) \tag{7}$$

This amounts to replace Y with the *fibrant* simplicial set $\text{sin}|Y|$ and it entails that the π_n , defined using such a fibrant replacement (see also *op.cit.* A.2.5.1), are then groups for $n \geq 1$ (abelian for $n > 1$). Thus in the definition of the homotopy groups of a Γ -space $M : \Gamma^{\text{op}} \rightarrow \mathcal{S}_*$ (see *op.cit.* Definition 2.2.1.2, and §2.2.1.1 where M is extended to an endofunctor of \mathcal{S}_*)

$$\pi_q M := \varinjlim_k \pi_{k+q} M(S^k) \tag{8}$$

the terms involved in the colimit are groups, hence $\pi_q M$ is an abelian group.

For our applications however, the simplification effected by the definition (7) hides certain finer features of Γ -spaces which become relevant for arithmetic constructions. We shall thus work directly in the category $\Gamma\mathcal{S}_*$ without performing this fibrant replacement.

2.1. HOMOTOPY FOR POINTED SIMPLICIAL SETS. In order to define the new homotopy $\pi_n^{\text{new}}(X, \star)$ for a general pointed simplicial set (X, \star) , we shall first reduce to the case of π_0^{new} . This reduction process by “decalage” is well known and is recalled here for convenience. We follow [11] Definition 1.9. One defines an endofunctor Ω of \mathcal{S}_* which associates to a pointed simplicial set (X, \star) the pointed simplicial set $\Omega(X, \star)$ defined as follows (with k a positive integer)

$$\Omega(X, \star)_k := \{x \in X_{1+k} \mid \partial_0(x) = \star, \partial_{i_0} \dots \partial_{i_k} x = \star, \forall i_j \in \{0, \dots, 1+k\}\} \tag{9}$$

with the simplicial structure given by faces

$$\partial_j : \Omega(X, \star)_k \rightarrow \Omega(X, \star)_{k-1}, \quad \partial_j(x) = \partial_{j+1}^X(x) \tag{10}$$

and degeneracies

$$s_j : \Omega(X, \star)_k \rightarrow \Omega(X, \star)_{k+1}, \quad s_j(x) = s_{j+1}^X(x). \quad (11)$$

The definition of the homotopy $\pi_n^{\text{new}}(X, \star)$ is then reduced to that of π_0^{new} for the simplicial set $\Omega^n(X)$ obtained after iterating the endofunctor Ω n -times :

$$\pi_n^{\text{new}}(X, \star) := \pi_0^{\text{new}}(\Omega^n(X)). \quad (12)$$

One shows by induction¹ on n that

$$\Omega^n(X, \star)_k = \{x \in X_{n+k} \mid \partial_j(x) = \star, \forall j < n, \partial_{i_0} \dots \partial_{i_k} x = \star, \forall i_j \in \{0, \dots, k+n\}\} \quad (13)$$

while the face and degeneracies are obtained as in (10) and (11) but using ∂_{j+n}^X and s_{j+n}^X . One describes directly the first levels of $\Omega^n(X)$ as follows

2.2. LEMMA. *Let (X, \star) be a pointed simplicial set.*

(i) *The 0-skeleton $(\Omega^n(X))_0$ is the set of simplices $x \in X_n$ with all $\partial_j(x)$ equal to the base point.*

(ii) *$(\Omega^n(X))_0$ coincides with $\text{Hom}_{\mathcal{S}_\star}(S^n, X) \subset X_n$ where S^n is obtained by collapsing the boundary $\partial\Delta[n]$ of the standard simplex to a single base point².*

(iii) *The 1-skeleton $(\Omega^n(X))_1$ is the set of $x \in X_{n+1}$ which fulfill the conditions*

$$\partial_i \partial_j(x) = \star, \quad \forall i, j, \quad \partial_j(x) = \star, \quad \forall j \in \{0, \dots, n-1\}.$$

(iv) *The boundaries $\partial_i : (\Omega^n(X))_1 \rightarrow (\Omega^n(X))_0$ for $i = 0, 1$ are given by ∂_n and ∂_{n+1} .*

(v) *The relation R on $(\Omega^n(X))_0 = \text{Hom}_{\mathcal{S}_\star}(S^n, X) \subset X_n$ given by*

$$xRy \iff \exists z \in (\Omega^n(X))_1 \text{ s.t. } \partial_0 z = x \text{ and } \partial_1 z = y$$

coincides with the relation of homotopy between n -simplices as in (3).

PROOF. (i) Follows from (13) for $k = 0$.

(ii) By Yoneda's lemma one checks that the morphisms $y \in \text{Hom}_{\mathcal{S}_\star}(S^n, X)$ i.e. the elements of $\text{Hom}_{\mathcal{S}_\star}(\Delta[n], X)$ which send $\partial\Delta[n]$ to the base point, are the same as the elements of the 0-skeleton $(\Omega^n(X))_0$.

(iii) Follows from (13) for $k = 1$.

(iv) Follows from $\partial_j = \partial_{j+n}^X$ for $j = 0, 1$.

(v) This follows from the previous part of the lemma since the relation (3) restricts to

$$R = \{(x, y) \in X_n \times X_n \mid \partial_j x = \partial_j y = \star \forall j \ \& \ \exists z \mid \partial_j z = \star \forall j < n, \partial_n z = x, \partial_{n+1} z = y\} \quad (14)$$

■

¹Note that a product $\partial_{i_0} \dots \partial_{i_k}$ can be reordered using the simplicial rules so that the indices fulfill $i_0 \geq i_1 \geq \dots \geq i_k$.

²This is not the definition used in [3], where S^n is defined as the n -fold smash product $S^1 \wedge \dots \wedge S^1$ of $S^1 = \Delta[1]/\partial\Delta[1]$. This distinction in the definition of the homotopy groups is irrelevant in the fibrant case since the geometric realizations are homeomorphic, but as in [11] our choice is more convenient to compute the set of maps using Yoneda's Lemma.

2.3. REMARK. *The geometric meaning of the above well known endofunctor Ω can be understood starting from the “decalage” (see [6] or [3] A.2.7) which gives a combinatorial model PX mimicking the path space of a simplicial set X by precomposing the functor X with the endofunctor $\text{Dec}_1 = [0] \amalg \bullet : \Delta \rightarrow \Delta$. This simply shifts the indices i.e. one has $(PX)_k = X_{k+1}$ and the indices of faces and degeneracies are shifted by 1. The link with ordinary paths is given by precomposing with the morphism of simplicial sets $\gamma : \Delta[1] \times \Delta[q] \rightarrow \Delta[q+1]$, associated as $\gamma := N(p)$ by the nerve functor N to*

$$p : [1] \times [q] \rightarrow [q+1], \quad p(0, j) := 0 \quad \forall j, \quad p(1, j) := j+1 \quad \forall j$$

Requiring that the two end points of the path associated to $x \in X_{k+1} = \text{Hom}_\Delta(\Delta[q+1], x)$ are equal to the base point \star (when X is pointed) gives exactly the conditions of (9) defining $\Omega(X)$. When X is fibrant one obtains in this way a model for its loop space.

For a *fibrant* simplicial pointed set X , the relation (14) is an *equivalence* relation and the quotient by this relation defines $\pi_0(\Omega^n(X))$ which is known to be a group, for $n \geq 1$ (see [10, 11], or Theorem 7.2 in Chapter III of [4]). Note also that when X is fibrant the above equivalence relation on $\text{Hom}_{\mathcal{S}_*}(S^n, X) \subset X_n$ coincides with the one defined by the two boundary maps from the 1-skeleton of the simplicial set $\underline{\text{Hom}}_{\mathcal{S}_*}(S^n, X)$ (see [11] Lemma 1B.3).

On the other hand, the simplicial sets X we consider here are *not necessarily fibrant* and the relation R is not in general transitive (nor symmetric). The easy solution to bypass this problem is to define $\pi_n^{\text{comb}}(X)$ as the quotient by the equivalence relation generated by the relation R in agreement with Definition 1.1. This provides a first notion of homotopy which suffices for the goal of the present paper. One has by construction

$$\pi_n^{\text{comb}}(X, \star) := \pi_0^{\text{comb}}(\Omega^n(X)). \tag{15}$$

We state simple properties of this combinatorial notion

2.4. PROPOSITION. (i) *Let X be a pointed simplicial set and $k > 0$ an integer. Then for any n*

$$\pi_n^{\text{comb}}(X \wedge k_+) = \pi_n^{\text{comb}}(X) \wedge k_+.$$

(ii) *Let X, Y be pointed simplicial sets, one has for any n*

$$\pi_n^{\text{comb}}(X \times Y) = \pi_n^{\text{comb}}(X) \times \pi_n^{\text{comb}}(Y).$$

PROOF. (i) An element $x \in (X \wedge k_+)_n$, $x \neq \star$ is of the form $x = (a, j)$ with $a \in X_n$ and $0 < j \leq k$. Two elements $x = (a, j)$ and $x' = (a', j')$ fulfill $(x, x') \in R$ as in (14) if and only if $j = j'$ and $(a, a') \in R_X$ since the boundaries preserve the index j .

(ii) This follows since $(X \times Y)_n = X_n \times Y_n$ and the boundaries act componentwise. ■

2.5. THE FINER NOTION $\pi_n^{(2)}(X)$ AND THE TOPOS $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$. For later applications to Arakelov divisors Definition 1.1 is too coarse and one would like to

- keep all the information about the relation R and
- still think of π_0^{new} as a set.

The idea of “topos” of Grothendieck [1] comes to the rescue providing a satisfactory answer. We consider the topos $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ of contravariant functors to the category of sets from the small category obtained by restricting the objects of Δ to $[0]$ and $[1]$ and keeping the same morphisms as in the following definition

2.6. DEFINITION. (i) Let $X : \Delta^{\text{op}} \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}$ be a simplicial set. We define $\pi_0^{(2)}(X)$ as the object of $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ which is the restriction of the functor X to the full subcategory of Δ with objects $[0], [1]$ and same morphisms as Δ .
(ii) Let X be a pointed simplicial set, then we define

$$\pi_n^{(2)}(X) := \pi_0^{(2)}(\Omega^n(X)).$$

It turns out that the topos $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ can also be described as the dual of the small category with a single object whose morphisms form the monoid \mathcal{M} with three elements $1, m_0, m_1$ and the multiplication table specified by the rule $m_j x = m_j$ for all $j \in \{0, 1\}$.

2.7. PROPOSITION. The topos $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ is the same as the presheaf category $\widehat{\mathcal{M}}$ of the monoid \mathcal{M} , i.e. the category of sets with a right action of \mathcal{M} .

PROOF. By definition an object F of the topos $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ is a pair of sets $F(0), F(1)$, with two maps $\partial_j : F(1) \rightarrow F(0)$, $j \in \{0, 1\}$ and a map $s : F(0) \rightarrow F(1)$ such that $\partial_j \circ s = \text{Id}$. This implies that $s : F(0) \rightarrow F(1)$ is an injection and one can thus view $F(0)$ as a subset of $F(1)$ and consider the two self-maps $T_j = s \circ \partial_j : F(1) \rightarrow F(1)$. They fulfill the rule

$$T_i \circ T_j = T_j, \quad \forall i, j \in \{0, 1\}$$

since $s \circ (\partial_i \circ s) \circ \partial_j = s \circ (\text{Id}) \circ \partial_j = s \circ \partial_j$. Thus one obtains an object in $\widehat{\mathcal{M}}$ where \mathcal{M} is the monoid defined by the opposite of the above rules. Conversely given an object X of $\widehat{\mathcal{M}}$, i.e. a set X endowed with a right action of \mathcal{M} one defines an object of $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ by setting $F(1) := X$, $F(0) := \text{Range}(T_j)$ which does not depend on the choice of $j \in \{0, 1\}$. One lets $s : F(0) \rightarrow F(1)$ be the inclusion as a subset, and $\partial_j : F(1) \rightarrow F(0)$ is given by T_j . One checks that $\partial_j \circ s = \text{Id}$. One obtains in this way two functors $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)} \rightarrow \widehat{\mathcal{M}}$ and $\widehat{\mathcal{M}} \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ which are inverse of each other. ■




Figure 1: Subobject classifier for $\mathbf{Sets}^{(2)}$.

2.8. REMARK. (i) Two presheaf categories \widehat{C} and \widehat{D} are equivalent if and only if the categories C and D have isomorphic Karoubi completions. The full subcategory of Δ on the objects $[0], [1]$ has two idempotents on $[1]$ with isomorphic image on $[2]$ and forms the “splitting idempotents” of \mathcal{M} thus Proposition 2.7 follows directly.

(ii) The topos $\mathbf{Sets}^{(2)}$ is closely related to the topos of quivers but is not the same. In fact the topos of quivers has two points given by the functors to the set of vertices and the functor to the set of edges. Similarly these two functors give the two points of the topos $\mathbf{Sets}^{(2)}$ but in the latter case the functor to the set of edges never takes the value \emptyset when the functor to the set of vertices takes a non-empty value.

The determination of the subobject classifier shows that the topos $\mathbf{Sets}^{(2)}$ is two valued and not boolean (see [8], VI).

2.9. LEMMA. The subobject classifier Ω of the topos $\mathbf{Sets}^{(2)}$ is the object with two vertices False, True and five edges which besides the two degenerate ones form the graph of Figure 1.

PROOF. It is a general fact (see [8], §I.4) that for a topos of the form $\widehat{\mathcal{M}}$ for a monoid \mathcal{M} , i.e. the topos of sets with a right action of \mathcal{M} , the subobject classifier is given by the set \mathcal{J} of right ideals of \mathcal{M} on which the right action of \mathcal{M} is defined by

$$J.m := \{n \in \mathcal{M} \mid mn \in J\}, \quad \forall J \in \mathcal{J}, m \in \mathcal{M}.$$

Taking the above \mathcal{M} with three elements $1, m_0, m_1$ and the multiplication table specified by the rule $m_j x = m_j$ for all $j \in \{0, 1\}$, one finds that \mathcal{J} contains five elements

$$\mathcal{J} = \{\emptyset, \{m_0\}, \{m_1\}, \{m_0, m_1\}, \mathcal{M}\}$$

and that the right action T_j of $m_j \in \mathcal{M}$ fixes \emptyset and \mathcal{M} (which are hence degenerate edges, i.e. vertices) while $T_j\{m_j\} = \mathcal{M}$ and $T_i\{m_j\} = \emptyset$ for $i \neq j$. Thus the set V of vertices contains two elements \emptyset and \mathcal{M} and the non-degenerate edges are the three edges shown in Figure 1.

The reason for renaming the vertices \emptyset as “False” and \mathcal{M} as “True” and for the choice of the labels of the edges comes from the construction of the classifying map associated to a subobject G' of an object G in $\mathbf{Sets}^{(2)}$. One finds that the classifying map f is obtained as follows as a map from G to Ω :

1. $\epsilon \in G' \Rightarrow f(\epsilon) = \text{True}$
2. $\epsilon \notin G'$ and $\partial_j \epsilon \notin G' \Rightarrow f(\epsilon) = \text{False}$
3. $\epsilon \notin G'$, $\partial_0 \epsilon \notin G'$ and $\partial_1 \epsilon \in G' \Rightarrow f(\epsilon) = \text{Repair}$
4. $\epsilon \notin G'$, $\partial_0 \epsilon \in G'$ and $\partial_1 \epsilon \notin G' \Rightarrow f(\epsilon) = \text{Doubt}$
5. $\epsilon \notin G'$, $\partial_0 \epsilon \in G'$ and $\partial_1 \epsilon \in G' \Rightarrow f(\epsilon) = \text{Check}$.

The terminology “False” and “True” is the standard one for the two extremes in subobject classifiers, the notations for the edges are suggestive but more arbitrary. ■

2.10. HOMOTOPY OF Γ -SPACES. If \mathcal{C} is a pointed category with initial and final object denoted $*$, one defines (see [3]) the category of Γ -objects of \mathcal{C} as the category $\Gamma\mathcal{C}$ of pointed covariant functors $\Gamma^{\text{op}} \rightarrow \mathcal{C}$. This construction applies to the category \mathcal{S}_* of pointed simplicial sets to yield the category $\Gamma\mathcal{S}_*$ of Γ -spaces. It also applies to the category $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*^{(2)}$ of pointed objects in $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$. We shall call Γ -2-sets the objects of $\Gamma\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*^{(2)}$.

2.11. PROPOSITION. (i) Let X be a Γ -space and $n \in \mathbb{N}$. Then the map $k \mapsto \pi_n^{(2)}(X(k_+))$ (resp. $k \mapsto \pi_n^{\text{comb}}(X(k_+))$) extends to a pointed covariant functor $\pi_n^{(2)}(X) : \Gamma^{\text{op}} \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*^{(2)}$ (resp. to a Γ -set).

(ii) For $n \in \mathbb{N}$, $\pi_n^{(2)}$ defines a functor $\pi_n^{(2)} : \Gamma\mathcal{S}_* \rightarrow \Gamma\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*^{(2)}$ from Γ -spaces to Γ -2-sets.

(iii) For $n \in \mathbb{N}$, π_n^{comb} defines a functor $\pi_n^{\text{comb}} : \Gamma\mathcal{S}_* \rightarrow \Gamma\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ from Γ -spaces to \mathfrak{s} -Mod.

PROOF. This follows from the naturality of Definitions 1.1 and 2.6. ■

The relation between $\pi_n^{(2)}$ and π_n^{comb} is given by

$$\pi_n^{\text{comb}} := \ell \circ \pi_n^{(2)} \quad (16)$$

i.e. composition with the functor

$$\ell : \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)} \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}, \quad \ell(X) := \lim_{\mathcal{C}^{\circ}} X(c)$$

which assigns to a 2-set its set of components. Here, \mathcal{C} is any of the small categories defining $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$ as the presheaf category $\widehat{\mathcal{C}}$ as in Proposition 2.7 and \mathcal{C}° is its opposite. Note that the functor ℓ does not correspond to a point of the topos $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}^{(2)}$.

2.12. Γ -SETS AS ENDOFUNCTORS . In this section we recall the construction of [3] of the endofunctor in the category \mathcal{S}_* associated to a Γ -space, in the case of discrete Γ -spaces, i.e. \mathfrak{s} -modules. By construction an \mathfrak{s} -module is a covariant functor $M : \Gamma^{\text{op}} \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ and, as in Section 2.1.2.1 of *op.cit.*, we view pointed sets as discrete pointed simplicial sets, i.e. as constant functors $\Delta^{\text{op}} \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$.

2.13. LEMMA. Let $M : \Gamma^{\text{op}} \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ be an \mathfrak{s} -module. Then the associated endofunctor of the category \mathcal{S}_* of pointed simplicial sets is obtained by composition with M viewed as an endofunctor of the category $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ of pointed sets.

PROOF. One first extends the functor $M : \Gamma^{\text{op}} \rightarrow \mathfrak{S}\text{ets}_*$ to an endofunctor $\mathfrak{S}\text{ets}_* \rightarrow \mathfrak{S}\text{ets}_*$ in pointed sets. This is done by taking a colimit on the finite subsets as explained in §2.2.1.1 of [3]. Then, one applies the technique described in *op.cit.* that uses, for a simplicial set $X = \{[q] \mapsto X_q\}$, the diagonal

$$M(X) := \{[q] \mapsto M(X_q)_q\}.$$

Since by construction $M(X_q)$ is a discrete simplicial set, it is the same in all degrees so that the index q in $M(X_q)_q$ disappears, thus we simply write $M(X_q)$. Hence starting with the pointed simplicial set $X : \Delta^{\text{op}} \rightarrow \mathfrak{S}\text{ets}_*$, we obtain a new pointed simplicial set by composition *i.e.*

$$X \mapsto M(X) = M \circ X : \Delta^{\text{op}} \rightarrow \mathfrak{S}\text{ets}_*. \tag{17}$$

In summary the result follows from §2.2.1.1 of *op.cit.* . ■

The basic example of an \mathfrak{s} -module is given in 2.1.2.1 of [3] where one associates to an abelian monoid A with a zero element, the functor $M = HA$

$$HA(k_+) = A^k, \quad Hf : HA(k_+) \rightarrow HA(n_+), \quad Hf(m)(j) := \sum_{f(\ell)=j} m_\ell \tag{18}$$

where $m = (m_1, \dots, m_k) \in HA(k_+)$. The zero element of A gives meaning to the empty sum. In the special case when the monoid A is an abelian group, the composition (17), *i.e.* the functor $HA \circ X : \Delta^{\text{op}} \rightarrow \mathfrak{S}\text{ets}_*$ factors through simplicial abelian groups (the functor HA is the composite of a more precise functor $A \mapsto \text{Ab}A$ to abelian groups with the forgetful functor from abelian groups to pointed sets, where the base point is the 0) and always fulfills the Kan extension property. The geometric realization $|HA \circ X|$ only uses the underlying simplicial pointed set but the finer structure as a simplicial abelian group, and the Dold-Kan correspondence in the form of Corollary 2.5 of [4], Chapter III, show that the homotopy groups of the geometric realization $|HA \circ X|$ are given by the (reduced) homology³ of the associated complex of abelian groups, *i.e.* $\pi_n(|HA \circ X|) = H_n(X, A)$. This suffices to conclude for instance that $|HA \circ S^n|$ is an Eilenberg-MacLane space $K(A, n)$.

2.14. THE HOMOLOGY WITH COEFFICIENTS IN AN \mathfrak{s} -MODULE . In our arithmetic context we are interested in \mathfrak{s} -modules M which are no longer of the form HA where A is an abelian group. In a first class of examples M is still of the form HA , where A is a monoid. A second class of examples are those constructed in [2] to specify the geometric structure of $\overline{\text{Spec } \mathbb{Z}}$ at the archimedean place. In all these cases it is no longer true that the composite $M \circ X$ is fibrant, even when the simplicial set X itself is fibrant. We shall use the equality $\pi_n(|HA \circ X|) = H_n(X, A)$ holding for abelian groups as the motivation to extend the definition of the homology of a pointed simplicial set with coefficients in an arbitrary \mathfrak{s} -module as follows

³For a pointed simplicial set $(X, *)$ we use the notation $H_n(X, A)$ for the reduced homology $H_n((X, *), A)$.

2.15. DEFINITION. Let M be an \mathfrak{s} -module, and X a pointed simplicial set. For any integer $n \geq 0$ one defines the homology $H_n(X, M)$ as the \mathfrak{s} -module

$$H_n(X, M)(k_+) := \pi_n^{\text{comb}}(M \circ (X \wedge k_+)). \quad (19)$$

Here, k_+ is viewed as a discrete simplicial pointed set *i.e.* constant in all degrees.

As in Lemma 2.13, M is viewed as an endofunctor of the category $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ and π_n^{comb} is defined in Definition 1.1. There is in fact a refined version of homology $H_n^{(2)}(X, M)$ using $\pi_n^{(2)}$ instead of π_n^{comb} but we shall not need it in the present paper.

The following result establishes several basic properties of the new homology. The \mathfrak{s} -module HM makes sense for any additive monoid with a zero element, and this applies to the monoid $\mathbb{B} := \{0, 1\}$ with $1 + 1 = 1$ so that the \mathfrak{s} -module $H\mathbb{B}$ in (v) below is well defined.

2.16. PROPOSITION. (i) For any $n \geq 0$, $H_n(X, M)$ is a covariant bifunctor

$$H_n : \mathcal{S}_* \times \mathfrak{s} - \text{Mod} \longrightarrow \mathfrak{s} - \text{Mod}.$$

(ii) Let M_1, M_2 be \mathfrak{s} -modules. One has a natural transformation

$$H_n(M_1 \circ X, M_2) \rightarrow H_n(X, M_2 \circ M_1)$$

which is an isomorphism when evaluated on 1_+ .

(iii) For any pointed simplicial set X one has $H_n(X, \mathfrak{s}) = \pi_n^{\text{comb}}(X) \wedge \mathfrak{s}$.

(iv) For $n \neq m$: $H_m(S^n, \mathfrak{s}) = \{*\}$ while for $n = m$ one has $H_m(S^n, \mathfrak{s}) = \mathfrak{s}$.

(v) For $n \neq m$: $H_m(S^n, H\mathbb{B}) = \{*\}$ while for $n = m$ one has $H_m(S^n, H\mathbb{B}) = H\mathbb{B}$.

PROOF. (i) By construction, $H_n(X, M)$ is a covariant functor of X for fixed M , and of M for fixed X . To prove that it is a bifunctor it suffices, using the bifunctor lemma (see [9] Proposition 1 Chapter II, §3), to show that it satisfies the interchange law which states that given morphisms $f \in \text{Hom}_{\mathcal{S}_*}(X, Y)$ and $h \in \text{Hom}_{\mathfrak{s}}(M, N)$ one has the equality

$$H_n(f, N) \circ H_n(X, h) = H_n(Y, h) \circ H_n(f, M) \in \text{Hom}_{\mathfrak{s}}(H_n(X, M), H_n(Y, N)). \quad (20)$$

Both sides of this formula are \mathfrak{s} -modules *i.e.* functors $\Gamma^{\text{op}} \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ thus it is enough to check the equality pointwise *i.e.* by evaluating both sides on k_+ for fixed k . Since $\pi_n^{\text{comb}} : \mathcal{S}_* \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ is a functor the equality follows provided one shows that the same equality holds if one replaces $H_n(X, M)$ by $F(X, M) := M \circ X$ which is a separately covariant functor to \mathcal{S}_* with arguments in \mathcal{S}_* and $\mathfrak{s} - \text{Mod}$. Again it is enough to check this equality pointwise *i.e.* replacing \mathcal{S}_* by $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ and $F(X, M) := M \circ X$ by $G(X, M) := M(X)$ which is a separately covariant functor to $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ with arguments in $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ and $\mathfrak{s} - \text{Mod}$. Since M and N are endofunctors of $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ and the morphism $h \in \text{Hom}_{\mathfrak{s}}(M, N)$ is a natural transformation from M to N one has, for any $f \in \text{Hom}_{\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*}(X, Y)$, the equality

$$N(f) \circ h_X = h_Y \circ M(f) \in \text{Hom}_{\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*}(M(X), N(Y))$$

which gives (20).

(ii) As in [3], (2.2.1.2 equation (2.2)), one has natural maps $M(X) \wedge Y \mapsto M(X \wedge Y)$. We apply this with $Y = k_+$ and thus obtain natural maps

$$\eta_k : M_1(X) \wedge k_+ \rightarrow M_1(X \wedge k_+), \quad \forall k.$$

This yields a natural morphism

$$M_2(\eta_k) : M_2(M_1(X) \wedge k_+) \rightarrow M_2(M_1(X \wedge k_+))$$

and by composition with π_n^{comb} one gets the natural transformation

$$\pi_n^{\text{comb}}(M_2(\eta_k)) : H_n(M_1 \circ X, M_2)(k_+) \rightarrow H_n(X, M_2 \circ M_1)(k_+)$$

which is, by construction, an isomorphism for $k = 1$.

(iii) Since the endofunctor of \mathfrak{Sets}_* associated to \mathfrak{s} is the identity, the result follows from Proposition 2.4 (i).

(iv) Using (iii) it is enough to determine $\pi_n^{\text{comb}}(S^m)$. The pointed simplicial set S^n is obtained by collapsing $\partial\Delta[n]$ to a base point. This means that one considers the sub-functor $[q] \mapsto \partial\Delta([q]) \subset \text{Hom}_\Delta([q], [n])$ given by the maps $[q] \rightarrow [n]$ which are not surjective and one identifies all the elements of $\partial\Delta([q])$ with the base point. For $h \in \text{Hom}_\Delta([q'], [q])$ one has $\partial\Delta([q]) \circ h \subset \partial\Delta([q'])$ so that the collapsing gives a pointed simplicial set. An element of $\text{Hom}_{\mathcal{S}_*}(S^n, X)$ is an element of $\text{Hom}_{\mathcal{S}_*}(\Delta[n], X)$ which maps $\partial\Delta[n]$ to the base point. This means, by Yoneda's lemma, an element $x \in X_n$ such that $\partial_j(x) = *$ for all j (since any map $[q] \rightarrow [n]$ which is not surjective factors through a $d_j : [n-1] \rightarrow [n]$). For $X = S^m$, such an $x \in X_n$ is, if it is not the base point, an element $\phi \in \text{Hom}_\Delta([n], [m])$ which is surjective and such that $\phi \circ d_j$ fails to be surjective for any j . This latter condition implies that ϕ is also injective and one concludes that $n = m$ and ϕ is the identity map. This gives $\pi_n^{\text{comb}}(S^m) = \{*\}$ for $n \neq m$. To prove that $\pi_n^{\text{comb}}(S^n) = \{*, \text{Id}\}$ one just needs to show that the element Id does not get identified with the base point under the equivalence relation generated by the relation (14), *i.e.*

$$xRy \iff \exists z \mid \partial_j z = * \forall j < n, \partial_n z = x, \partial_{n+1} z = y\}.$$

Any $z \neq *$ in S_{n+1}^n is given by a surjective map $s_i \in \text{Hom}_\Delta([n+1], [n])$ such that $s_i(i) = s_i(i+1)$ and the condition $\partial_j z = * \forall j < n$ shows that the index i is equal to $i = n$. It follows that $\partial_n z = \partial_{n+1} z$ and that the relation R is the diagonal.

(v) The endofunctor $H\mathbb{B}$ associates to a pointed set E the (pointed) set of all finite subsets of E which contain the base point $*$, and to a map $f : E \rightarrow F$ the direct image map $Z \mapsto f(Z)$. Note the equivalence,

$$f(Z) = \{*\} \iff f(x) = *, \quad \forall x \in Z \tag{21}$$

It follows that there are only two elements $u \in (H\mathbb{B} \circ S^n)_n = H\mathbb{B}(S^n) = H\mathbb{B}(\{*, \text{Id}\})$, namely the base point $*$ and the subset $u = \{*, \text{Id}\}$. Let us show that these two elements

are not equivalent under the equivalence relation generated by the relation (14) for the simplicial pointed set $H\mathbb{B} \circ S^n$. An element of $(H\mathbb{B} \circ S^n)_{n+1} = H\mathbb{B}(S_{n+1}^n)$ is a subset $z = \{\ast, s_{i_1}, \dots, s_{i_k}\}$ of the set S_{n+1}^n described in the proof of (iv). The condition $\partial_j z = \ast \forall j < n$ shows that all indices i_j are equal to n so that either $z = \ast$ or $z = \{\ast, s_n\}$. This shows, as in the proof of (iv), that the relation R is diagonal and $\pi_n^{\text{comb}}(H\mathbb{B} \circ S^n) = H\mathbb{B}(1_+) = \mathbb{B}$. Let then $k > 0$ be an integer and E a pointed set. One has a natural isomorphism $H\mathbb{B} \circ (E \wedge k_+) \simeq (H\mathbb{B} \circ E)^k$. It follows that for any pointed simplicial set X one has a natural isomorphism $H\mathbb{B} \circ (X \wedge k_+) \rightarrow (H\mathbb{B} \circ X)^k$. Then by Proposition 2.4 (ii) one gets

$$\pi_n^{\text{comb}}(H\mathbb{B} \circ (S^n \wedge k_+)) = (\pi_n^{\text{comb}}(H\mathbb{B} \circ S^n))^k = H\mathbb{B}(k_+).$$

By construction the natural identifications are compatible with the structures of Γ -sets. This shows that $H_n(S^n, H\mathbb{B}) = H\mathbb{B}$. The proof of (iv) together with (21) show that $H_n(S^n, H\mathbb{B}) = \{\ast\}$ for $m \neq n$. ■

Definition 2.15 provides a meaning to the following equality (22) whose two sides are \mathfrak{s} -modules.

2.17. THEOREM. *Let A be an abelian group, and X a pointed simplicial set. For any integer $n \geq 0$ one has the equality of \mathfrak{s} -modules*

$$H_n(X, HA) = H(H_n(X, A)) \quad (22)$$

where $H_n(X, A)$ is the (reduced) abelian group homology of X with coefficients in A .

PROOF. For any simplicial set Y the composite $HA \circ Y$ is a simplicial abelian group and hence has the Kan extension property (see [11], Theorem 2.2). It follows that the combinatorial homotopy $\pi_n^{\text{comb}}(HA \circ Y)$ coincides with the usual homotopy $\pi_n(|HA \circ Y|)$ of the geometric realization

$$\pi_n^{\text{comb}}(HA \circ Y) = \pi_n(|HA \circ Y|). \quad (23)$$

Moreover the group law of these homotopy groups coincides with the abelian group law inherited from the simplicial abelian group structure (see *op.cit.* Proposition 2.4). The Dold-Kan correspondence (see [4], Chapter III Corollary 2.5) gives a canonical bijection

$$\delta_Y : H_n(Y, A) \rightarrow \pi_n^{\text{comb}}(HA \circ Y). \quad (24)$$

Furthermore this bijection is a natural transformation of covariant functors from pointed simplicial sets to pointed sets. More precisely given a morphism $\psi : Y \rightarrow Y'$ of pointed simplicial sets, one obtains the equality

$$\pi_n^{\text{comb}}(HA(\psi)) \circ \delta_Y = \delta_{Y'} \circ H_n(\psi, A). \quad (25)$$

Indeed, it is enough to check this equality on cycles $c \in Z_n(Y, A)$ which are Moore normalized, *i.e.* $\partial_j c = 0 \forall j$. The element $\delta_Y(c)$ is then given by the combinatorial class directly

associated to c viewed as an element of $\text{Hom}_{\mathcal{S}^*}(S^n, HA \circ Y) = \text{Hom}_{\mathcal{S}^*}((\Delta^n, \partial\Delta^n), HA \circ Y)$ which is a subset of $(HA \circ Y)_n = HA(Y_n)$. Let $c = \sum a_j y_j$ where $a_j \in A$ and $y_j \in Y_n$. Then $\pi_n^{\text{comb}}(HA(\psi)) \circ \delta_Y(c)$ is represented by the combinatorial class obtained from $\delta_Y(c)$ by applying the functor $HA(\psi)$. This gives $HA(\psi)(\sum a_j y_j) = \sum a_j \psi(y_j)$, as one sees using the definition (18) of the functor HA . But one has similarly $H_n(\psi, A)(c) = \sum a_j \psi(y_j)$, thus one gets the required equality (25).

From (24) one gets the bijection $\delta_X : H_n(X, A) \rightarrow \pi_n^{\text{comb}}(HA \circ X)$. Let then $k > 0$ be an integer and E a pointed set. One has a natural isomorphism $HA \circ (E \wedge k_+) \simeq (HA \circ E)^k$ since both sides consist of maps $(x, j) \mapsto \phi(x, j) \in A$, $x \in E$, $j \in \{1, \dots, k\}$ with finite support and such that $\phi(*, j) = 0$ for all j . Thus one obtains a natural isomorphism of simplicial sets

$$HA \circ (X \wedge k_+) = (HA \circ X)^k.$$

The same equality holds for the geometric realizations, and using (23) one derives

$$\pi_n^{\text{comb}}(HA \circ (X \wedge k_+)) = \pi_n(|HA \circ (X \wedge k_+)|) = \pi_n(|(HA \circ X)^k|) = H_n(X, A)^k.$$

At the set-theoretic level this coincides with $H(H_n(X, A))(k_+)$. In fact one can obtain the same result more directly as a consequence of (24) and of the equality of (reduced) homology groups $H_n(X_1 \vee X_2, A) = H_n(X_1, A) \oplus H_n(X_2, A)$.

It remains to show that given a morphism $\phi : k_+ \rightarrow m_+$ in Γ^{op} the associated map

$$\pi_n^{\text{comb}}(HA \circ (X \wedge k_+)) \rightarrow \pi_n^{\text{comb}}(HA \circ (X \wedge m_+)),$$

is the same as the map $HK(\phi) : H_n(X, A)^k \rightarrow H_n(X, A)^m$ associated to the group law of $K = H_n(X, A)$ and the functor HK . Using (25) it is enough to show that $HK(\phi)$ equals the homology map

$$H_n(\text{Id}_X \wedge \phi, A) : H_n(X \wedge k_+, A) \rightarrow H_n(X \wedge m_+, A).$$

With e_j , $j \in \{1, \dots, k\}$, the canonical basis of $H_0(k_+)$, and $e_0 := 0$, the above map is given by

$$H_n(\text{Id}_X \wedge \phi, A)(\sum c_j \otimes e_j) = \sum c_j \otimes e_{\phi(j)}$$

and using the definition (18) of the functor HK one gets the required equality. ■

2.18. REMARK. *In homotopy theory the homotopy groups π_n are abelian groups for $n > 1$. The group operation arises, at the combinatorial level, from the Kan extension property of fibrant simplicial sets together with combinatorial constructions involving simplices. Definition 2.15 does not involve any of these constructions and yet Theorem 2.17 shows that one recovers the same group law on the homotopy groups π_n from the Γ -set (\mathfrak{s} -module) obtained using the functorial nature of the map $k_+ \mapsto X \wedge k_+$. The reason behind this equality of structures is the interchange law which is fulfilled by the group law of the homotopy group π_n and the group law induced by the abelian coefficients. In that sense, Definition 2.15 takes into account the \mathfrak{s} -module structure of the coefficients to obtain a replacement of the group structure of homotopy groups. We shall see in the next sections a striking example where this additional structure is put to work.*

3. The archimedean place and the Gromov norm

In this section we show that the singular homology $H_*(X, \mathbb{R})$ of a topological space inherits a natural semi-norm from the filtration of the \mathfrak{s} -module $H\mathbb{R}$ by the sub- \mathfrak{s} -modules $\|H\mathbb{R}\|_\lambda$ associated to the archimedean place of $\overline{\text{Spec } \mathbb{Z}}$ as constructed in [2]. Moreover we prove that this semi-norm is equivalent to the Gromov semi-norm on singular homology.

3.1. $\overline{\text{Spec } \mathbb{Z}}$ AT THE ARCHIMEDEAN PLACE. In [2] we showed how to endow the Arakelov compactification $\overline{\text{Spec } \mathbb{Z}}$ with a structure sheaf of \mathfrak{s} -algebras, which coincides with the standard structure sheaf of $\text{Spec } \mathbb{Z}$ on the dense open set $\text{Spec } \mathbb{Z} \subset \overline{\text{Spec } \mathbb{Z}}$ using the fully faithful functor H from rings to \mathfrak{s} -algebras. The new feature is the structure of this sheaf at the archimedean place which is obtained using the following proposition⁴ of [2]

3.2. PROPOSITION. (i) *Let R be a semiring, and $\| \cdot \|$ a sub-multiplicative seminorm on R . Then HR is naturally endowed with a structure of \mathfrak{s} -subalgebra $\|HR\|_1 \subset HR$ defined as follows*

$$\|HR\|_1 : \Gamma^{\text{op}} \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_* \quad \|HR\|_1(F) := \{\phi \in HR(F) \mid \sum_{F \setminus \{*\}} \|\phi(x)\| \leq 1\}. \quad (26)$$

(ii) *Let E be an R -semimodule and $\| \cdot \|^E$ a seminorm on E such that $\|a\xi\| \leq \|a\|\|\xi\|$, $\forall a \in R, \forall \xi \in E$, then for any $\lambda \in \mathbb{R}_+$ the following defines a module $\|HE\|_\lambda^E$ over $\|HR\|_1$*

$$\|HE\|_\lambda^E : \Gamma^{\text{op}} \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_* \quad \|HE\|_\lambda^E(F) := \{\phi \in HE(F) \mid \sum_{F \setminus \{*\}} \|\phi(x)\|^E < \lambda\}. \quad (27)$$

The first statement of Proposition 3.2 is applied for the ring $R = \mathbb{Q}$ of rational numbers and its archimedean absolute value to construct the stalk at ∞ of the structure sheaf. One obtains in this way a sheaf $\mathcal{O} \subset H\mathbb{Q}$ of \mathfrak{s} -algebras over $\overline{\text{Spec } \mathbb{Z}}$. The second statement of Proposition 3.2 is then applied to the one-dimensional real vector space \mathbb{R} to obtain, given an Arakelov divisor $D = D_{\text{finite}} + D_\infty$, the sheaf $\mathcal{O}(D)$ of \mathcal{O} -modules over $\overline{\text{Spec } \mathbb{Z}}$

$$\mathcal{O}(D)(\Omega) := \|H\mathcal{O}(D_{\text{finite}})(\Omega \setminus \{\infty\})\|_{e^a}, \quad D_\infty = a\{\infty\}. \quad (28)$$

Thus the \mathfrak{s} -modules at work at the archimedean place depend on a positive real parameter $\lambda > 0$ and are implemented by the functor $\|H\mathbb{R}\|_\lambda : \Gamma^{\text{op}} \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ which associates to a pointed set F the pointed set

$$\|H\mathbb{R}\|_\lambda(F) = \{x : F \rightarrow \mathbb{R} \mid \#\{j, x(j) \neq 0\} < \infty \ \& \ \sum |x(j)| < \lambda\}. \quad (29)$$

In fact (29) describes also the extension of $\|H\mathbb{R}\|_\lambda$ as an endofunctor of $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$. There is an obvious analogue of (29) when \mathbb{R} is replaced by \mathbb{Q} and this analogue is what is needed in (28); on the other hand it is more natural to work with the local field \mathbb{R} associated to the archimedean place of \mathbb{Q} .

⁴With the nuance that in (27) we use the strict inequality.

3.3. SIMPLICIAL VOLUME. We recall briefly the notion of simplicial volume introduced by M. Gromov [5]. Let first X be a topological space and $C_*(X, \mathbb{R})$ the associated singular chain complex with real coefficients. One defines the ℓ^1 -norm on singular chains as follows

$$\|c\|_1 := \sum |a_j|, \quad \forall c = \sum a_j \sigma_j, \quad \sigma_j \in \text{Top}(\Delta^*, X) \tag{30}$$

The induced semi-norm on the singular homology $H_*(X, \mathbb{R})$ is the quotient semi-norm

$$\|\alpha\|_1 := \inf_{c \in \alpha} \|c\|_1. \tag{31}$$

The Gromov norm $|M|$ of an oriented closed connected manifold of dimension n is then defined as the semi-norm of its fundamental class $|M| := \|[M]\|_1$. A fundamental result of the theory ([5], [13] Thm 6.2) is the proportionality principle:

3.4. THEOREM. (*M. Gromov*) *Let Σ be any compact oriented hyperbolic manifold of dimension $n > 1$, then one has*

$$|\Sigma| = \frac{v(\Sigma)}{v_n}$$

where $v(\Sigma)$ is the volume of Σ and v_n is the maximal volume of straight simplices in hyperbolic space.

We refer to [13] chapter 6 for the description of the straightening of singular simplices and singular chains. The constant v_2 is equal to π and one thus has

3.5. COROLLARY. *Let Σ be a Riemann surface of genus $g > 1$, then $|\Sigma| = 4(g - 1)$.*

The fact that the norm does not vanish is dual to the boundedness of cohomology and this holds in the hyperbolic case, thus for $k > 1$ the semi-norm (31) is in fact a norm on the homology $H_k(M, \mathbb{R})$ when M is an hyperbolic manifold (see *op.cit.*).

3.6. MOORE NORMALIZATION. Let A be a simplicial abelian group. The standard complex (still denoted A for simplicity) of abelian groups associated to A is defined using the boundary map

$$\partial := \sum_0^n (-1)^j d_j : A_n \rightarrow A_{n-1}. \tag{32}$$

The associated normalized complex NA is defined as follows

$$NA_n := \cap_0^{n-1} \text{Ker } d_j \subset A_n, \quad d := d_n : NA_n \rightarrow NA_{n-1} \tag{33}$$

(the simplicial identity $d_n d_{n-1} = d_{n-1} d_{n-1}$ shows that it defines a complex). For each n one lets $D_n \subset A_n$ be the subgroup generated by the ranges of the degeneracies. The boundary map ∂ of (32) fulfills $\partial(D_n) \subset D_{n-1}$ and induces a map

$$\partial : A_n/D_n \rightarrow A_{n-1}/D_{n-1}.$$

The corresponding quotient complex A/D is the complex modulo degeneracies. By construction one has two morphisms of complexes $i : NA \rightarrow A$ and $p : A \rightarrow A/D$. Moreover

(see [4], Theorem 2.1), the morphism $p \circ i : NA \rightarrow A/D$ is an isomorphism of chain complexes. It follows that the composite morphism $\nu := (p \circ i)^{-1} \circ p : A \rightarrow NA$ is a projection. As in the proof of Theorem 2.4 of [4], one constructs explicitly a chain map $f^{(n)} : A_n \rightarrow NA_n$ as the composition

$$f^{(n)} := f_{n-2}^{(n)} \circ \dots \circ f_j^{(n)} \circ f_0^{(n)} \quad (34)$$

where $f_j^{(n)} : A_n \rightarrow A_n$ is defined as $f_j^{(n)} = \text{Id} - s_{j+1}d_{j+1}$. Moreover one also constructs explicitly a chain homotopy $T_k : A_k \rightarrow A_{k+1}$ such that

$$\text{Id} - f^{(n)} = T \circ \partial + \partial \circ T. \quad (35)$$

Since each $f_j^{(n)}$ acts as the identity in the quotient A_n/D_n the same holds for $f^{(n)}$ and one obtains the equality $\nu_n = f^{(n)}$.

3.7. LEMMA. *Let X be pointed simplicial set. Let the simplicial vector space $A = H\mathbb{R} \circ X$ be endowed with the norm*

$$\|\phi\| := \sum |\phi(x)|, \quad \forall \phi \in A_n = H\mathbb{R}(X_n). \quad (36)$$

(i) *The linear map $\nu_n = f^{(n)} : A_n \rightarrow NA_n$ is of norm $\leq 2^{n-1}$.*

(ii) *Let $c \in Z_n(A)$ be a cycle. Then $\nu_n(c)$ is a homologous normalized cycle and $\|\nu_n(c)\| \leq 2^{n-1}\|c\|$.*

PROOF. (i) The statement follows from (34) and the inequalities

$$\|f_j^{(n)}\| \leq \|\text{Id}\| + \|s_{j+1}d_{j+1}\| \leq 2, \quad \|f \circ g\| \leq \|f\|\|g\|.$$

(ii) This follows from (i) and (35). ■

On the real vector space $H_n(X, \mathbb{R})$ we consider the following semi-norm which is induced by the ℓ^1 -norm (36) on the normalized complex NA_n :

$$\|c\|^{\text{nor}} := \inf\{\|\phi\| \mid \phi \in NA_n, \phi \sim c\} \quad (37)$$

where $\phi \sim c$ means that $\phi \in NA_n$ is homologous to the cycle $c \in Z_n(A)$. By applying Lemma 3.7 one obtains the basic inequalities

$$\|c\|_1 \leq \|c\|^{\text{nor}} \leq 2^{n-1}\|c\|_1, \quad \forall c \in H_n(X, \mathbb{R}). \quad (38)$$

3.8. EQUIVALENCE WITH THE GROMOV NORM. The filtration of the \mathfrak{s} -module $H\mathbb{R}$ by the sub- \mathfrak{s} -modules $\|H\mathbb{R}\|_\lambda \subset H\mathbb{R}$, $\lambda \in \mathbb{R}_+$, of (29) provides, for any pointed simplicial set X and integer $n \geq 0$ natural morphisms of \mathfrak{s} -modules

$$\rho_{n,\lambda} : H_n(X, \|H\mathbb{R}\|_\lambda) \rightarrow H_n(X, H\mathbb{R}), \quad (39)$$

and a filtration by the ranges of the $\rho_{n,\lambda}$. Theorem 2.17 gives a natural isomorphism $H_n(X, H\mathbb{R}) = H(H_n(X, \mathbb{R}))$.

3.9. THEOREM. *Let X be a pointed simplicial set. For any integer $n \geq 0$ and $\lambda \in \mathbb{R}_+$, the range of the natural morphism of \mathfrak{s} -modules is*

$$\rho_{n,\lambda}(H_n(X, \|H\mathbb{R}\|_\lambda)) = \|H(H_n(X, \mathbb{R}))\|_\lambda^{\text{nor}} \quad (40)$$

where $\|c\|^{\text{nor}}$ is the semi-norm defined in (37).

PROOF. It follows from (29) that the endofunctor $\|H\mathbb{R}\|_\lambda$ assigns to a pointed set (X, \star) the set of maps with finite support

$$\|H\mathbb{R}\|_\lambda(X) := \{\phi : X \rightarrow \mathbb{R} \mid \phi(\star) = 0, \#\{x \mid \phi(x) \neq 0\} < \infty, \sum |\phi(x)| < \lambda\}.$$

By construction the range $\rho_{n,\lambda}(H_n(X, \|H\mathbb{R}\|_\lambda))$ is a sub-functor of $H(H_n(X, \mathbb{R}))$ thus to show (40) it is enough to prove that for any integer $k > 0$ one has

$$\rho_{n,\lambda}(H_n(X, \|H\mathbb{R}\|_\lambda))(k_+) = \|H(H_n(X, \mathbb{R}))\|_\lambda^{\text{nor}}(k_+). \quad (41)$$

The right hand side of (41) is given by k -tuples $(\gamma_j)_{1 \leq j \leq k}$, $\gamma_j \in H_n(X, \mathbb{R})$ such that $\sum \|\gamma_j\|^{\text{nor}} < \lambda$, *i.e.*

$$\exists \phi_j \in H\mathbb{R} \circ X_n \mid d_i \phi_j = 0 \forall i, \phi_j \sim \gamma_j, \sum_{X_n \times \{1, \dots, k\}} |\phi_j(x)| < \lambda.$$

For the left hand side of (41) one has

$$\|H\mathbb{R}\|_\lambda(X_n \wedge k_+) =$$

$$\{(\psi_j)_{j \in \{1, \dots, k\}} \mid \psi_j : X_n \rightarrow \mathbb{R}, \psi_j(\star) = 0, \#\{x \mid \psi_j(x) \neq 0\} < \infty, \sum |\psi_j(x)| < \lambda\}$$

and moreover the simplicial structure satisfies

$$d_i((\psi_j)_{j \in \{1, \dots, k\}}) = (d_i \psi_j)_{j \in \{1, \dots, k\}}.$$

Thus the 0-chains $\text{Hom}_{\mathcal{S}_\star}(S^n, \|H\mathbb{R}\|_\lambda(X \wedge k_+))$ are exactly the same as the ones involved in the right hand side of (41) and one gets (40). ■

For a topological space X one lets $\text{sin}X$ be the associated simplicial set of singular simplices

$$\text{sin}X = \{[n] \mapsto \text{Top}(\Delta^n, X)\}$$

where the standard simplex Δ^n of dimension n is given concretely as

$$\Delta^n = \{(\lambda_0, \dots, \lambda_n) \mid \lambda_j \geq 0, \sum \lambda_j = 1\}$$

Then, Definition 2.15 extends to topological spaces and arbitrary \mathfrak{s} -modules as

$$H_n(X, M) := H_n(\text{sin}X, M) \quad (42)$$

3.10. COROLLARY. *Let X be a topological space. The filtration of the singular homology group $H_n(X, \mathbb{R})$ by the \mathfrak{s} -modules $H_n(X, \|H\mathbb{R}\|_\lambda)$ defines a semi-norm which is equivalent to the Gromov norm.*

PROOF. This follows from Theorem 3.9 and the basic inequalities (38). ■

4. Equality for Riemann surfaces of genus $g > 1$

We show that for a compact oriented 2-dimensional manifold Σ of genus $g > 1$, the normalized norm (37) on singular homology agrees with the Gromov norm. Since these two norms are equivalent and the Gromov norm vanishes except on $H_2(\Sigma, \mathbb{R})$ it is enough to prove the equality for the fundamental class $[\Sigma] \in H_2(\Sigma, \mathbb{R})$. The difficulty is to construct singular cycles c in the homology class $[\Sigma]$ which not only have ℓ^1 -norm $\|c\|_1$ close to the expected value $4(g-1)$ but are also *normalized*, *i.e.* such that all boundaries vanish $\partial_j(c) = 0$. This is achieved in three steps. In section 4.1 we deal with the relative situation of the building block K and construct a normalized cycle relative to its boundary ∂K . In section 4.4 we assemble together g copies of K and obtain a surface of genus g and a normalized cycle representing the fundamental class whose ℓ^1 -norm is $4g$. The third step is standard and uses cyclic covers to improve the estimate to the expected value $4(g-1)$.

4.1. MOORE NORMALIZATION FOR THE BUILDING BLOCK. A compact oriented 2-dimensional manifold Σ of genus $g > 1$ is obtained by gluing together g copies of a building block K which we now describe. This building block is the quotient of the convex polygon $\text{Conv}(0, 1, 2, 3, 4, 5)$ of Figure 2 by the equivalence relation R generated by

$$\Delta(\{1, 2\})(x) \sim_R \Delta(\{4, 3\})(x), \quad \Delta(\{2, 3\})(x) \sim_R \Delta(\{5, 4\})(x), \quad \forall x \in \Delta^1$$

where given $n+1$ points (P_0, \dots, P_n) in the real affine plane $E = \mathbb{R}^2$, one denotes

$$\Delta(\{P_0, \dots, P_n\}) \in \text{Top}(\Delta^n, E), \quad (\lambda_0, \dots, \lambda_n) \mapsto \sum \lambda_j P_j.$$

By transitivity one finds that the five vertices $(1, 2, 3, 4, 5)$ are equal modulo R , since $1 \sim_R 4 \sim_R 3 \sim_R 2 \sim_R 5$. Thus one has by construction a continuous map

$$\gamma : \text{Conv}(0, 1, 2, 3, 4, 5) \rightarrow K. \quad (43)$$

The building block K thus consists of 4 triangles with the common vertex 0 and where the external sides are identified following the rules

$$\begin{aligned} \Delta(\{1, 2\}) &\sim \Delta(\{4, 3\}), & \Delta(\{2, 1\}) &\sim \Delta(\{3, 4\}), \\ \Delta(\{2, 3\}) &\sim \Delta(\{5, 4\}), & \Delta(\{3, 2\}) &\sim \Delta(\{4, 5\}). \end{aligned} \quad (44)$$

It is shown geometrically in Figure 3 as a subset of the 2-torus (before the identifications of the edges) and in Figures 4 and 5 after these identifications have been performed. These Figures keep track of the natural triangulations.

By construction one has $\Delta(\{P_0, \dots, P_n\}) \in \text{Top}(\Delta^n, \text{Conv}(P_j))$ where $\text{Conv}(P_j)$ is the convex hull of the points P_j . The composition

$$\Delta'(\{P_0, \dots, P_n\}) := \gamma \circ \Delta(\{P_0, \dots, P_n\}) \in \text{Top}(\Delta^n, K) \quad (45)$$

defines singular simplices *i.e.* elements of $\text{sin } K$. From (44) one obtains the equalities

$$\begin{aligned} \Delta'(\{1, 2\}) &= \Delta'(\{4, 3\}), & \Delta'(\{2, 1\}) &= \Delta'(\{3, 4\}), \\ \Delta'(\{2, 3\}) &= \Delta'(\{5, 4\}), & \Delta'(\{3, 2\}) &= \Delta'(\{4, 5\}) \end{aligned} \quad (46)$$




Figure 2: Basic polygon in $E = \mathbb{R}^2$.




Figure 3: Basic polygon inside the fundamental domain for \mathbb{Z}^2 acting on \mathbb{R}^2 .




Figure 4: Triangulated building block K .




Figure 5: Neighborhood of point P .

To each pair (i, j) of > 0 integer indices we associate the simplicial chain $c(0, i, j)$

$$c(0, i, j) := \Delta'(\{0, i, j\}) + \Delta'(\{i, j, 0\}) + 2\Delta'(\{j, 0, i\}) - \Delta'(\{j, i, 0\}) - \Delta'(\{0, j, i\}) - 2\Delta'(\{i, 0, j\}) \quad (47)$$

The boundaries of $c(0, i, j)$ are described as follows

4.2. LEMMA. *The following equalities hold*

$$\begin{aligned} \partial_0(c(0, i, j)) &= 2\Delta'(\{0, i\}) - 2\Delta'(\{0, j\}) - \Delta'(\{i, 0\}) + \Delta'(\{j, 0\}) + \Delta'(\{i, j\}) - \Delta'(\{j, i\}) \\ \partial_1(c(0, i, j)) &= -\Delta'(\{0, i\}) + \Delta'(\{0, j\}) + \Delta'(\{i, 0\}) - \Delta'(\{j, 0\}) + 2\Delta'(\{j, i\}) - 2\Delta'(\{i, j\}) \\ \partial_2(c(0, i, j)) &= \Delta'(\{0, i\}) - \Delta'(\{0, j\}) + 2\Delta'(\{j, 0\}) - 2\Delta'(\{i, 0\}) + \Delta'(\{i, j\}) - \Delta'(\{j, i\}) \end{aligned}$$

PROOF. The result follows by linearity of the ∂_j and the equalities

$$\partial_0\Delta'(\{a, b, c\}) = \Delta'(\{b, c\}), \quad \partial_1\Delta'(\{a, b, c\}) = \Delta'(\{a, c\}), \quad \partial_2\Delta'(\{a, b, c\}) = \Delta'(\{a, b\})$$

■




Figure 6: Surface of genus 2.




Figure 7: Domain for surface of genus 3.

We now combine the above simplicial chains and use the rules (46) to get a chain which is normalized relative to the boundary ∂K of K .

4.3. LEMMA. *Let $c_{(1,5)} := \sum_{1 \leq i \leq 4} c(0, i, i + 1)$. One has*

$$\begin{aligned} \partial_0 c_{(1,5)} &= 2\Delta'(\{0, 1\}) - 2\Delta'(\{0, 5\}) - \Delta'(\{1, 0\}) + \Delta'(\{5, 0\}) \\ \partial_1 c_{(1,5)} &= -\Delta'(\{0, 1\}) + \Delta'(\{0, 5\}) + \Delta'(\{1, 0\}) - \Delta'(\{5, 0\}) \\ \partial_2 c_{(1,5)} &= \Delta'(\{0, 1\}) - \Delta'(\{0, 5\}) - 2\Delta'(\{1, 0\}) + 2\Delta'(\{5, 0\}) \end{aligned}$$

PROOF. The cancelations follow from the equalities

$$\begin{aligned} \sum_{1 \leq i \leq 4} (\Delta'(\{0, i\}) - \Delta'(\{0, i + 1\})) &= \Delta'(\{0, 1\}) - \Delta'(\{0, 5\}) \\ \sum_{1 \leq i \leq 4} (\Delta'(\{i, 0\}) - \Delta'(\{i + 1, 0\})) &= \Delta'(\{1, 0\}) - \Delta'(\{5, 0\}) \end{aligned}$$

and from the following one which uses the rules (46)

$$\sum_{1 \leq i \leq 4} (\Delta(\{i, i + 1\}) - \Delta(\{i + 1, i\})) = 0$$

since (46) shows that the following terms all vanish

$$\Delta(\{1, 2\}) - \Delta(\{4, 3\}), \Delta(\{2, 3\}) - \Delta(\{5, 4\}), \Delta(\{3, 4\}) - \Delta(\{2, 1\}), \Delta(\{4, 5\}) - \Delta(\{3, 2\})$$

We thus get the required formulas. ■

4.4. MOORE NORMALIZATION FOR A RIEMANN SURFACE OF GENUS $g > 1$. Let $g > 1$ and P be obtained (see Figure 7) as the union of g copies $P(w) = P_{(1+4w, 5+4w)}$ for $0 \leq w < g$, of the basic polygon $\text{Conv}(0, 1, 2, 3, 4, 5)$ of Figure 2, where the side $(0, 5 + 4w)$ is common to $P(w)$ and $P(w + 1)$ for $w < g - 1$ and is common to $P(g - 1)$ and $P(0)$ for $w = g - 1$, while the external sides are identified pairwise as in P . The quotient of P by these identifications is a surface $\Sigma(g)$ of genus g .

4.5. LEMMA. *The singular chain*

$$c = c_{(1,5)} + c_{(5,9)} + \dots + c_{(1+4w,5+4w)} + \dots + c_{(1+4(g-1),1)}$$

is closed and normalized, i.e. one has $\partial_j c = 0$ for $j \in \{0, 1, 2\}$.

PROOF. By Lemma 4.3 one gets for $0 \leq w \leq g-1$, and with $5 + 4(g-1) \sim 1$,

$$\partial_0 c_{(1+4w,5+4w)} = 2\Delta'(\{0, 1 + 4w\}) - 2\Delta'(\{0, 5 + 4w\}) - \Delta'(\{1 + 4w, 0\}) + \Delta'(\{5 + 4w, 0\})$$

which gives

$$\partial_0 c = \partial_0 \sum_{0 \leq w \leq g-1} c_{(1+4w,5+4w)} = \sum_{0 \leq w \leq g-1} \partial_0 c_{(1+4w,5+4w)} = 0.$$

The same reasoning applies to show that $\partial_j c = 0$ for $j \in \{1, 2\}$. ■

4.6. LEMMA. *The singular chain c of Lemma 4.5 represents the singular homology class $8[\Sigma]$ ($[\Sigma]$ = fundamental class of Σ).*

PROOF. The result follows since each chain $c(0, i, i+1)$ as in (47) is homologous to $8\Delta'(\{0, i, i+1\})$ while the $4g$ triangles $\Delta'(\{0, i, i+1\})$ for $1 \leq i \leq 4g$ give a triangulation of Σ . ■

4.7. LEMMA. *The ℓ^1 -norm of the singular chain c of Lemma 4.5 is $\leq 32g$.*

PROOF. This follows from the triangle inequality and the definition (47) of the chain $c(0, i, j)$ whose ℓ^1 -norm is ≤ 8 . ■

4.8. THEOREM. *Let Σ be a compact Riemann surface and $[\Sigma]$ its fundamental class in homology. Then $[\Sigma]$ belongs to the range of the canonical map $H_2(\Sigma, \|\mathbb{H}\mathbb{R}\|_\lambda) \rightarrow H_2(\Sigma, \mathbb{R})$ if and only if λ is larger than the Gromov norm of $[\Sigma]$.*

PROOF. The result follows from Theorem 3.9 if one shows that the fundamental class $[\Sigma]$ fulfills the equality

$$\|[\Sigma]\|^{\text{nor}} = \|[\Sigma]\|_1$$

The inequality \geq follows from (38). Moreover, as recalled in section 3.3, for a surface of genus g the Gromov norm $\|[\Sigma]\|_1$ is equal to $4(g-1)$. Thus it remains to show that $\|[\Sigma]\|^{\text{nor}} \leq 4(g-1)$. By applying Lemmas 4.6 and 4.7 one obtains the inequality $\|[\Sigma]\|^{\text{nor}} \leq 4g$. One then applies a standard technique which is to use the same inequality for the covering space Σ' of Σ associated to an infinite cyclic subgroup of the fundamental group $\pi_1(\Sigma)$. The genus of a cyclic cover Σ' of degree n is $g' = n(g-1) + 1$ since the Euler characteristic is multiplied by n . Thus the inequality $\|[\Sigma']\|^{\text{nor}} \leq 4g'$ entails

$$n\|[\Sigma]\|^{\text{nor}} \leq 4g' = 4(n(g-1) + 1).$$

By passing to the limit when $n \rightarrow \infty$ one obtains the desired inequality $\|[\Sigma]\|^{\text{nor}} \leq 4(g-1)$. ■

References

- [1] M. Artin, A. Grothendieck, J-L. Verdier, eds. (1972), *SGA₄*, LNM 269-270-305, Berlin; New York: Springer-Verlag.
- [2] A. Connes, C. Consani, *Absolute algebra and Segal's Gamma sets*, J. Number Theory 162 (2016), 518–551.
- [3] B. Dundas, T. G. Goodwillie, R. McCarthy, *The local structure of algebraic K-theory*. Algebra and Applications, 18. Springer-Verlag London, Ltd., London, 2013.
- [4] P. G. Goerss, J. F. Jardine, *Simplicial homotopy theory*. Progress in Mathematics, 174. Birkhauser Verlag, Basel, 1999.
- [5] M. Gromov, *Volume and bounded cohomology*, Inst. Hautes Etudes Sci. Publ. Math. (1982), no. 56, 5–99 (1983).
- [6] L. Illusie, *Complexe cotangent et déformations*. Lecture Notes in Mathematics, Vol. 283. Springer-Verlag, Berlin-New York, 1972.
- [7] M. Kapranov, A. Smirnov, *Cohomology determinants and Reciprocity Laws* (preprint).
- [8] S. MacLane, I. Moerdijk, *Sheaves in geometry and logic. A first introduction to topos theory*. Corrected reprint of the 1992 edition. Universitext. Springer-Verlag, New York, 1994.
- [9] S. MacLane, *Categories for the working mathematician*. Graduate Texts in Mathematics, Vol. 5. Springer-Verlag, New York-Berlin, 1971. ix+262 pp.
- [10] P. May *Simplicial objects in algebraic topology*. Van Nostrand Mathematical Studies, No. 11 D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London 1967.
- [11] J. C. Moore *Seminar on algebraic homotopy theory*, Princeton, 1956 (mimeographed notes).
- [12] G. Segal, *Categories and cohomology theories*. Topology, 13 (1974), 293–312.
- [13] W. Thurston, *Geometry and topology of 3-manifolds*, Lecture Notes Princeton (1978).

I.H.E.S. and Ohio State University

*Department of Mathematics, The Johns Hopkins University
Baltimore, MD 21218 USA*

Email: alain@connes.org
kc@math.jhu.edu

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

INFORMATION FOR AUTHORS L^AT_EX₂ε is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

T_EXNICAL EDITOR. Michael Barr, McGill University: michael.barr@mcgill.ca

ASSISTANT T_EX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Julie Bergner, University of Virginia: jeb2md@virginia.edu

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Gabriella Böhm, Wigner Research Centre for Physics: bohm.gabriella@wigner.mta.hu

Valeria de Paiva, Nuance Communications Inc: valeria.depaiva@gmail.com

Richard Garner, Macquarie University: richard.garner@mq.edu.au

Ezra Getzler, Northwestern University: getzler@northwestern.edu

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Dirk Hofmann, Universidade de Aveiro: dirk@ua.pt

Pieter Hofstra, Université d' Ottawa: phofstra@uottawa.ca

Anders Kock, University of Aarhus: kock@math.au.dk

Joachim Kock, Universitat Autònoma de Barcelona: kock@mat.uab.cat

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: matias.menni@gmail.com

Ieke Moerdijk, Utrecht University: i.moerdijk@uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Kate Ponto, University of Kentucky: kate.ponto@uky.edu

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: ross.street@mq.edu.au

Tim Van der Linden, Université catholique de Louvain: tim.vanderlinden@uclouvain.be