

A CLOSED MODEL STRUCTURE ON THE CATEGORY OF WEAKLY UNITAL DG CATEGORIES, II

PIERGIORGIO PANERO AND BORIS SHOIKHET

ABSTRACT. In this paper, which is subsequent to our previous paper (but can be read independently from it), we continue our study of the closed model structure on the category of small weakly unital dg categories (in the sense of Kontsevich-Soibelman) over a field. In our previous paper, we constructed a closed model structure on the category of weakly unital dg categories, imposing a technical condition on the weakly unital dg categories, namely the idempotency $\text{id}_x \cdot \text{id}_x = \text{id}_x$ for any object. Although this condition led us to a great simplification, it was redundant and had to be dropped.

Here we get rid of this condition, and provide a closed model structure in full generality. The new closed model category is as well cofibrantly generated, and it is proven to be Quillen equivalent to the closed model category of (strictly unital) dg categories, given by Tabuada. Dropping the idempotency condition makes the construction of the closed model structure more distant from loc.cit., and requires new constructions. One of them is a pre-triangulated hull of a wu dg category, which in turn is shown to be a wu dg category as well.

One example of a weakly unital dg category which naturally appears is the bar-cobar resolution of a dg category. We supply this paper with a refinement of the classical bar-cobar resolution of a unital dg category which is strictly unital (appendix B). A similar construction can be applied to constructing a cofibrant resolution of weakly unital dg categories.

1. Introduction

Many algebraic constructions, dealing with unital dg algebras (or, more generally, unital dg categories) give rise to only weakly ones. The simplest example is the bar-cobar resolution $R(A) = \text{Cobar}(\text{Bar}(A))$ of a dg algebra A (resp., a dg category A): for unital A , $R(A)$ is only weakly unital. Another example which we keep in mind is a further generalisation of the twisted tensor product of small dg categories [Sh], which is supposed to have better homotopical and monoidal properties, and which exists only in the weakly unital context. It would be beneficial to have a more relaxed (than the closed model category [Tab1] of small dg categories) closed model category, which is formed by small

We are thankful to Bernhard Keller for his encouragement. The work of both authors was partially supported by the FWO Research Project Nr. G060118N. The work of B.Sh. was partially supported by the HSE University Basic Research Program.

Received by the editors 2020-10-05 and, in final form, 2021-04-18.

Transmitted by Clemens Berger. Published on 2021-04-19.

2020 Mathematics Subject Classification: 18N40, 18G35.

Key words and phrases: dg-category, closed model category, weak units.

© Piergiorgio Panero and Boris Shoikhet, 2021. Permission to copy for private use granted.

weakly unital dg categories, and which is Quillen equivalent to the closed model category of small dg categories loc.cit. In this paper, we completely solve this problem.

Among the three definitions of a weakly unital dg category, existing in literature (see [Ly2] for an overview of all of them), the definition given by Kontsevich and Soibelman [KS, Sect. 4.2] seems to be the only one which gives rise to a closed model structure. In this paper, we work with this definition (we recall it in Section 2).

In this paper, we provide a cofibrantly generated Quillen model structure on the category $\mathbb{C}_{dgwu}(\mathbb{k})$ of Kontsevich-Soibelman weakly unital dg categories [KS, Sect. 4.2] over a field \mathbb{k} (we recall the definition in Section 2). We also establish a Quillen equivalence between the model category $\mathbb{C}_{dgwu}(\mathbb{k})$ and the category $\mathbb{C}_{dg}(\mathbb{k})$ of unital dg categories over \mathbb{k} (with Tabuada’s model structure [Tab1] on it).

In our previous paper [PS], we constructed a cofibrantly generated Quillen model structure on the category $\mathbb{C}_{dgwu}^0(\mathbb{k})$ of small Kontsevich-Soibelman weakly unital dg categories¹ over a field \mathbb{k} with an extra condition saying that $\text{id}_X \circ \text{id}_X = \text{id}_X$ for any object X . We also proved that there was a Quillen equivalence $L: \mathbb{C}_{dgwu}^0(\mathbb{k}) \rightleftarrows \mathbb{C}_{dg}(\mathbb{k}) : R$, where $\mathbb{C}_{dg}(\mathbb{k})$ is the closed model structure on the category of small (unital) dg categories over \mathbb{k} . The condition $\text{id}_X \circ \text{id}_X = \text{id}_X$ had a technical nature and was imposed to avoid some difficulties in the proofs we experienced for the general case of $\mathbb{C}_{dgwu}(\mathbb{k})$. However, this condition is rather artificial and is not fulfilled in any reasonable example, so the next step is to get rid of it. We complete it here.

Our results are counterparts of the main results of [PS] in this general setting, though the proofs need some essentially new ideas and constructions. Among them, we draw reader’s attention to the pre-triangulated hull of a weakly unital dg category (Section 3), and a weakly unital replacement of the Kontsevich dg category \mathcal{K} (Section 4.4).

We prove

1.1. THEOREM. [proven in Theorem 4.17] *For a field \mathbb{k} , there is a cofibrantly generated Quillen model structure on $\mathbb{C}_{dgwu}(\mathbb{k})$.*

1.2. THEOREM. [proven in Prop. 5.2 and Theorem 5.3] *There is a Quillen equivalence*

$$L: \mathbb{C}_{dgwu}(\mathbb{k}) \rightleftarrows \mathbb{C}_{dg}(\mathbb{k}) : R$$

where $\mathbb{C}_{dg}(\mathbb{k})$ is endowed with the Tabuada closed model structure [Tab1].

Here we outline in more detail our results and the organisation of the paper.

In Section 2, we recall the main definitions from [PS], and the results from loc.cit., used here. These are small completeness and small cocompleteness of $\mathbb{C}_{dgwu}(\mathbb{k})$, and the monadic description of it. Namely, there is a monad on the category of unital dg graphs $\mathbb{G}_{dgu}(\mathbb{k})$, such as the algebras over this monad are the weakly unital dg categories. Moreover, this monad is associated with a non-symmetric dg operad, called \mathcal{O}' . Here one of the main results is cited from loc.cit. as Theorem 2.5 here. It says that a natural map

¹Our notations here are different from the ones in [PS]. Our $\mathbb{C}_{dgwu}(\mathbb{k})$ here was denoted by $\text{Cat}'_{dgwu}(\mathbb{k})$ in [PS], and our $\mathbb{C}_{dgwu}^0(\mathbb{k})$ was $\text{Cat}_{dgwu}(\mathbb{k})$ in [PS].

gives a quasi-isomorphism of dg operads $\mathcal{O}' \rightarrow \mathcal{A}ssoc_+$ to the operad of unital dg algebras. This computation is essentially used in the proof that the Quillen pair of Prop. 5.2 is a Quillen equivalence. This Section makes it possible reading this paper independently on [PS].

In Section 3, we prove that to any weakly unital dg-category C we can functorially associate the pretriangulated hull of it C^{pretr} , and this non unital dg-category is in turn weakly unital. Here the main point, which makes the problem non-trivial, is that, for a closed morphism f in $C(x, y)$, the identity morphism $\text{id}_{\text{Cone}(f)}$ within the “naive” definition, fails to be closed, if $\text{id}_y \circ f \neq f \circ \text{id}_x$. This construction comes out naturally when one perturbs the naive construction making $\text{id}_{\text{Cone}(f)}$ closed. This tool allows us to work with cones of any closed morphism in C , and these are key points for crucial Lemma 4.8 and Lemma 4.16.

In Section 4, we construct a closed model structure on $\mathbb{C}_{dgwu}(\mathbb{k})$. Here the strategy is rather close to [Tab1] and [PS], though the generating cofibrations I and generating acyclic cofibrations J should be re-defined to the weakly unital context, making them “fattened”. This “fattening” goes straightforwardly for all generating (acyclic) cofibrations except to the one in which the Kontsevich dg category \mathcal{K} is involved. The refined wu version of \mathcal{K} is not straightforward (see Section 4.4, and (20)). It is a dg category which classifies a homotopy equivalence in a wu dg category (Lemma 4.7). We also prove crucial Lemma 4.8, connecting \mathcal{K} and weakly unital pre-triangulated dg hull. This Lemmas are used in the proof of key-lemma Lemma 4.16, which completes the story.

In Section 5, we establish a Quillen equivalence between $\mathbb{C}_{dgwu}(\mathbb{k})$ and $\mathbb{C}_{dg}(\mathbb{k})$. Here we essentially use Theorem 2.5, saying that the dg operad \mathcal{O}' is quasi-isomorphic $\mathcal{A}ssoc_+$.

Finally, Section 6 contains a computational proof of Proposition 3.3.

We supply the paper with three Appendices. In Appendix A, we just fix signs in the identities on the Taylor coefficients of an A_∞ morphism. Appendix B contains a (seemingly new but elementary) construction of a unital cofibrant resolution of a unital dg algebra, easily generalizing to the case of dg categories. Appendix C details a computation of particular coequalizers, following from the general discussion in [PS]; this computation is used in the proof of Theorem 5.3.

2. Weakly unital dg categories

Recall the definition of a weakly unital dg category [KS, 4.2].

Let A be a non unital dg category. Denote by $A \oplus \mathbb{k}_A$ the strictly unital dg category where $Ob(A \oplus \mathbb{k}_A) = Ob(A)$ and

$$\text{Hom}_{A \oplus \mathbb{k}_A}(x, y) = \begin{cases} \text{Hom}_A(x, y) & \text{if } x \neq y \\ \text{Hom}_A(x, x) \oplus \mathbb{k}1_x & \text{if } x = y. \end{cases}$$

One has a natural imbedding $i : A \rightarrow A \oplus \mathbb{k}_A$, sending x to x , and $f \in A(x, x)$ to the pair $(f, 0) \in (A \oplus \mathbb{k}_A)(x, x)$. We denote by 1_x the generator of \mathbb{k}_x .

2.1. DEFINITION. A weakly unital dg category A over \mathbb{k} is a non-unital dg category A over \mathbb{k} with a distinguished closed element $\text{id}_x \in A^0(x, x)$ for any object x in A , such that there exists an A_∞ -functor $p : A \oplus \mathbb{k}_A \rightarrow A$ which is the identity on the objects, such that $p \circ i = \text{id}_A$, $p_1(1_x) = \text{id}_x, \forall x \in \text{Ob}(A)$, and $p_n(f_1, \dots, f_n) = 0$ for $n \geq 2$ and all f_i morphisms in the image $i(A)$.

2.2. DEFINITION. Let A, C be two weakly unital dg categories, with the structure maps $p^A : A \oplus \mathbb{k}_A \rightarrow A$ and $p^C : C \oplus \mathbb{k}_C \rightarrow C$. A weakly unital dg functor $F : A \rightarrow C$ is a non-unital dg functor $F : A \rightarrow C$ such that the following diagram commutes:

$$\begin{array}{ccc}
 A \oplus \mathbb{k}_A & \xrightarrow{F \oplus \mathbb{k}_F} & C \oplus \mathbb{k}_C \\
 p^A \downarrow & & \downarrow p^C \\
 A & \xrightarrow{F} & C
 \end{array} \tag{1}$$

In this way, we define the category $\mathbb{C}_{dgwu}(\mathbb{k})$. Its full subcategory, for which $\text{id}_x \circ \text{id}_x = \text{id}_x$ for any object x , is denoted by $\mathbb{C}_{dgwu}^0(\mathbb{k})$.

It follows from the definition that:

$$\begin{aligned}
 F(\text{id}_x) &= \text{id}_{F(x)} & \forall x \in \text{Ob}(A) \\
 F(p_n^A(f_1, \dots, f_n)) &= p_n^C(F(f_1), \dots, F(f_n)) \quad f_i \in A \oplus \mathbb{k}_A, i = 1 \dots n
 \end{aligned} \tag{2}$$

2.3. EXAMPLE. Let C be a strictly unital dg category. We denote by $i(C)$ the weakly unital dg category, for which the A_∞ functor $p : C \oplus \mathbb{k}_C \rightarrow C$ is a natural dg functor, so that all $p_n^{i(C)}, n \geq 2$ are equal to 0. In this way we get a fully-faithful embedding $i : \mathbb{C}_{dg}(\mathbb{k}) \rightarrow \mathbb{C}_{dgwu}(\mathbb{k})$.

This definition is due to Kontsevich-Soibelman loc.cit. There are two other definitions, due to Lyubashenko and Fukaya, correspondingly. We refer the reader to [Ly], [LyMa] for discussion of all three definitions, and to [COS] for interplay between them.

An advantage of the Kontsevich-Soibelman definition is that the category of small KS wu dg categories carries a closed model structure, unlike for the two other definitions. Such closed model structure was provided in [PS] for $\mathbb{C}_{dgwu}^0(\mathbb{k})$, and is done here for the general case of $\mathbb{C}_{dgwu}(\mathbb{k})$, see Theorem 4.17 below.

We recall the definition of an A_∞ morphism and recall the signs in the A_∞ identities in Appendix A, see (38).

Here we write down, for further reference, formulas for dp_2 and dp_3 , coming from (38).

$$dp_2(f, 1_x) + p_2(df, 1_x) = f - f \circ \text{id}_x, \quad dp_2(1_x, f) + p_2(1_x, df) = f - \text{id}_x \circ f \tag{3}$$

$$\begin{aligned}
 dp_3(f, g, 1_x) - (-1)^{|g|}p_3(df, g, 1_x) - p_3(f, dg, 1_x) &= f \circ p_2(g, 1_x) - p_2(f \circ g, 1_x) \\
 dp_3(f, 1_x, g) - (-1)^{|g|}p_3(df, 1_x, g) - p_3(f, 1_x, dg) &= -(-1)^{|g|}p_2(f, 1_x) \circ g + f \circ p_2(1_x, g) \\
 dp_3(1_x, f, g) - (-1)^{|g|}p_3(1_x, df, g) - p_3(1_x, f, dg) &= -(-1)^{|g|}p_2(1_x, f) \circ g + p_2(1_x, f \circ g) \\
 dp_3(1_x, 1_x, f) - p_3(1_x, 1_x, df) &= \text{id}_x \circ p_2(1_x, f) - (-1)^{|f|}p_2(1_x, 1_x) \circ f \\
 dp_3(1_x, f, 1_x) - p_3(1_x, df, 1_x) &= \text{id}_x \circ p_2(f, 1_x) - p_2(1_x, f) \circ \text{id}_x - p_2(f, 1_x) + p_2(1_x, f) \\
 dp_3(f, 1_x, 1_x) - p_3(df, 1_x, 1_x) &= f \circ p_2(1_x, 1_x) - p_2(f, 1_x) \circ \text{id}_x \\
 dp_3(1_x, 1_x, 1_x) &= \text{id}_x \circ p_2(1_x, 1_x) - p_2(1_x, 1_x) \circ \text{id}_x
 \end{aligned}
 \tag{4}$$

2.4. THE MONADIC STRUCTURE AND THE DG OPERAD \mathcal{O}' . Here we recall some results of [PS] which we use in this paper.

With the concept of a weakly unital dg category is associated a monad acting on the category of oriented dg graphs such that a weakly unital dg category amounts to the same thing as an algebra over this monad. This description is used in loc.cit. to prove that the category $\mathbb{C}_{dgwu}(\mathbb{k})$ is small complete and small cocomplete. The monad itself is defined via a non-symmetric dg operad called \mathcal{O}' . Here we briefly recall the corresponding definitions and results.

An oriented dg graph Γ over \mathbb{k} is given by a set V_Γ of vertices, and a complex $\Gamma(x, y)$ for any ordered pair $x, y \in V_\Gamma$. A morphism $F: \Gamma_1 \rightarrow \Gamma_2$ is given by a map of sets $F_V: V_{\Gamma_1} \rightarrow V_{\Gamma_2}$, and by a map of complexes $F_E: \Gamma_1(x, y) \rightarrow \Gamma_2(F_V(x), F_V(y))$, for any $x, y \in V_{\Gamma_1}$. We denote by $\mathbb{G}_{dg}(\mathbb{k})$ the category of the dg graphs over \mathbb{k} . A *unital* dg graph Γ is an oriented dg graph such that there is an element $\text{id}_x \in \Gamma(x, x)$, closed of degree 0, for any $x \in V_\Gamma$. A map of unital dg graphs is a map F of the underlying dg graphs such that $F(\text{id}_x) = \text{id}_{F(x)}$, for any $x \in V_\Gamma$. We denote by $\mathbb{G}_{dgu}(\mathbb{k})$ the category of unital dg graphs over \mathbb{k} .

There is a natural forgetful functor $U: \mathbb{C}_{dgwu}(\mathbb{k}) \rightarrow \mathbb{G}_{dgu}(\mathbb{k})$, where $U(C)$ is a graph Γ with $V_\Gamma = \text{Ob}(C)$, and $\Gamma(x, y) = C(x, y)$.

This functor admits a left adjoint $F: \mathbb{G}_{dgu}(\mathbb{k}) \rightarrow \mathbb{C}_{dgwu}(\mathbb{k})$. It is constructed via a dg operad \mathcal{O}' .

Consider the non- Σ the dg operad \mathcal{O}' defined as the quotient-operad of the free operad generated by the composition operations:

- (a) the composition operation $m \in \mathcal{O}'(2)^0$
- (b) $p_{n; i_1, \dots, i_k} \in \mathcal{O}'(n-k)^{-n+1}$, $0 \leq k \leq n$, $1 \leq i_1 < i_2 < \dots < i_k \leq n$, with the following meaning: For a weakly unital dg category C , the operation $p_{n; i_1, \dots, i_k}(f_1, \dots, f_{n-k})$ is defined as

$$p_n(f_1, \dots, f_{i_1-1}, \underset{i_1}{1_{x_1}}, f_{i_1}, \dots, f_{i_2-2}, \underset{i_2}{1_{x_2}}, f_{i_2-1}, \dots, \underset{i_k}{1_{x_k}}, f_{i_k-k+1}, \dots, f_{n-k}) \tag{5}$$

where by 1_{x_i} s are denoted the morphisms $1_{x_i} \in \mathbb{k}_C$ for the corresponding objects $x_i \in C$.

by the following relations:

- (i) the associativity of m , and $dm = 0$
 - (ii) $p_{n;i_1, \dots, i_k} = 0$ if $k = 0$
 - (iii) $p_{1;-} = \text{id}$
 - (iv) the A_∞ morphism relation for $dp_{n;i_1, \dots, i_k}$
- (6)

We use the notation $j = p_{1,1}$, the degree zero 0-ary operation generating the weak unit. It follows from (iv) that $dj = 0$. The reader is referred to [PS, Sect. 1.2.3] for the precise form of the relation in (iv). It expresses the relations like (3), (4) in the operadic terms, using the correspondence (5).

Morally, the dg operad \mathcal{O}' comprises all universal operations one can define on a weakly unital dg category.

Denote by $\mathcal{A}ssoc_+$ the operad of unital associative \mathbb{k} -algebras. We proved the following theorem:

2.5. THEOREM. *The natural map of dg operads $\mathcal{O}' \rightarrow \mathcal{A}ssoc_+$, sending all $p_{n;i_1, \dots, i_k}$, $n \geq 2$, to 0, sending $j = p_{1,1}$ to the 0-ary unit generating operation, and sending m to m , is a quasi-isomorphism.*

PROOF. The proof is a rather long and tricky computation with different spectral sequences, see [PS, Theorem 1.13]. ■

The left adjoint functor $F: \mathbb{G}_{dgu}(\mathbb{k}) \rightarrow \mathbb{C}_{dgu}(\mathbb{k})$ is defined in two steps, as follows. Given a unital dg graph Γ , consider the free \mathcal{O}' -algebra $T_{\mathcal{O}'}(\Gamma)$, generated by Γ (see [PS, Sect. 1.2.3]). It is a weakly unital dg category with objects V_Γ . The 0-ary operation j generates an element $j_x \in T_{\mathcal{O}'}(x, x)$, for any $x \in V_\Gamma$. After that, define $F(\Gamma)$ as the dg quotient-category

$$F(\Gamma) = T_{\mathcal{O}'}(\Gamma)/(j_x - \text{id}_x, x \in V_\Gamma) \tag{7}$$

In this way, we identify $\text{id}_x \in \Gamma(x, x)$ with the “weak unit” j_x generated by \mathcal{O}' .

One has:

2.6. PROPOSITION. *There is an adjunction:*

$$\mathbb{C}_{dgu}(F(\Gamma), C) \simeq \mathbb{G}_{dgu}(\Gamma, U(C)) \tag{8}$$

Note that for Γ a (non-unital) dg graph, one defines a unital dg graph Γ_+ , formally adding \mathbb{k}_x to $\Gamma(x, x)$, for any $x \in V_\Gamma$. Then

$$F(\Gamma_+) \simeq T_{\mathcal{O}'}(\Gamma)$$

3. The pretriangulated hull of a weakly unital dg category

Recall that the pretriangulated hull of a dg category C was introduced by Bondal-Kapranov [BK] (see also [Dr, 2.4, Remark]). A dg category C is *pretriangulated* if $H^0(C)$ is triangulated. Explicitly, it means that the functors $Z \rightarrow \underline{\text{Hom}}(C(Z, X) \xrightarrow{f} C(Z, Y))$ and $Z \rightarrow C(Z, X)[n]$ defined for any closed morphism $f: X \rightarrow Y$ in C and for any object $X \in C$, $n \in \mathbb{Z}$, correspondingly, are representable. In this case, the representing objects are $\text{Cone}(f)$ and $X[n]$.

The pretriangulated hull C^{pretr} of a dg category C is a pretriangulated dg category with a dg functor $C \rightarrow C^{\text{pretr}}$, which is universal for dg functors from C to pretriangulated dg categories [BK].

Explicitly, it is constructed as follows. An object of C^{pretr} is a “one-sided twisted complexes”, which are formal finite sums $(\oplus_{i=1}^n X_i[r_i], q)$ where q has components $q_{ij} \in C^{r_i-r_j+1}(X_i, X_j)$, which are zero for $i \geq j$, such that $dq + q^2 = 0$. Let $X = (\oplus X_i[r_i], q)$, $X' = (\oplus X'_i[r'_i], q')$ be two objects of C^{pretr} , a morphism $\phi \in C^{\text{pretr}}(X, X')$ of degree k is defined as the collection $\phi_{ij}: X_i[r_i] \rightarrow X'_j[r'_j]$ of degree d (in general, non-zero for any i, j), and $d\phi = d_C\phi + q' \circ \phi - (-1)^k \phi \circ q$. The composition is defined as the matrix product.

The dg functor $C \rightarrow C^{\text{pretr}}$ is defined on objects as $X \mapsto (X[0], q = 0)$, and on morphisms accordingly. We recall that $\text{Cone}(f) \in C^{\text{pretr}}$, $f: X \rightarrow Y$ a closed morphism in C , is defined as $\text{Cone}(f) = (X \oplus Y[-1], q = f)$ (that is, $q_{12} = f$, $q_{11} = q_{22} = q_{21} = 0$).

We want to define the pretriangulated hull of a *weakly unital* dg category, which is a weakly unital dg category as well. If we just repeated the definition given above, we would experience the following problem. Let $f \in C(x, y)$ be a closed morphism, we do *not* assume that $f \circ \text{id}_x = f$ or $\text{id}_y \circ f = f$. Defining $\text{Cone}(f) = (X \oplus Y[-1], f)$, there should be a weak identity morphism $\text{id}_{\text{Cone}(f)}$, which is a closed morphism of degree 0. A natural candidate is given by $\text{id}_X: X \rightarrow X, \text{id}_Y: Y \rightarrow Y$. But then $d(\text{id}_{\text{Cone}(f)}) = f \circ \text{id}_X - \text{id}_Y \circ f \neq 0$.

We remedy this problem as follows. For a closed morphism f in C , define $\text{Cone}(f) = (X \oplus Y[1], f)$ as above, but re-define $\text{id}_{\text{Cone}(f)}$. Namely, *define* $\text{id}_{\text{Cone}(f)}$ as having 3 non-zero components:

$$\begin{aligned} \text{id}_{\text{Cone}(f)} &= (\text{id}_X, \text{id}_Y, \varepsilon \in C^0(X, Y[1])) \\ &\text{where } \varepsilon = p_2(f, 1_x) - p_2(1_y, f) \end{aligned} \tag{9}$$

where p_2 is the second Taylor component of the A_∞ morphism $p: C \oplus \mathbb{k}_C \rightarrow C$, see Definition 2.1.

Then one has:

$$d(\text{id}_{\text{Cone}(f)}) = f \circ \text{id}_X - \text{id}_Y \circ f + d\varepsilon = 0 \tag{10}$$

(Recall that $dp_2(f, 1_x) = p_1(f) - f \circ \text{id}_X = f - f \circ \text{id}_X$, and similarly for $dp_2(1_y, f)$).

Thus, at the first step we define, inspired by this example, the identity morphism id_X , for $X = (\oplus X_i[r_i], q_{ij})$, and check that $d(\text{id}_X) = 0$. After that, we construct an A_∞ functor $P: C^{\text{pretr}} \oplus \mathbb{k}_{C^{\text{pretr}}} \rightarrow C^{\text{pretr}}$, making C^{pretr} a weakly unital dg category.

(We denote by p the structure A_∞ functor for C , and by P the structure A_∞ functor for C^{pretr}).

3.1. DEFINITION. [Pretriangulated hull of a wudg category, I] *Let \mathcal{C} be a wudg category. We define the underlying non-unital dg category of pretriangulated hull of \mathcal{C} as in the strictly unital case:*

- (a) *objects are formal expressions $(\bigoplus_{i=1}^n X_i[r_i], q_{ij})$, where $X_i \in \mathcal{C}$, $r_i \in \mathbb{Z}$, $q_{ij} \in \mathcal{C}^{1+r_j-r_i}(X_i, X_j) = \mathcal{C}^1(X_i[r_i], X_j[r_j])$ such that $q_{ij} = 0$ if $i \geq j$ and $dq + q \circ q = 0$,*
- (b) *the space of degree k morphisms $C^{\text{pretr}}(X, X')$, for*

$$X = (X_i[r_i], q_{ij}), X' = (X'_j[r'_j], q'_{ij}),$$

is defined as the space of matrices $\phi = (\phi_{ij} : C^k(X_i[r_i] \rightarrow X'_j[r'_j]))$, the composition is matrix multiplication and the differential is $d\phi := d_C\phi + q' \circ \phi - (-1)^k\phi \circ q$.

Now we define, for any object $X \in C^{\text{pretr}}$, an “identity” morphism id_X (which is required to be a closed morphism of degree 0), and construct an A_∞ morphism $P : C^{\text{pretr}} \oplus \mathbb{k}_{C^{\text{pretr}}} \rightarrow C^{\text{pretr}}$, making it a weakly unital dg category. In fact, we start with A_∞ morphism P , then $\text{id}_X := P_1(1_X)$.

Let X^0, \dots, X^n be objects of C^{pretr} , and let $\phi^i : X^{i-1} \rightarrow X^i$ be either a morphism in C^{pretr} or $1_{X^{i-1}}$ (in which case $X^i = X^{i-1}$).

We are going to define $P_n(\phi^n, \dots, \phi^1)$. Let us introduce some notations. We visualize the string

$$X^0 \xrightarrow{\phi^1} X^1 \xrightarrow{\phi^2} X^2 \rightarrow \dots \xrightarrow{\phi^n} X^n$$

as a planar diagram whose horizontal arrows are $q^i_{k\ell}$, where $X^i = (\bigoplus X^i_k[r^i_k], q^i_{k\ell})$, and whose other arrows are the components of ϕ^i , $i = 1, \dots, n$, see (11).

We refer to the arrows $q^i_{k\ell}$ as *horizontal*, and the other arrows, called *essential*, are either components of ϕ^i s or morphisms 1_X for some $X \in \mathcal{C}$.

Now we associate to any couple (X^0_a, X^n_b) of starting and ending objects, a set $\mathcal{P}ath_{sab}$ of **all** the possible paths from X^0_a to X^n_b , see (11), (12).

By definition, a *path* $\kappa \in \mathcal{P}ath_{sab}$ is a sequence of arrows $\kappa = (\kappa_1, \dots, \kappa_\ell)$, either horizontal or essential, such that (a) for any $1 \leq s \leq n$ there is exactly 1 essential arrow which is a components of ϕ^s , and these n essential arrows stand respecting the order, (b) the arrows between two successive essential arrows, which are components of ϕ^s and ϕ^{s+1} (here $\phi^s = 1_X$ is allowed), are horizontal arrows in X^s , which form a composable chain (there are allowed more than 1 arrows in this chain), (c) the first arrow starts at X^0_a , and the last one ends at X^n_b . It follows in particular that a path is represented by a composable chain of arrows.

If some $\phi^i = 1_{X^{i-1}}$, the corresponding arrow in $\phi^i_{k\ell} : X^{i-1}_k \rightarrow X^{i-1}_\ell$ is defined as $1_{X^{i-1}_k}$ for $k = \ell$, and 0 otherwise.

For example, in (11) the sequence $(q_{i\ell}, q_{\ell m}, \phi_{m\ell}, q'_{\ell m}, \phi'_{mm}, q''_{mj})$ is a path, and in (12)

the sequence $(\phi_{ii}, q'_{il}, q'_{lk}, \phi'_{ki}, q''_{ij})$ is a path (here for both diagrams $n = 2$).

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & X_i^0 & \xrightarrow{q_{i\ell}} & X_l^0 & \xrightarrow{q_{\ell m}} & X_m^0 & \longrightarrow & \dots X_j^0 & \longrightarrow & \dots \\
 & & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \\
 \dots & \longrightarrow & X_i^1 & \longrightarrow & X_l^1 & \xrightarrow{q'_{\ell m}} & X_m^1 & \longrightarrow & X_j^1 & \longrightarrow & \dots \\
 & & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \\
 \dots & \longrightarrow & X_i^2 & \longrightarrow & X_l^2 & \longrightarrow & X_m^2 & \xrightarrow{q''_{mj}} & X_j^2 & \longrightarrow & \dots
 \end{array} \tag{11}$$

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & X_i^0 & \longrightarrow & X_l^0 & \longrightarrow & X_m^0 & \longrightarrow & X_j^0 & \longrightarrow & \dots \\
 & & \downarrow \phi_{ii} & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \\
 \dots & \longrightarrow & X_i^1 & \xrightarrow{q_{i\ell}} & X_l^1 & \longrightarrow & X_m^1 & \longrightarrow & X_j^1 & \longrightarrow & \dots \\
 & & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \\
 \dots & \longrightarrow & X_i^2 & \longrightarrow & X_l^2 & \longrightarrow & X_m^2 & \longrightarrow & X_j^2 & \longrightarrow & \dots
 \end{array} \tag{12}$$

Below we assume for $n \geq 2$ that at least one of morphisms $\phi^i: X^{i-1} \rightarrow X^i$ is $1_{X^{i-1}}$; otherwise, (13) below gives 0.

Define

$$P_n^{ij}(\phi_n, \dots, \phi_1) := \sum_{\kappa \in \text{Paths}_{ij}} (-1)^{|\kappa|} p_l(\kappa_l, \dots, \kappa_1) \tag{13}$$

(Recall that p denotes the structure A_∞ morphism for C).

To define the integer $|\kappa|$, we introduce some notations. Let $\kappa = (\kappa_1, \dots, \kappa_\ell)$, and let the n arrows $\kappa_{d_1}, \dots, \kappa_{d_n}$ be essential. Assume that κ_{d_s} is an arrow in $\mathcal{C}(X_{a_s}^{s-1}[r^s], X_{b_s}^s[r^{s+1}])$. Define $t_s = r^{s+1} - r^s$ (we set $t_s = 0$ if $\kappa_{d_s} = 1_X$).

The integer $|\kappa|$ is given by

$$|\kappa| = \sum_{s=1}^n (\deg \phi^s + t_s + 1) N_s \tag{14}$$

where N_s is the number of the horizontal arrows standing *leftwards* to the s -th essential arrow κ_{d_s} in the sequence $(\kappa_1, \dots, \kappa_\ell)$.

3.2. LEMMA. *The maps $P_n^{ij}(\phi_n, \dots, \phi_1)$ are homogeneous of degree $\sum \deg \phi_i - n + 1$. Thus, they are the components of a morphism $P_n(\phi_n, \dots, \phi_1): X^0 \rightarrow X^n$ of degree $\sum \deg \phi_i - n + 1$ in the category C^{pretr} .*

PROOF. Let $\kappa = (\kappa_1, \dots, \kappa_\ell) \in \mathcal{P}aths_{ij}$, we have to compute the degree of $p_\ell(\kappa_n, \dots, \kappa_1)$. One has:

$$\deg p_\ell(\kappa_\ell, \dots, \kappa_1) = \sum_{r=1}^{\ell} \deg \kappa_r - \ell + 1$$

Among $\kappa_1, \dots, \kappa_\ell$ exactly n arrows are ϕ_{st}^i s, the remaining $\ell - n$ are q_{st}^i and have degree 1 in C^{pretr} . On the other hand, $\deg \phi_{st}^i = \deg \phi^i$ and does not depend on s, t . Therefore,

$$\deg p_\ell(\kappa_\ell, \dots, \kappa_1) = \sum_{\kappa_r \neq q_{st}^j} \deg \kappa_r - n + 1 = \sum_{r=1}^n \deg \phi_r - n + 1$$

■

3.3. PROPOSITION. *Let C be a weakly unital dg category, C^{pretr} the non-unital dg category from Definition 3.1. Taken for all $n \geq 1$ and all ϕ_1, \dots, ϕ_n , the morphisms $P_n(\phi_n, \dots, \phi_1)$ are the Taylor components of an A_∞ morphism $P: C^{\text{pretr}} \oplus \mathbb{k}_{C^{\text{pretr}}} \rightarrow C^{\text{pretr}}$, making C^{pretr} a weakly unital dg category, with $\text{id}_X := P_1(1_X)$.*

We prove this Proposition in Section 6.

3.4. DEFINITION. [Pretriangulated hull of a wudg category, II] *The pretriangulated hull C^{pretr} of a weakly unital dg category C is the non-unital dg category C^{pretr} (see Definition 3.1) with the weakly unital structure given in Proposition 3.3.*

It is instructive to unwind the definition $\text{id}_X = P_1(1_X)$ and get an explicit formula $\text{id}_X, X \in C^{\text{pretr}}$.

Let $X = (\oplus X_i[r_i], q_{ij}) \in C^{\text{pretr}}$. We want to find the (ij) -component $(\text{id}_X)_{ij}$. Let $i \leq j$. Define

$$\begin{aligned} (\text{id}_X)_{ij} &= \sum_{i=\ell_0 < \ell_1 < \dots < \ell_k=j} \sum_{r=0}^k \\ &(-1)^r p_k(q_{\ell_{k-1}j}, q_{\ell_{k-2}\ell_{k-1}}, \dots, q_{\ell_r\ell_{r+1}}, 1_{X_{i_r}}, q_{\ell_{r-1}\ell_r}, q_{\ell_{r-2}\ell_{r-1}}, \dots, q_{i\ell_1}) \end{aligned} \tag{15}$$

Then

$$(\text{id}_X)_{ij} = \begin{cases} \text{the rhs of (15)} & i \leq j \\ 0 & i > j \end{cases} \tag{16}$$

The reader easily checks that for the case $\text{id}_{\text{Cone}(f)}$ (15) gives (9).

4. A Closed Model Structure on $\mathbb{C}_{dgwu}(\mathbb{k})$

In this section we provide a cofibrantly generated Quillen model structure on $\mathbb{C}_{dgwu}(\mathbb{k})$. The reader is referred to [Ho], [Hi], [GS] for general introduction to (cofibrantly generated) closed model categories. A simpler counterpart of the material of this Section is our treatment [PS] of the closed model structure on the category $\mathbb{C}_{dgwu}^0(\mathbb{k})$. However, the

passage from $\mathbb{C}_{dgwu}^0(\mathbb{k})$ to $\mathbb{C}_{dgwu}(\mathbb{k})$ requires several essentially new ideas, such as the weakly unital pretriangulated hull, introduced in Section 3, and a “fattened version” \mathcal{K} of the Kontsevich dg category \mathcal{K} (see Section 4.4).

Define *weak equivalences* W in $\mathbb{C}_{dgwu}(\mathbb{k})$ as the weakly unital dg functors $F : \mathcal{C} \rightarrow \mathcal{D}$ such that the following two conditions hold:

- (W1) for any two objects $x, y \in \mathcal{C}$, the map of complexes $\mathcal{C}(x, y) \rightarrow \mathcal{D}(Fx, Fy)$ is a quasi-isomorphism of complexes,
- (W2) the functor $H^0(F) : H^0(\mathcal{C}) \rightarrow H^0(\mathcal{D})$ is an equivalence of \mathbb{k} -linear categories.

Remark that for a weakly unital dg category \mathcal{C} , the category $H^0(\mathcal{C})$ is strictly unital and the functor $H^0(F)$ is well-defined.

Define *fibrations* Fib in $\mathbb{C}_{dgwu}(\mathbb{k})$ as the weakly unital dg functors $F : \mathcal{C} \rightarrow \mathcal{D}$ such that the following two conditions hold:

- (F1) for any two objects $x, y \in \mathcal{C}$, the map of complexes $\mathcal{C}(x, y) \rightarrow \mathcal{D}(Fx, Fy)$ is component-wise surjective,
- (F2) for any $x \in \mathcal{C}$ and a closed degree 0 arrow $g : Fx \mapsto z$ in \mathcal{D} (z not necessarily in the image of F), such that g becomes an isomorphism in $H^0(\mathcal{D})$, there is an object $y \in \mathcal{C}$ and a closed degree 0 arrow $f : x \mapsto y$ inducing an isomorphism in $H^0(\mathcal{C})$ and such that $F(f) = g$.

We define also a class *Surj* of maps in $\mathbb{C}_{dgwu}(\mathbb{k})$ as follows: a weakly unital dg functor $F : \mathcal{C} \rightarrow \mathcal{D}$ belongs to *Surj* if F is surjective on objects and if (F1) holds.

4.1. LEMMA. *A weakly unital dg functor $F : \mathcal{C} \rightarrow \mathcal{D}$ belongs to $Fib \cap W$ if and only if it belongs to $Surj \cap (W1)$.*

PROOF. A proof can be found in [PS, Lemma 2.1]. ■

4.2. LEMMA. *Let X be a weakly unital dg category, $x \in X$ an object. Suppose there are two degree -1 maps $h_1, h_2 \in X^{-1}(x, x)$ such that $dh_i = id_x, i = 1, 2$. Then there is $t \in X^{-2}(x, x)$ such that $dt = h_1 - h_2$.*

PROOF. Consider $t' = h_1h_2$. We find (using (3)):

$$\begin{aligned} dt' &= id_x \circ h_2 - h_1 \circ id_x \\ &= -dp_2(1, h_2) + h_2 - p_2(1, id_x) + dp_2(h_1, 1) - h_1 + p_2(id_x, 1) \end{aligned} \tag{17}$$

If we manage to prove that $p_2(id, 1) - p_2(1, id)$ is a boundary, we’re done. Consider

$$\begin{aligned} H &:= p_2(p_2(id, 1), 1) + p_2(1, p_2(1, id)) - p_3(1, id, 1) + p_3(id, id, 1) + p_3(1, id, id) \\ &\quad - p_3(1, 1, id) - p_3(id, 1, 1) + p_3(1, 1, 1) \end{aligned} \tag{18}$$

We compute dH using (3) and (4), the differential of each particular summand in (18) is displayed as [...]:

$$\begin{aligned}
 dH := & \left[\underbrace{p_2(id, 1) \circ id}_{\dots\dots\dots} + \underbrace{p_2(id \circ id, 1)} \right] + \left[\underbrace{id \circ p_2(1, id)}_{\dots\dots\dots} - \underbrace{p_2(1, id \circ id)}_{\dots\dots\dots} \right] + \left[\underbrace{p_2(1, id) \circ id}_{\dots\dots\dots} \right. \\
 & \left. - \underbrace{id \circ p_2(id, 1)}_{\dots\dots\dots} + \boxed{p_2(id, 1) - p_2(1, id)} \right] + \left[- \underbrace{p_2(id \circ id, 1)}_{\dots\dots\dots} + \underbrace{id \circ p_2(id, 1)}_{\dots\dots\dots} \right] + \\
 & \left[\underbrace{p_2(1, id \circ id)}_{\dots\dots\dots} - \underbrace{p_2(1, id) \circ id}_{\dots\dots\dots} \right] + \left[\underbrace{p_2(1, 1) \circ id}_{\dots\dots\dots} - \underbrace{id \circ p_2(1, id)}_{\dots\dots\dots} \right] + \\
 & \left[- \underbrace{id \circ p_2(1, 1)}_{\dots\dots\dots} + \underbrace{p_2(id, 1) \circ id}_{\dots\dots\dots} \right] + \left[\underbrace{id \circ p_2(1, 1)}_{\dots\dots\dots} - \underbrace{p_2(1, 1) \circ id}_{\dots\dots\dots} \right] = \\
 & p_2(id, 1) - p_2(1, id).
 \end{aligned}$$

Therefore $t := -t' - p_2(1, h_2) + p_2(h_1, 1) + H \in X^{-2}(x, x)$ is such that $dt = h_1 - h_2$. ■

4.3. REMARK. Lemma 4.2 is a refined version of Lemma 2.8 and discussion below it of [PS]. When $id_x \circ id_x = id_x$ is not assumed, the computation turns out to be more tricky, as we've seen in Lemma 4.2.

4.4. THE WEAKLY UNITAL DG CATEGORY \mathcal{K}' . The wu dg category \mathcal{K}' introduced below is a weakly unital counterpart of the dg category \mathcal{K} , due to Kontsevich in [K1, Lecture 6], and subsequently used by Tabuada in his closed model structure on $\mathbb{C}_{dg}(\mathbb{k})$ [Tab1].

Recall here the definition.

The dg category \mathcal{K} is the strictly unital dg category with two objects 0 and 1, and freely generated by $f \in \mathcal{K}^0(0, 1), g \in \mathcal{K}^0(1, 0), h_0 \in \mathcal{K}^{-1}(0, 0), h_1 \in \mathcal{K}^{-1}(1, 1), r \in \mathcal{K}^{-2}(0, 1)$, whose differentials are

$$df = dg = 0, dh_0 = g \circ f - id_0, dh_1 = f \circ g - id_1, dr = h_1 \circ f - f \circ h_0 \tag{19}$$

Denote by I_2 the \mathbb{k} -linear envelope of the ordinary category with two objects 0 and 1, and with a unique morphism (including the identity one) between any ordered pair of objects. There is a dg functor $p_{\mathcal{K}}: \mathcal{K} \rightarrow I_2$, which is the identity map on the objects, and sends h_1, h_2, r to 0.

The following well-known result says that \mathcal{K} is a semi-free resolution of I_2 :

4.5. LEMMA. *The dg functor $p_{\mathcal{K}}: \mathcal{K} \rightarrow I_2$ is a quasi-equivalence.*

The proof can be found in [Dr, 3.7].

4.6. DEFINITION. *Denote by \mathcal{K}' the weakly unital dg category with two objects 0 and 1, whose morphisms are freely generated by the following morphisms:*

- a morphism $f \in (\mathcal{K}')^0(0, 1)$,
- a morphism $g \in (\mathcal{K}')^0(1, 0)$,
- a morphism $h_0 \in (\mathcal{K}')^{-1}(0, 0)$,

- a morphism $h_1 \in (\mathcal{K}')^{-1}(1, 1)$,
- a degree -2 morphism $r \in (\mathcal{K}')^{-2}(0, 1)$

whose differentials are given as

$$\begin{aligned} df &= dg = 0 \\ dh_0 &= g \circ f - \text{id}_0, \quad dh_1 = f \circ g - \text{id}_1 \\ dr &= h_1 \circ f - f \circ h_0 + p_2(1, f) - p_2(f, 1) \end{aligned} \tag{20}$$

A version of a lemma in [K1, Lecture 6] holds as well in the setting of weakly unital dg categories:

4.7. LEMMA. *Let C be a weakly unital dg category, and $\xi \in C^0(x, y)$ be a closed degree 0 morphism, such that $[\xi] \in H^0(C)$ is a homotopy equivalence. Then there is a weakly unital dg functor $F : \mathcal{K}' \rightarrow C$, such that $F(f) = \xi$.*

PROOF. By definition of being a homotopy equivalence, there exist $\eta \in C^0(y, x)$, $h_x \in C^{-1}(x, x)$ and $h_y \in C^{-1}(y, y)$ such that:

$$\begin{aligned} dh_x &= \eta \circ \xi - \text{id}_x \\ dh_y &= \xi \circ \eta - \text{id}_y \end{aligned} \tag{21}$$

Now we are looking for a morphism $r \in C^{-2}(x, y)$ such that $dr = h_y \circ \xi - \xi \circ h_x + p_2(1, \xi) - p_2(\xi, 1)$. We define $A := h_y \circ \xi - \xi \circ h_x + p_2(1, \xi) - p_2(\xi, 1)$. Clearly $dA = 0$.

Then we take $h'_y := h_y - A \circ \eta$ and $r := A \circ h_x - p_2(A, 1)$, so that we easily get:

$$dh'_y = dh_y - dA\eta + Ad\eta = \xi\eta - \text{id}_y. \tag{22}$$

and also

$$\begin{aligned} dr &= dA \circ h_x - A \circ dh_x - A \circ \text{id}_x + A - p_2(dA, 1) \\ &= -A \circ (\eta\xi - \text{id}_x) - A \circ \text{id}_x + A \\ &= -A \circ \eta \circ \xi + h_y \circ \xi - \xi \circ h_x + p_2(1, \xi) - p_2(\xi, 1) \\ &= (h_y - A \circ \eta) \circ \xi - \xi \circ h_x + p_2(1, \xi) - p_2(\xi, 1) \\ &= h'_y \circ \xi - \xi \circ h_x + p_2(1, \xi) - p_2(\xi, 1). \end{aligned} \tag{23}$$

We are done. ■

We prove a lemma which we will be used later in the proof of Theorem 4.9 and (implicitly) in Theorem 4.17:

4.8. LEMMA. *Let C be a weakly unital dg category. There is a bijection between the set of weakly unital dg functors from \mathcal{K}' to C and the set of pairs (ξ, h) , where $\xi \in C^0(x, y)$ is a closed morphism and h is a contraction of $\text{Cone}(\xi)$ in C^{pretr} .*

PROOF. A *wu dg* functor $F: \mathcal{K}' \rightarrow \mathcal{C}$ amounts to the following morphisms in \mathcal{C} : $\xi = F(f), \eta = F(g), h_{11} = F(h_0), h_{22} = F(h_1), h_{12} = F(r)$ such that

$$\begin{aligned} d(\xi) = 0, \quad d\eta = 0, \quad dh_{11} = \eta \circ \xi - id_x, \quad dh_{22} = \xi \circ \eta - id_y, \\ dh_{12} = h_{22} \circ \xi - \xi \circ h_{11} + p_2(1, \xi) - p_2(\xi, 1) \end{aligned} \tag{24}$$

A contraction to $id_{Cone(\xi)}$ (see (9)) is the datum of a morphism $H: Cone(\xi) \rightarrow Cone(\xi)$ of degree -1 such that

$$dH = Id_{Cone(\xi)}, \tag{25}$$

Then

$$H = (h_{11} \in C^{-1}(X, X), h_{22} \in C^{-1}(Y, Y), h_{12} \in C^{-2}(X, Y), h_{21} \in C^0(Y, X))$$

as in the diagram below

$$\begin{array}{ccc} X & \xrightarrow{\xi} & Y \\ & \searrow^{h_{12}} & \downarrow^{h_{22}} \\ h_{11} \downarrow & & \\ X & \xrightarrow{\xi} & Y \\ & \swarrow_{h_{21}} & \end{array} \tag{26}$$

We see by a direct calculation that 24 is equivalent to 25. ■

4.9. PROPOSITION. *The *wu dg* category \mathcal{K}' has the same homotopy type as the Kontsevich dg category \mathcal{K} . More precisely, regarding \mathcal{K} as an object in $\mathcal{C}_{dgwu}(\mathbb{k})$, the natural projection $p: \mathcal{K}' \rightarrow \mathcal{K}$, sending all $p_n(-)$, $n \geq 2$, to 0, is a weak equivalence.*

PROOF. Consider ascending filtrations $\{\Phi_i(a, b)\}_{i \geq 0}$ of $\mathcal{K}'(a, b)$ and $\{F_i(a, b)\}_{i \geq 0}$ of $\mathcal{K}(a, b)$, $a, b \in \{0, 1\}$, such that $p(\Phi_i(a, b)) \subset F_i(a, b)$, $i \geq 0$. We prove that the corresponding spectral sequences converge, and that the map p induces an isomorphism in the E_1 sheets. The result will follow from the latter statement.

Define $F_i(a, b)$ as the dg vector space generated by all monomials with $\leq i$ factors r . Define $\Phi_i(a, b)$ similarly, but we count all occurrences of r in expressions $p_j(\dots, r, \dots)$ as a “factor r ”. It is clear that $d(F_i(a, b)) \subset F_i(a, b)$ and $d(\Phi_i(a, b)) \subset \Phi_i(a, b)$, and that $p(\Phi_i(a, b)) \subset F_i(a, b)$.

Also, it is clear that both spectral sequences converge, by dimensional reasons (the spectral sequences live in the quarter “ $x \leq 0, y \leq 0$ ”).

We have:

4.10. LEMMA. *The map p induces an isomorphism in the E_1 sheets.*

PROOF. For both cases, the differential in E_0 is the same as it would be if $dr = 0$. Therefore, to compute E_1 we assume that $dr = 0$ for both cases.

Denote by $\tilde{\mathcal{K}}'$ (corresp. $\tilde{\mathcal{K}}$) the semi-free *wu dg* category (corresp., the semi-free unital dg category) with two objects $\{0, 1\}$, the generators f, g, h_0, h_1, r , as in (19), (20), and in

which the differential of the generators is given by *the same* formulas:

$$\begin{aligned} df &= dg = 0 \\ dh_0 &= g \circ f - \text{id}_0, \quad dh_1 = f \circ g - \text{id}_1 \\ dr &= 0 \end{aligned} \tag{27}$$

Now the statement follows from the fact that the projection of dg operads $\mathcal{O}' \rightarrow \mathcal{O}$ is a quasi-isomorphism in any arity, see [PS, Section 4] (here \mathcal{O} is a dg operad, defined analogously to \mathcal{O}' , see loc.cit.). ■

4.11. **REMARK.** The argument employed in the proof of Lemma 4.10 can not be used directly for $p: \mathcal{K}' \rightarrow \mathcal{K}$ (without any spectral sequence argument), because dr is given by *different* formulas in (19) and (20). More precisely, the equation for dr for \mathcal{K}' is a *deformation* of that for \mathcal{K} . Consequently, it does not follow directly from the quasi-isomorphism $\mathcal{O}' \rightarrow \mathcal{O}$ that $p: \mathcal{K}'(a, b) \rightarrow \mathcal{K}(a, b)$ is a quasi-isomorphism.

Recall that I_2 is the \mathbb{k} -linear envelope of the ordinary category with two objects, 0 and 1, and having exactly 1 morphism between any ordered pair of objects. Recall that Lemma 4.5 says that the projection $p_{\mathcal{K}}: \mathcal{K} \rightarrow I_2$ is a weak equivalence.

4.12. **COROLLARY.** *The natural projection $\mathcal{K}' \rightarrow I_2$, equal to the composition $\mathcal{K}' \xrightarrow{p} \mathcal{K} \xrightarrow{p_{\mathcal{K}}} I_2$, is a weak equivalence.*

PROOF. The dg functor p is a weak equivalence by Proposition 4.9. The dg functor $p_{\mathcal{K}}: \mathcal{K} \rightarrow I_2$, is a weak equivalence by Lemma 4.5. ■

4.13. **THE SETS I AND J .** Denote by $D(n)$ the complex $0 \rightarrow \mathbb{k}[n] \xrightarrow{\text{id}} \mathbb{k}[n-1] \rightarrow 0$, it is $D(n) = \text{Cone}(\text{id}: \mathbb{k}[n] \rightarrow \mathbb{k}[n])$.

Denote $S(n-1) = \mathbb{k}[n-1]$. Consider the natural imbedding $i: S(n-1) \rightarrow D(n)$ of complexes.

Denote by A the weakly unital dg category with a single object 0 and generated (over the dg operad \mathcal{O}') by id_0 . Denote by κ the weakly unital dg functor

$$\kappa: A \rightarrow \mathcal{K}',$$

sending 0 to 0. It follows from Corollary 4.12 that κ is a weak equivalence.

Denote by \mathcal{B} the weakly unital dg category with two objects 0 and 1 and generated over \mathcal{O}' by morphisms id_0 and id_1 .

Let $P(n)$ be the dg graph with two objects 0 and 1, and with morphisms $P(n)(0, 1) = D(n)$, $P(n)(0, 0) = 0$, $P(n)(1, 1) = 0$, $P(n)(1, 0) = 0$. Denote by $\mathcal{P}(n)$ the weakly unital dg category generated by $P(n)$: $\mathcal{P}(n) = \text{FU}(P(n))$.

Denote by $\alpha(n)$ the weakly unital dg functor

$$\alpha(n): \mathcal{B} \rightarrow \mathcal{P}(n),$$

sending 0 to 0 and 1 to 1.

Let $C(n)$ be the dg graph with two objects 0 and 1, and with morphisms $C(n)(0, 1) = S(n-1)$, $C(n)(0, 0) = 0$, $C(n)(1, 1) = 0$, $C(n)(1, 0) = 0$. Denote by $\mathcal{C}(n)$ the weakly unital dg category generated by $C(n)$: $\mathcal{C}(n) := FU(C(n))$.

Let $b(n) : C(n) \rightarrow P(n)$ be a map of dg graphs sending 0 to 0, 1 to 1, and such that $S(n-1) = C(n)(0, 1) \xrightarrow{i} P(n)(0, 1) = D(n)$ is the imbedding i . Denote by $\beta(n)$ the weakly unital dg functor

$$\beta(n) := FU(b(n)) : \mathcal{C}(n) \xrightarrow{i} \mathcal{P}(n).$$

Denote by Q the natural weakly unital dg functor

$$Q : \emptyset \rightarrow \mathcal{A}.$$

Let I be a set of morphisms in $\mathbb{C}_{dgwu}(\mathbb{k})$ which comprises the weakly unital dg functors Q and $\beta(n)$, $n \in \mathbb{Z}$.

Let J be a set of morphisms in $\mathbb{C}_{dgwu}(\mathbb{k})$ which comprises κ and $\alpha(n)$, $n \in \mathbb{Z}$.

The set I and J are referred to as the sets of *generating cofibrations* and of *generating acyclic cofibrations*, correspondingly.

4.14. LEMMA. *A weakly unital dg functor $\phi : \mathcal{C} \rightarrow \mathcal{D}$ has RLP with respect to all $\alpha(n)$, $n \in \mathbb{Z}$ if and only if ϕ obeys (F1). A weakly unital dg functor $\phi : \mathcal{C} \rightarrow \mathcal{D}$ has RLP with respect to all $\beta(n)$, $n \in \mathbb{Z}$ if and only if ϕ obeys $(F1) \cap (W1)$.*

PROOF. The proof is straightforward and can be found in [PS, Sec.2.1.3]. ■

Recall the standard terminology (conventional for the theory of closed model categories): a dg functor $\phi : \mathcal{C} \rightarrow \mathcal{D}$ belongs to *I-inj* (resp., to *J-inj*) if it has the RLP with respect to all morphisms in I (resp., in J).

4.15. PROPOSITION. *One has*

$$I\text{-inj} = \text{Surj} \cap (W1) = J\text{-inj} \cap W$$

PROOF. Thanks to Lemma 4.14, the first equality follows from the fact that a dg functor $\phi : \mathcal{C} \rightarrow \mathcal{D}$ has the RLP for Q if and only if it is surjective on objects, which is straightforward.

The second equality is far more sophisticated, and its proof is based on the following lemma.²

4.16. LEMMA. *One has $Fib = J\text{-inj}$.*

²Lemma 4.16 is one of the most subtle places in our constructions; in particular, the theory of weakly unital pre-triangulated hull developed in Section 3, and Lemma 4.2, were designed especially for its proof.

PROOF. The inclusion $J\text{-inj} \subseteq \text{Fib}$ follows from Lemma 4.7.

In order to prove the inclusion $\text{Fib} \subseteq J\text{-inj}$ consider $\phi : \mathcal{C} \rightarrow \mathcal{D}$ in Fib . Axiom (F1) is equivalent to the *RLP* with respect to $\alpha(n), n \in \mathbb{Z}$, thence we only need to prove the *RLP* with respect to κ for ϕ . We are given a weakly unital dg functor $F : \mathcal{K}' \rightarrow \mathcal{D}$. We can apply (F2) to $\xi = F(f) \in \mathcal{D}^0(\phi(x), z)$, and so we get a morphism $\eta \in \mathcal{C}^0(x, y)$ which is a homotopy equivalence and $\phi(\eta) = \xi, \phi(y) = z$. (Recall that f, g, h_0, h_1, r are generators for \mathcal{K}' , see (20)). We should construct a weakly unital dg functor $\hat{F} : \mathcal{K}' \rightarrow \mathcal{C}$ such that $\phi \circ \hat{F} = F$ and $\hat{F}(f) = \eta$. By Lemma 4.8 having a weakly unital dg functor $F : \mathcal{K}' \rightarrow \mathcal{D}, F(f) = \xi$ is equivalent to having a contraction of $\text{Cone}(\xi)$ in $\mathcal{D}^{\text{pretr}}$, i.e. we have $h \in \mathcal{D}^{\text{pretr}}(\text{Cone}(\xi), \text{Cone}(\xi))$ such that $dh = id_{\text{Cone}(\xi)}$. By (F2) we know that $\text{Cone}(\eta)$ is also contractible, so we also have a morphism $\tilde{h}_1 \in \mathcal{C}^{\text{pretr}}(\text{Cone}(\eta), \text{Cone}(\eta))$, such that $d\tilde{h}_1 = id_{\text{Cone}(\eta)}$. Even though we don't know whether $\phi^{\text{pretr}}(\tilde{h}_1) = h$, we still have $d\phi^{\text{pretr}}(\tilde{h}_1) = id_{\text{Cone}(\xi)}$.

By Lemma 4.2, one has $\phi^{\text{pretr}}(\tilde{h}_1) - h = dt$. By (F1), we find a lift \tilde{t} of t , and set $\tilde{h} := \tilde{h}_1 - d\tilde{t}$. One clearly has $d\tilde{h} = id_{\text{Cone}(\eta)}$ and $\phi(\tilde{h}) = h$. This gives the desired lift $\hat{F} : \mathcal{K}' \rightarrow \mathcal{C}$ such that $\phi \circ \hat{F} = F$, by Lemma 4.8. ■

Now we have $J\text{-inj} \cap W = \text{Fib} \cap W = \text{Surj} \cap (W1)$, and we're done. ■

The following theorem is one of our main results:

4.17. THEOREM. *The category*

$$\mathbb{C}_{dgwu}(\mathbb{k})$$

admits a cofibrantly generated closed model structure whose weak equivalences and fibrations are as above, and whose sets of generating cofibrations and generating acyclic cofibrations are I and J .

4.18. PROOF OF THEOREM 4.17. We refer the reader to standard textbooks on closed model categories for the notations *I-cell, J-cell, I-cof, J-cof*, see e.g. [Ho, 2.1.3], [Hi, 11.1]. Recall that $I\text{-cell} \subset I\text{-cof}$ and $J\text{-cell} \subset J\text{-cof}$.

Recall [Ho, Th. 2.1.19] which the proof is based on:

4.19. THEOREM. *Let C be a small complete and cocomplete category. Suppose that W is a subcategory of C , and I and J are sets of maps. Assume that the following conditions hold:*

1. *the subcategory W has 2-out of-3 property and is closed under retracts,*
2. *the domains of I are small relative to $I\text{-cell}$,*
3. *the domains of J are small relative to $J\text{-cell}$,*
4. *$J\text{-cell} \subset W \cap I\text{-cof}$,*
5. *$I\text{-inj} = W \cap J\text{-inj}$.*

Then there is a cofibrantly generated closed model structure on C , for which the morphisms W of \mathcal{W} are weak equivalences, I are generating cofibrations, J are generating acyclic cofibrations. Its fibrations are defined as J -inj.

PROOF OF THEOREM 4.17. The category $\mathbb{C}_{dgwu}(\mathbb{k})$ is small complete and small cocomplete by [PS, Theorem 1.19]. The conditions (1) – (3) are clear. Condition (5) was proved in Proposition 4.15. It follows from (5) that I -inj $\subset J$ -inj, and so J -cof $\subset I$ -cof. It only remains to prove that J -cell $\subseteq W$.

PROOF OF J -cell $\subseteq W$. We have to prove that the push-out of a morphism in J is a weak equivalence. We consider two cases: when the morphism in J is $\alpha(n)$, $n \in \mathbb{Z}$, and when it is κ .

First case.

The first case we need to consider is shown in the push-out diagram:

$$\begin{array}{ccc} \mathcal{B} & \xrightarrow{g} & \mathcal{X} \\ \alpha(n) \downarrow & & \downarrow f \\ \mathcal{P}(n) & \longrightarrow & \mathcal{Y} \end{array} \tag{28}$$

where g is an arbitrary map. We need to prove that f is a weak equivalence.

Clearly $Ob(\mathcal{X}) = Ob(\mathcal{Y})$, and f acts by the identity maps on the objects. We are left to show that, for any objects $a, b \in Ob(\mathcal{X})$, the map of complexes $f(a, b) : \mathcal{X}(a, b) \rightarrow \mathcal{Y}(a, b)$ is a quasi-isomorphism. For objects $0, 1$ in $Ob(\mathcal{B})$, let denote by $u = g(0)$ and $v = g(1)$. By Proposition C.1, one has the following description for the hom-complexes of \mathcal{Y} :

$$\begin{aligned} \mathcal{Y}(a, b) := & \mathcal{X}(a, b) \bigoplus \mathcal{O}'(3) \otimes \mathcal{X}(v, b) \otimes D(n) \otimes \mathcal{X}(a, u) \\ & \bigoplus \mathcal{O}'(5) \otimes \mathcal{X}(v, b) \otimes D(n) \otimes \mathcal{X}(v, u) \otimes D(n) \otimes \mathcal{X}(a, u) \bigoplus \dots \end{aligned} \tag{29}$$

The map $f(a, b)$ sends $\mathcal{X}(a, b)$ to the first summand. The other summands have trivial cohomology by the Künneth formula, since the acyclicity of $D(n)$.

Second case.

As the second case we consider the following push-out diagram:

$$\begin{array}{ccc} A & \xrightarrow{h} & \mathcal{X} \\ \kappa \downarrow & & \downarrow f \\ \mathcal{K}' & \longrightarrow & \mathcal{Y} \end{array} \tag{30}$$

where h is an arbitrary map.

One has $Ob(\mathcal{Y}) = Ob(\mathcal{X}) \sqcup 1_{\mathcal{K}'}$. It is clear that $H^0(f)$ is essentially surjective. One has to prove that the f is locally a quasi-isomorphism: $f(a, b) : \mathcal{X}(a, b) \rightarrow \mathcal{Y}(a, b)$, $\forall a, b \neq 1_{\mathcal{K}'}$.

Denote $h(0_A) = u$.

By Theorem 4.9, we know that \mathcal{K}' is a resolution of the \mathbb{k} -linear envelope of the ordinary category with two objects $0, 1$ and with only one morphism between any pair of objects. In particular, $\mathcal{K}'(0, 0)$ is quasi-isomorphic to $\mathbb{k}[0]$. Therefore

$$\bar{\mathcal{K}}' := \mathcal{K}'(0, 0)/\mathbb{k}[0] \tag{31}$$

is a complex acyclic in all degrees.

By Proposition C.1, we have:

$$\begin{aligned} \mathcal{Y}(a, b) := & \mathcal{X}(a, b) \bigoplus \mathcal{O}(3) \otimes \mathcal{X}(u, b) \otimes \bar{\mathcal{K}}' \otimes \mathcal{X}(a, u) \\ & \bigoplus \mathcal{O}(5) \otimes \mathcal{X}(u, b) \otimes \bar{\mathcal{K}}' \otimes \mathcal{X}(u, u) \otimes \bar{\mathcal{K}}' \otimes \mathcal{X}(a, u) \bigoplus \dots \end{aligned} \tag{32}$$

It is a direct sum of complexes, among which all but the first one are acyclic, due to the acyclicity of $\bar{\mathcal{K}}'$. It completes the proof that f is a quasi-equivalence. ■

Theorem 4.17 is proven. ■

5. A Quillen equivalence between $\mathbb{C}_{dgwu}(\mathbb{k})$ and $\mathbb{C}_{dg}(\mathbb{k})$

Let \mathcal{C} and \mathcal{D} be two model categories. Recall that a *Quillen pair* of functors $L : \mathcal{C} \rightleftarrows \mathcal{D} : R$ is an adjoint pair of functors such that L preserves cofibrations and acyclic cofibrations, or equivalently, R preserves fibrations and acyclic fibrations, [Ho, 1.3], [Hi, 8.5]. These two conditions are sufficient to show that the Quillen pair of functors descends to a pair of adjoint functors

$$L : Ho(\mathcal{C}) \rightleftarrows Ho(\mathcal{D}) : R \tag{33}$$

between the homotopy categories.

When \mathcal{C} is cofibrantly generated, there is a manageable criterium for an adjoint pair of functors to be a Quillen pair [[Ho], Lemma 2.1.20]:

5.1. PROPOSITION. *Let \mathcal{C}, \mathcal{D} be closed model categories, with \mathcal{C} cofibrantly generated with generating cofibrations I and generating acyclic cofibrations J . Let $L : \mathcal{C} \rightleftarrows \mathcal{D} : R$ be an adjoint pair of functors. Assume that $L(f)$ is a cofibration for all $f \in I$ and $L(f)$ is a trivial cofibration for all $f \in J$. Then (L, R) is a Quillen pair.*

Let $C \in \mathbb{C}_{dgwu}(\mathbb{k})$. Define

$$L_1(C) := C/I,$$

where I is the dg-category ideal generated by all $p_n(\cdot), n \geq 2$. Clearly $L_1(C) \in \mathbb{C}_{dg}(\mathbb{k})$.

This assignment $C \mapsto L_1(C)$ gives rise to a functor $L_1 : \mathbb{C}_{dgwu}(\mathbb{k}) \rightarrow \mathbb{C}_{dg}(\mathbb{k})$.

Let $\mathcal{D} \in \mathbb{C}_{dg}(\mathbb{k})$. Define $R_1 : \mathbb{C}_{dg}(\mathbb{k}) \rightarrow \mathbb{C}_{dgwu}(\mathbb{k})$ as the fully-faithful embedding from Example 2.3.

5.2. PROPOSITION. *The following statements are true:*

(1) *there is an adjunction*

$$\mathrm{Hom}_{\mathbb{C}_{dg}(\mathbb{k})}(L_1(\mathcal{C}), \mathcal{D}) \cong \mathrm{Hom}_{\mathbb{C}_{dgwu}(\mathbb{k})}(\mathcal{C}, R_1(\mathcal{D}))$$

(2) *the functors*

$$L_1 : \mathbb{C}_{dgwu}(\mathbb{k}) \rightleftarrows \mathbb{C}_{dg}(\mathbb{k}) : R_1$$

form a Quillen pair of functors.

PROOF. (1) : any morphism $F : C \rightarrow R_1(D)$ sends $p_n(\cdot), n \geq 2$ to 0, since $D \in \mathbb{C}_{dg}(\mathbb{k})$, and therefore this morphism is the same as a morphism $L_1(C) \mapsto D$.

(2) : the dg categories $\{L_1(\beta(n)), L_1(Q)\}$ form exactly the set I of generating cofibrations for the Tabuada closed model structure on $\mathbb{C}_{dg}(\mathbb{k})$, and the dg categories $\{L_1(\alpha(n)), L_1(\kappa)\}$ form the set of generating acyclic cofibrations for this model structure. The statement follows then by Proposition 5.1. ■

Recall that a Quillen pair $L : \mathcal{C} \rightleftarrows \mathcal{D} : R$ is called a *Quillen equivalence* if it satisfies the following condition:

For a cofibrant $X \in \mathcal{C}$ and a fibrant $Y \in \mathcal{D}$, a morphism $f : LX \rightarrow Y$ is a weak equivalence in \mathcal{D} if and only if its adjoint morphism $X \rightarrow RY$ also is, [Ho, 1.3.3], [Hi, 8.5]. This condition implies the corresponding adjoint pair of functors between the homotopy categories $\mathbb{33}$ is an adjoint *equivalence* of categories.

5.3. THEOREM. *The Quillen pair of functors*

$$L_1 : \mathbb{C}_{dgwu}(\mathbb{k}) \rightleftarrows \mathbb{C}_{dg}(\mathbb{k}) : R_1$$

is a Quillen equivalence.

PROOF. Let $X \in \mathbb{C}_{dgwu}(\mathbb{k})$ be cofibrant and $Y \in \mathbb{C}_{dg}(\mathbb{k})$ fibrant. We have to prove that $f : L_1X \rightarrow Y$ is a weak equivalence in $\mathbb{C}_{dg}(\mathbb{k})$ if and only if the adjoint morphism $f^* : X \rightarrow R_1Y$ is a weak equivalence in $\mathbb{C}_{dgwu}(\mathbb{k})$,

It is enough to prove the statement for the case when X is an I -cell. Indeed, by the small object argument, for any X there exist an I -cell X' such that $p : X' \rightarrow X$ is an acyclic fibration. The Quillen left adjoint L maps the weak equivalences between cofibrant object to weak equivalences, by [Hi, Prop. 8.5.7]. Therefore, $L(p) : L(X') \rightarrow L(X)$ is a weak equivalence. There is a map $i : X \rightarrow X'$ such that $p \circ i = \mathrm{id}$, given by the RLP. By 2-of-3 axiom, i is a weak equivalence, and $L(i)$ also is.

We assume that X is an I -cell. Denote by V the graded graph of generators of X . We need to prove that for any objects $x, x' \in X$, the cone $M := \mathrm{Cone}(f : L_1X(x, x') \rightarrow Y(fx, fx'))$ is acyclic if and only if the cone $N := \mathrm{Cone}(f^* : X(x, x') \rightarrow R_1Y(f^*x, f^*x'))$. Denote by $\tilde{\mathcal{O}} := \mathrm{Ker}(P : \mathcal{O}' \rightarrow \mathcal{A}ssoc_+)$, where P is the dg operad map sending all $p_n(\dots), n \geq 2$ to 0.

There is a canonical map $\omega : N \rightarrow M$, and $\mathrm{Cone}(\omega)$ is quasi-isomorphic to $F_{\tilde{\mathcal{O}}}(V)(x, x')$,

where $F_{\tilde{\mathcal{O}}}(V)$ is the free algebra over $\tilde{\mathcal{O}}$ generated by V , with an extra differential coming from the differential in the I -cell X . Since the dg operad \mathcal{O}' is quasi isomorphic to $\mathcal{A}ssoc_+$, $\tilde{\mathcal{O}}$ is acyclic. Therefore $F_{\tilde{\mathcal{O}}}(V)$ is acyclic by the Künneth formula, and so M is quasi isomorphic to N , by the acyclicity of $Cone(\omega)$. We conclude that M is acyclic if and only if N is. ■

6. A proof of Proposition 3.3

Here we prove Proposition 3.3.

Recall the maps $P_n, n \geq 1$ defined in (13).

Recall that by Lemma 3.2 one has

$$deg(P_n(\phi_1, \dots, \phi_n))_{ij} = |\phi_1| + \dots + |\phi_n| - n + 1 = deg(p_l(\kappa_1, \dots, \kappa_l)), \forall \kappa \in \mathcal{P}aths_{ij}$$

Proposition 3.3 reads:

6.1. PROPOSITION. *The maps $\{P_n\}, n \geq 1$ are Taylor components of an A_∞ functor*

$$P: C^{\text{pretr}} \oplus \mathbb{k}_{C^{\text{pretr}}} \rightarrow C^{\text{pretr}}$$

PROOF. Proving the statement amounts to proving the following identities, for any $n \geq 1$ (see (38) for the sign convention):

$$\begin{aligned} dP_n(\phi_n, \dots, \phi_1) + \sum_{a+b=n} (-1)^{(a-1)(\sum_{i=1}^b |\phi_i|)+b-1} P_a(\phi_n, \dots, \phi_{b+1}) P_b(\phi_b, \dots, \phi_1) = \\ \sum_{k=1}^{n-1} (-1)^{k-1} P_{n-1}(\phi_n, \dots, \phi_{k+1} \circ \phi_k, \dots, \phi_1) + \sum_{k=1}^n (-1)^{n-1+\sum_{i=1}^{k-1} |\phi_i|} P_n(\phi_n, \dots, d\phi_k, \dots, \phi_1), \end{aligned} \tag{34}$$

where

$$dP_n(\phi_n, \dots, \phi_1) = d_{naive}P_n(\phi_n, \dots, \phi_1) + q' \circ P_n(\phi_n, \dots, \phi_1) - (-1)^{n-1} P_n(\phi_n, \dots, \phi_1) \circ q \tag{35}$$

Writing down explicitly the first line of equation 34 (and dropping the signs to \pm for simplicity), we get

$$\begin{aligned} dP_n(\phi_n, \dots, \phi_1) &= d_{naive}P_n(\dots) + q' \circ P_n(\dots) + (-1)^{n-1} P_n(\dots) \circ q = \\ &= d_{naive} \left(\sum_{\kappa \in \mathcal{P}aths} \pm p_m(\kappa_m, \dots, \kappa_1) \right) + q' \circ \sum_{\kappa \in \mathcal{P}aths} \pm p_{m'}(\kappa_{m'}, \dots, \kappa_1) \\ &+ (-1)^{n-1} \sum_{\kappa \in \mathcal{P}aths} p_{m''}(\kappa_{m''}, \dots, \kappa_1) \circ q = \\ &\sum_{\kappa \in \mathcal{P}aths} \left(\sum_{a+b=m} \pm p_a(\dots) \circ p_b(\dots) + \sum_{i=1}^{n-1} \pm p_{m-1}(\dots, m(\kappa_{i+1}, \kappa_i), \dots) + \sum_{i=1}^m \pm p_n(\dots, d\kappa_i, \dots) \right) + \\ &q' \circ \sum_{\kappa \in \mathcal{P}aths} \pm p_{m'}(\kappa_{m'}, \dots, \kappa_1) + (-1)^{n-1} \sum_{\kappa \in \mathcal{P}aths} \pm p_{m''}(\kappa_{m''}, \dots, \kappa_1) \circ q \end{aligned} \tag{36}$$

We stress that inside the terms $\sum_{\kappa \in \mathcal{P}aths} (\sum_{i=1}^{n-1} (-1)^s p_{m-1}(\dots, m(\kappa_{i+1}, \kappa_i), \dots))$ the composition term might be of following three types:

- $q_{jl} \circ q_{kj}$
- $q \circ \phi$ or $\phi \circ q$
- $\phi_i \circ \phi_{i+1}$.

Similarly inside the terms $\sum_{\kappa \in \mathcal{P}aths} (\sum_{i=1}^m (-1)^t p_n(\dots, d\kappa_i, \dots))$ the term which is differentiated might be of the following two types:

- $d\phi$
- dq_{kl} .

We also write down the possible terms of $\sum_{\kappa \in \mathcal{P}aths} (\sum_{a+b=m} \pm p_a(\dots) p_b(\dots))$:

- $p_a(\dots) p_b(\dots)$
- $q \circ p_m$ or $p_l(\dots) \circ q$.

Thence we can write for the r.h.s. of (36)

$$\begin{aligned}
 (\text{r.h.s. of (36)}) = & \sum_{\kappa \in \mathcal{P}aths} \pm \left(\underbrace{\sum_i \pm p_n(\dots, d\phi, \dots)}_i \pm \sum_i \pm p_n(\dots, dq_{kl}, \dots) \pm \sum_i \pm p_n(\dots, q_{jl} \circ q_{kj}, \dots) \right) + \\
 & \sum_{\kappa \in \mathcal{P}aths} \pm \left(\pm \sum_i \pm p_n(\dots, q \circ \phi, \dots) \pm \sum_i \pm p_n(\dots, \phi \circ q, \dots) \pm \sum_i \pm p_n(\dots, \phi_{i+1} \circ \phi_i, \dots) \right) + \\
 & \sum_{\kappa \in \mathcal{P}aths} \pm \left(\underbrace{\pm q' \circ p_m(\dots)}_a \pm \underbrace{\pm p_l(\dots) \circ q}_a \pm \sum_a \pm p_a(\dots) \circ p_b(\dots) \right) + \tag{37} \\
 & \underbrace{q' \circ \sum_{\kappa \in \mathcal{P}aths} \pm p_{m'}(\kappa_1, \dots, \kappa_{m'})}_{\kappa \in \mathcal{P}aths} + \sum_{\kappa \in \mathcal{P}aths} \pm p_{m''}(\kappa_1, \dots, \kappa_{m''}) \circ q \stackrel{?}{=} \\
 & \sum_{a+b=n} \pm P_a(\dots) \circ P_b(\dots) + \sum_i \pm P_{n-1}(\dots, m(\phi_{i+1}, \phi_i), \dots) + \sum_i \pm P_n(\dots, d\phi, \dots)
 \end{aligned}$$

We have to prove the equation marked by question sign. The dashed underlined terms gets cancelled by the Maurer-Cartan condition on the q_{ij} : indeed if there exists $\kappa \in \mathcal{P}aths_{ij}$ which contains q_{kl} , there will also exist a path $\kappa' \in \mathcal{P}aths_{ij}$ containing all the terms $q_{jl} \circ q_{kj}$, since they have got same domain and codomain as q_{kl} and in $\mathcal{P}aths_{ij}$ we were

considering all the possible paths.

Therefore we are left with the following expressions:

$$\begin{aligned} & \pm \sum_i \sum_{\kappa \in \mathcal{P}aths} p_n(\dots, m(\phi_{i+1}, \phi_i), \dots) \pm \sum_{a+b=n} \sum_{\kappa \in \mathcal{P}aths} \pm p_a(\dots) \circ p_b(\dots) + \\ & \sum_i \sum_{\kappa \in \mathcal{P}aths} \pm (p_n(\dots, d\phi_i, \dots) + p_n(\dots, q' \circ \phi_i, \dots) + p_n(\dots, \phi_i \circ q, \dots)) = \\ & \sum_{a+b=n} \pm P_a(\dots) \circ P_b(\dots) + \sum_i \pm P_{n-1}(\dots, m(\phi_{i+1}, \phi_i), \dots) + \sum_i \pm P_n(\dots, d\phi, \dots), \end{aligned}$$

which shows the desired equation, up to signs. The correctness of signs (which were explicitly defined in (13) and (34)) is checked by a long but routine computation. ■

A.

In the paper, we use only A_∞ functors between *dg categories*.

We need to specify the signs in the A_∞ identity. The common sign conventions can be found in [Kel, 3.4] for the *left to right* order formulas, and in [Ly1, 2.4] for the *right to left* order formulas. As we adapt here the right to left formalism, our signs agree with the ones in [Ly1].

A.1. DEFINITION. *Let C, D be (non-unital) dg categories over \mathbb{k} . An A_∞ functor $F: C \rightarrow D$ is given by:*

- a map of sets $F_0: \text{Ob}(C) \rightarrow \text{Ob}(D)$,
- for any $n \geq 1$ and a sequence of objects $x_0, \dots, x_n \in \text{Ob}(C)$ of degree $1 - n$

$$F_n: C(x_{n-1}, x_n) \otimes \cdots \otimes C(x_1, x_2) \otimes C(x_0, x_1) \rightarrow D(F_0(x_0), F_0(x_n))[1 - n]$$

such that one has:

$$\begin{aligned} & d(F_n(f_n \otimes \cdots \otimes f_1)) + \\ & \sum_{a+b=n} (-1)^{b-1+(a-1)(|f_1|+\cdots+|f_b|)} F_a(f_n \otimes \cdots \otimes f_{b+1}) \cdot F_b(f_b \otimes \cdots \otimes f_1) = \\ & \sum_{k=0}^{n-1} (-1)^{n-1+|f_1|+\cdots+|f_k|} F_n(f_n \otimes \cdots \otimes f_{k+2} \otimes d(f_{k+1}) \otimes f_k \otimes \cdots \otimes f_1) + \\ & \sum_{k=0}^{n-2} (-1)^k F_{n-1}(f_n \otimes \cdots \otimes f_{k+3} \otimes (f_{k+2} \circ f_{k+1}) \otimes f_k \otimes \cdots \otimes f_1) \end{aligned} \quad (38)$$

A.2. DEFINITION. Let C, D be (unital) dg categories over \mathbb{k} , $F: C \rightarrow D$ an A_∞ functor, $\{F_i\}_{i \geq 1}$ its Taylor components.

(1) F is called strongly unital if $F_1(\text{id}_X) = \text{id}_{F(X)}$ for any object $X \in C$, and

$$F_n(\dots, \text{id}_X, \dots) = 0$$

for any object X and $n \geq 2$,

(2) F is called weakly unital if $F_1(\text{id}_X) = \text{id}_{F(X)}$ for any object $X \in C$ (and the second condition is dropped).

B. A cofibrant resolution of a unital dg algebra

Here we provide a canonical unital cofibrant dg algebra, quasi-isomorphic to a unital dg algebra A over a field \mathbb{k} . (We consider only the case of a dg algebra for simplicity, the construction is directly generalised for the case of a small dg category).

The classical bar-cobar resolution of A fails to be unital, it is only weakly unital dg algebra. A well-known explicit unital construction comes from the *curved* version of bar-cobar duality, due to L.Positselski [Pos] (see also [Ly3]). A drawback of this construction is that it is not canonical.

We haven't seen this construction in the literature. In our opinion, it deserves to be included as an appendix to this paper, by two reasons. At first, it "replaces" the bar-cobar resolution, which fails to be strictly unital, which was one of our starting points. At second, it can be easily generalised to a cofibrant resolution of $i(C)$ in $\mathbb{C}_{dgwu}(\mathbb{k})$, for $C \in \mathbb{C}_{dg}(\mathbb{k})$.

Let A be a unital dg algebra over \mathbb{k} . Consider the dg algebra $A_+ = A \oplus \mathbb{k}[1]$. It is a unital dg algebra with unit 1_A , and the product of A with $\mathbb{k}[1]$, as well as of $\mathbb{k}[1]$ with itself, is defined as 0.

Consider the bar-complex

$$\text{Bar}(A_+) = \bigoplus_{n \geq 1} A_+[1]^{\otimes n}$$

which is a dg coalgebra. We use the notation ξ for a generator of $\mathbb{k}[1]$. Then a general monomial element of $\text{Bar}(A_+)$ is denoted as

$$a_1 \otimes \dots \otimes a_{i_1} \otimes \xi \otimes a_{i_1+1} \otimes \dots \otimes a_{i_2} \otimes \xi \otimes a_{i_2+1} \otimes \dots$$

Now consider the *unital* dg algebra

$$C_0(A) = \text{Cobar}_+(\text{Bar}(A_+))$$

where, for a dg coalgebra B ,

$$\text{Cobar}_+(B) = \mathbb{k} \oplus \bigoplus_{n \geq 1} B[-1]^{\otimes n}$$

with the cobar-differential. It is a unital dg algebra. Denote by $1_{\mathbb{k}}$ the unit of \mathbb{k} . It is the unit of $C_0(A)$. We denote the product in $\text{Cobar}_+(B)$ by \boxtimes .

Consider a derivation d_ξ of degree $+1$ of $C_0(A)$ whose only non-zero Taylor coefficient is linear, and is defined as

$$\begin{aligned} d_\xi|_A &= 0, \quad d_\xi(\xi) = 1_{\mathbb{k}} - 1_A \\ d_\xi(x_1 \otimes \cdots \otimes x_n) &= \sum_{\ell=1}^k \pm x_1 \otimes \cdots \otimes x_{i_\ell-1} \otimes 1_A \otimes x_{i_\ell+1} \otimes \cdots \otimes x_n \end{aligned} \quad (39)$$

where $x_{i_1} = \cdots = x_{i_k} = \xi$ and other $x_i \in A$

One has

B.1. LEMMA. *The differential d_ξ squares to 0, and d_ξ commutes with $d_{\text{Bar}} + d_{\text{Cobar}}$. Consequently, $d_{\text{tot}} := d_{\text{Bar}} + d_{\text{Cobar}} + d_\xi$ squares to 0.*

PROOF. It is a direct computation. ■

We denote

$$C(A) = (C_0(A), d_{\text{Bar}} + d_{\text{Cobar}} + d_\xi)$$

It is a unital dg algebra.

B.2. PROPOSITION. *There is a unital dg algebra map $p: C(A) \rightarrow A$ which is a quasi-isomorphism.*

PROOF. We start with computing the cohomology of $(C(A), d_{\text{tot}})$. The differential $d_{\text{Bar}} + d_{\text{Cobar}}$ preserves the total number of ξ -factors in a (homogeneous in ξ) element of $C(A)$. It makes $C(A)$ a bicomplex. Define

$$\text{deg}_{\text{Bar}}(x_1 \otimes \cdots \otimes x_n) = -n + \sum_i \text{deg}_0 x_i$$

where $\text{deg}_0(a) = \text{deg}_A(a)$ and $\text{deg}_0(\xi) = 0$. Next, define

$$\text{deg}_1(\omega_1 \boxtimes \cdots \boxtimes \omega_k) = k + \sum_i \text{deg}_{\text{Bar}}(\omega_i)$$

$$\text{deg}_{\text{Cobar}}(\omega_1 \boxtimes \cdots \boxtimes \omega_k) = k$$

$$\text{deg}_{\text{Bar}}(\omega_1 \boxtimes \cdots \boxtimes \omega_k) = \sum_i \text{deg}_{\text{Bar}}(\omega_i)$$

where $\omega_i \in \text{Bar}(A_+)$. Finally, define

$$\text{deg}_\xi(\alpha) = -(\#\xi \text{ in } \alpha), \quad \alpha \in C(A)$$

$$\text{deg}_{\text{tot}}(\alpha) = \text{deg}_1(\alpha) + \text{deg}_\xi(\alpha)$$

where \deg_{tot} is the degree of α in $C(A)$.

Thus $C(A)$ becomes a bicomplex, with $C(A)^{a,b}$ defined as the spaces of elements $\alpha \in C(A)$ with $\deg_1 \alpha = a$, $\deg_\xi(\alpha) = b$.

We compute the cohomology of $C(A)$ by using a spectral sequence, which computes the cohomology of $d_{\text{Bar}} + d_{\text{Cobar}}$ at first. The bicomplex lives in the *II* and *III* quarters, so the spectral sequence converges.

The term $E_1^{a,b}$ of the spectral sequence is equal to $H^a(C(A)^{\bullet,b}, d_{\text{Bar}} + d_{\text{Cobar}})$. Thus, we have to compute the cohomology of the complex $(C(A)^{\bullet,b}, d_{\text{Bar}} + d_{\text{Cobar}})$. Denote this complex by C_b^\bullet .

The complex C_b^\bullet (for b fixed) is by its own a bicomplex, with differentials d_{Bar} and d_{Cobar} . Thus $C_b^{m,n}$ consist of all elements $\alpha \in C_b^\bullet$ with $\deg_{\text{Bar}}(\alpha) = m$ and $\deg_{\text{Cobar}}(\alpha) = n$ (in this case, $\deg_{\text{tot}}(\alpha) = m + n + b$).

The spectral sequence, whose first differential is d_{Cobar} , converges (the other possible spectral sequence, whose first differential is d_{Bar} , generally diverges).

We denote by $\mathbb{E}(b)_k^{m,n}$ the k -th term of this spectral sequence. We have

$$\mathbb{E}(b)_1^{m,n} = H^m(\mathbb{E}(b)_1^{\bullet,n}, d_{\text{Cobar}})$$

B.3. LEMMA. *One has:*

$$\mathbb{E}(b)_1^{m,n} = \begin{cases} 0, & m \neq 0, 1 \text{ or } m = 0, n \neq 0 \\ \mathbb{k}, & b = 0, m = 0, n = 0 \\ \mathbb{k}[1], & b = -1, m = 1, n = -1 \\ A^n, & b = 0, m = 1 \\ 0, & b \neq 0, -1 \end{cases} \quad (40)$$

where $(-)^n$ stands for degree n elements.

PROOF. The argument is standard and comes from the following observation. Let V be a (graded) vector space, consider the cofree non-unital coalgebra $T_{\geq 1}^\vee(V) = \bigoplus_{n \geq 1} V^{\otimes n}$. Then $\text{Cobar}(T_{\geq 1}^\vee(V))$ is quasi-isomorphic to $V[-1]$. This statement is proven using Koszul duality. ■

It follows that the spectral sequence \mathbb{E}_\bullet collapses at the \mathbb{E}_1 term.

Now turn back to the spectral sequence E_\bullet .

B.4. LEMMA. *One has*

$$E_1^{a,b} = \begin{cases} \mathbb{k}[1], & a = 0, b = -1 \\ (\mathbb{k} \oplus A)^a, & b = 0 \\ 0, & \text{otherwise} \end{cases} \quad (41)$$

The differential d_1 is induced by d_ξ . It looks like

$$\mathbb{k}[1] \xrightarrow{d_1} \mathbb{k} \oplus A, \quad d_1: 1 \mapsto 1_{\mathbb{k}} - 1_A$$

Its cohomology is isomorphic to A . The spectral sequence E . collapses at the E_2 term.

It completes the computation of cohomology of $C(A)$.

Now define a map of dg algebras $p: C(A) \rightarrow A$ on generators

$$\begin{aligned} p|_{(\mathbb{k}[1] \oplus A)^{\otimes n}} &= 0, \quad n \geq 2 \\ p(\xi) &= 0 \\ p|_A &= \text{id} \\ p(1_{\mathbb{k}}) &= 1_A \end{aligned} \tag{42}$$

and extend it to $C(A)$ as a map of algebras.

It follows from the previous computation that p is a quasi-isomorphism. ■

C. Push-outs in the category $\mathbb{C}_{dgwu}(\mathbb{k})$

Consider the following push-out diagram:

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{i_1} & \mathcal{C}_1 \\ i_2 \downarrow & & \\ \mathcal{C}_2 & & \end{array} \tag{43}$$

where $i_a : \mathcal{C} \hookrightarrow \mathcal{C}_a, a = 1, 2$ are embeddings of wu dg categories. We provide an explicit formula for the push-out \mathcal{E} . This formula is essentially used in the proof of Theorem 4.17. Note that this colimit \mathcal{E} is equivalently the colimit of the following coequalizer diagram:

$$\mathcal{C} \rightrightarrows \mathcal{C}_1 \oplus \mathcal{C}_2 \tag{44}$$

where the maps are $(i_1, 0)$ and $(0, i_2)$. The category structure on $\mathcal{C}_1 \oplus \mathcal{C}_2$ is defined as

$$\begin{aligned} \text{Ob}(\mathcal{C}_1 \oplus \mathcal{C}_2) &:= \text{Ob}(\mathcal{C}_1) \sqcup \text{Ob}(\mathcal{C}_2) \\ \text{Hom}_{\mathcal{C}_1 \oplus \mathcal{C}_2}(x, y) &= \begin{cases} \text{Hom}_{\mathcal{C}_a}(x, y) & \text{if } x, y \in \text{Ob}(\mathcal{C}_a) \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

In [PS, Prop.1.18] we considered the general coequalizers in the category $\mathbb{C}_{dgwu}(\mathbb{k})$. Here we provide the corresponding description for the case of the coequalizer (44). Here we essentially use that i_1, i_2 are fully-faithful functors. The derivation of this description from loc.cit. is straightforward.

We use notation \bar{a} which is defined as $\bar{1} = 2, \bar{2} = 1$.

C.1. PROPOSITION. Assume i_1, i_2 in (43) are fully faithful. Then the push-out weakly unital dg category \mathcal{E} has the following description.

The objects of \mathcal{E} are given by the coequalizer of sets:

$$Ob(\mathcal{E}) = Ob(\mathcal{C}_1) \sqcup Ob(\mathcal{C}_2) / \sim$$

where \sim is the equivalence relation generated by: $i_1(x) \sim i_2(x), \forall x \in Ob(\mathcal{C})$.

Consider $x, y \in Ob(\mathcal{C}_a), a = 1, 2$. Then:

$$\begin{aligned} \mathcal{E}(x, y) = & \mathcal{C}_a(x, y) \bigoplus_{u, v \in \mathcal{C}} \bigoplus \mathcal{O}'(3) \otimes \mathcal{C}_a(v, y) \otimes \bar{\mathcal{C}}_a(u, v) \otimes \mathcal{C}_a(x, u) \bigoplus \\ & \bigoplus_{u, v, u_1, v_1 \in \mathcal{C}} \mathcal{O}'(5) \otimes \mathcal{C}_a(v_1, y) \otimes \bar{\mathcal{C}}_a(u_1, v_1) \otimes \mathcal{C}_a(u, v_1) \otimes \bar{\mathcal{C}}_a(v, u) \otimes \mathcal{C}_a(x, v) \bigoplus \dots \end{aligned} \tag{45}$$

where we identify an object $u \in \mathcal{C}$ with its images $i_a(u) \in \mathcal{C}_a$, and

$$\bar{\mathcal{C}}_a(u, v) := \mathcal{C}_a(u, v) / i_a(\mathcal{C}(x, y)).$$

For $x \in Ob(\mathcal{C}_1) \setminus Ob(\mathcal{C}), y \in Ob(\mathcal{C}_2) \setminus Ob(\mathcal{C})$, one has:

$$\begin{aligned} \mathcal{E}(x, y) := & \bigoplus_{u \in \mathcal{C}} \mathcal{O}'(2) \otimes \mathcal{C}_2(u, y) \otimes \mathcal{C}_1(x, u) \\ & \bigoplus_{u, v, w \in \mathcal{C}} \bigoplus \mathcal{O}'(4) \otimes \mathcal{C}_2(w, y) \otimes \bar{\mathcal{C}}_1(v, w) \otimes \bar{\mathcal{C}}_2(u, v) \otimes \mathcal{C}_1(x, u) \bigoplus \dots \end{aligned} \tag{46}$$

References

- [Ba] M.A.Batanin, Monoidal globular categories as a natural environment for the theory of weak n-categories, *Adv. Math.* **136**(1) (1998), 39-103
- [BK] A. I. Bondal and M. M. Kapranov, *Enhanced triangulated categories*. Math. USSR Sbornik, vol. 70 (1991), no. 1, 93107. (Russian original: vol. 181 (1990), no. 5).
- [COS] A.Canonaco, M.Ornaghi, P.Stellari, Localizations of the category of A_∞ categories and internal Homs, preprint arXiv 1811.07830
- [Dr] V.Drinfeld, DG quotients of DG categories, *J. Algebra* **272**(2), (2004), 643-691
- [DS] W. G. Dwyer, J. Spalinski, Homotopy theories and model categories, in: *Handbook on Algebraic Topology*, Elsevier, 1995
- [Fa] G. Faonte, *Simplicial Nerve of an A_∞ -category*, Theory and Applications of Categories, Vol. 32, 2017, No. 2, pp 31-52.
- [GJ] P.G.Goerss, J.F.Jardine, *Simplicial Homotopy Theory*, Birkhäuser Progress in Mathematics, Vol. 174, 1999

- [GS] P.Goerss and K.Schemmerhorn, Model Categories and Simplicial Methods, Notes from lectures given at the University of Chicago, August 2004, available at P.Goerss' webpage
- [Hi] Ph.S.Hirschhorn, *Model Categories and Their Localizations*, AMS Mathematical Surveys and Monographs, Vol. 99, 2003
- [Ho] M.Hovey, *Model categories*, AMS Math. Surveys and Monographs, vol. 62, 1999
- [K1] M.Kontsevich, Lectures at ENS, Paris, Spring 1998: notes taken by J.Bellaiche, J.-F.Dat, I.Marin, G.Racinet and H.Randriambololona, unpublished
- [K2] M.Kontsevich, Homological algebra of Mirror Symmetry, *Proceedings of the International Congress of Mathematicians*, Zürich 1994, vol. I, Birkhauser 1995, 120-139
- [KS] M.Kontsevich, Y.Soibelman, Notes on A_∞ algebras, A_∞ categories, and non-commutative geometry, in: "Homological Mirror Symmetry: New Developments and Perspectives" (A.Kapustin et al. (Eds.)), Lect. Notes in Physics 757 (Springer, Berlin Heidelberg 2009) 153-219
- [Kel] B.Keller, Introduction to A_∞ algebras and modules, *Homology, Homotopy, and Applications*, **3**(2001), 1-35
- [Ly1] V.Lyubashenko, Category of A_∞ categories, *Homology Homotopy Appl.*, Vol. 5(1) (2003), 1-48
- [Ly2] V.Lyubashenko, Homotopy unital A_∞ algebras, *J. Algebra* 329 (2011) no. 1, 190-212
- [Ly3] V.Lyubashenko, Bar and cobar constructions for curved algebras and coalgebras. *Mat. Stud.*, **40**(2) (2013), 115131
- [PS] P. Panero, B. Shoikhet, *A Quillen Model Structure on the Category of Kontsevich-Soibelman Weakly Unital DG Categories*, preprint arXiv 1907.07970
- [Pos] L. Positselski, Weakly curved A_∞ -algebras over a topological local ring. *Mm. Soc. Math. Fr. (N.S.)* No. 159 (2018), vi+206 pp.
- [Sh] B. Shoikhet, On the twisted tensor product of small dg categories, to appear in *Journal of Non-commutative Geometry*, preprint arxiv:1803.01191
- [Tab1] G.Tabuada, A Quillen model structure on the category of dg categories C. R. Math. Acad. Sci. Paris, 340 (2005), 15-19
- [Tab2] G.Tabuada, On Drinfeld's dg quotient, *Journal of Algebra*, **323** (2010), 1226-1240
- [To] B. Toën, The homotopy theory of dg-categories and derived Morita theory, *Invent. Math.* **167**(3), (2007), 615-667

P.P.:

Universiteit Antwerpen, Campus Middelheim, Wiskunde en Informatica, Gebouw G
Middelheimlaan 1, 2020 Antwerpen, België

e-mail: Piergiorgio.Panero@uantwerpen.be

B.Sh.:

Universiteit Antwerpen, Campus Middelheim, Wiskunde en Informatica, Gebouw G
Middelheimlaan 1, 2020 Antwerpen, België

Laboratory of Algebraic Geometry, National Research University Higher School of Economics,
Moscow, Russia

e-mail: Boris.Shoikhet@uantwerpen.be

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

INFORMATION FOR AUTHORS \LaTeX 2 ϵ is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

MANAGING EDITOR. Geoff Cruttwell, Mount Allison University: gcruttwell@mta.ca

TEXNICAL EDITOR. Michael Barr, McGill University: michael.barr@mcgill.ca

ASSISTANT TEX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Julie Bergner, University of Virginia: jeb2md@virginia.edu

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Gabriella Böhm, Wigner Research Centre for Physics: bohm.gabriella@wigner.mta.hu

Valeria de Paiva, Nuance Communications Inc: valeria.depaiva@gmail.com

Richard Garner, Macquarie University: richard.garner@mq.edu.au

Ezra Getzler, Northwestern University: getzler@northwestern.edu

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Dirk Hofmann, Universidade de Aveiro: dirk@ua.pt

Pieter Hofstra, Université d' Ottawa: phofstra@uottawa.ca

Anders Kock, University of Aarhus: kock@math.au.dk

Joachim Kock, Universitat Autònoma de Barcelona: kock@mat.uab.cat

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: matias.menni@gmail.com

Ieke Moerdijk, Utrecht University: i.moerdijk@uu.nl

Susan Niefield, Union College: niefiels@union.edu

Kate Ponto, University of Kentucky: kate.ponto@uky.edu

Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: ross.street@mq.edu.au

Tim Van der Linden, Université catholique de Louvain: tim.vanderlinden@uclouvain.be