

ENRICHED LOCALLY GENERATED CATEGORIES

I. DI LIBERTI AND J. ROSICKÝ

ABSTRACT. We introduce the notion of \mathcal{M} -locally generated category for a factorization system $(\mathcal{E}, \mathcal{M})$ and study its properties. We offer a Gabriel-Ulmer duality for these categories, introducing the notion of nest. We develop this theory also from an enriched point of view. We apply this technology to Banach spaces showing that it is equivalent to the category of models of the nest of finite-dimensional Banach spaces.

Contents

1	Introduction	661
2	Locally generated categories	663
3	Extended Gabriel-Ulmer duality	669
4	Enriched locally generated categories	673
5	Enriched Gabriel-Ulmer duality	677
6	Banach spaces	678

1. Introduction

Locally presentable categories were introduced by Gabriel and Ulmer [12] in 1971 and, since then, their importance was steadily growing. Today, they form an established framework to do category theory in the daily practice of the working mathematician. The reason of their success is merely evidence-based. On the one hand, they are technically handy, allowing transfinite constructions (e.g. the small object argument) and thus offering pleasing versions of relevant tools (e.g. the adjoint functor theorem). On the other hand, a vast majority of relevant categories happen to be locally presentable (with some very disappointing exceptions, like the category of topological spaces).

Gabriel and Ulmer [12] also introduced locally generated categories. While locally presentable categories are based on the classical algebraic concept of a (finitely) presentable object, locally generated categories are based on (finitely) generated objects. This makes them easier to handle than locally presentable ones. Every locally finitely

The first author was supported by the Grant Agency of the Czech Republic project EXPRO 20-31529X and RVO: 67985840. The second author was supported by the Grant Agency of the Czech Republic under the grant 19-00902S.

Received by the editors 2022-03-20 and, in final form, 2022-03-29.

Transmitted by Giuseppe Rosolini. Published on 2022-03-31.

2020 Mathematics Subject Classification: 18C30, 18C35, 18D20, 18C15, 46A03, 46B04, 54B30.

Key words and phrases: locally generated category, nest, Gabriel-Ulmer duality, enriched category, metric spaces, Banach spaces, accessible monad.

© I. Di Liberti and J. Rosický, 2022. Permission to copy for private use granted.

generated category is locally λ -presentable for some regular cardinal. Yet, a category can be locally finitely generated without being locally finitely presentable. Despite this, locally generated categories were somewhat neglected. They were mentioned in [1] and, later, generalized in [2]. In a nutshell, a \mathcal{M} -locally generated category is a cocomplete category \mathcal{K} , equipped with a factorization system $(\mathcal{E}, \mathcal{M})$, that is generated by a set of \mathcal{M} -generated objects. In [12] they were only dealing with (Strong Epi, Mono), in [2], monomorphisms were replaced by any right part of a proper factorization system $(\mathcal{E}, \mathcal{M})$.

We slightly generalize [2] by taking any factorization system $(\mathcal{E}, \mathcal{M})$. A locally λ -presentable category is then a $\mathcal{K}^{\rightarrow}$ -locally λ -generated category. More importantly, we extend the (nearly forgotten) Gabriel-Ulmer duality for locally generated categories to our setting. While the famous Gabriel-Ulmer duality for locally finitely presentable categories is based on the fact that they are sketched by a finite limit sketch, their duality for locally finitely generated categories uses finite limits and multiple pullbacks of monomorphisms. This explains why locally finitely generated categories do not need to be locally finitely presentable. In our situation, we use multiple pullbacks of morphisms from \mathcal{M} .

Our main goal is to apply these ideas to Banach spaces. For this, we have to make our theory enriched. While enriched locally presentable categories were treated by Kelly [17], enriched locally generated categories have not been considered. Our motivation was [3, Remark 7.8] where finite-dimensional Banach spaces, which are not finitely presentable, were shown to be finitely generated with respect to isometries when Banach spaces are enriched over **CMet**, the category of complete metric spaces. The relevant factorization system is (dense maps, isometries). Our main accomplishment is that Banach spaces can be sketched using finite-dimensional Banach spaces equipped with finite weighted limits and multiple pullbacks of isometries. This has turned out to be more delicate than expected, because **CMet** is not a well-behaved enrichment base.

In Section 2 we introduce the notion of a locally generated category, providing examples and properties. For this, we need the concept of a λ -convenient factorization system $(\mathcal{E}, \mathcal{M})$. We discuss the main properties of λ -generated objects with respect to \mathcal{M} . We provide a recognition theorem (2.24) for locally λ -generated categories which meets the spirit of the original definition of Gabriel and Ulmer. In Section 3 we introduce the notion of a λ -nest, a sketch-like gadget that we use to recast a suitable version of Gabriel-Ulmer duality for \mathcal{M} -locally generated categories. We introduce the notion of a model of a nest, and show that the category of models of a nest is a \mathcal{M} -locally generated category. In Sections 4 and 5 we present an enriched version of Sections 2 and 3, when the enrichment base \mathcal{V} is locally λ -presentable as a closed category. Section 6 discusses the case of Banach spaces in detail, focusing on their axiomatizability via the $(\aleph_0, \mathbf{CMet})$ -nest **Ban_{fd}**.

Acknowledgement. The authors are grateful to A. Chirvasitu for a very valuable exchange of ideas on C^* -algebras.

2. Locally generated categories

Locally λ -generated categories were introduced by Gabriel and Ulmer [12] as cocomplete categories \mathcal{K} having a strong generator consisting of λ -generated objects such that every λ -generated object has only a set of strong quotients. Here, an object A is λ -generated if its hom-functor $\mathcal{K}(A, -) : \mathcal{K} \rightarrow \mathbf{Set}$ preserves λ -directed colimits of monomorphisms. A category is *locally generated* if it is locally λ -generated for some regular cardinal λ . [12] showed that locally generated categories coincide with locally presentable ones. Every locally λ -presentable category is locally λ -generated but a locally λ -generated category does not need to be locally λ -presentable. Thus the passage to locally generated categories can lower the defining cardinal λ .

The definition of a locally λ -generated category was reformulated in [1]: a cocomplete category \mathcal{K} is locally λ -generated if it has a set \mathcal{A} of λ -generated objects such that every object is a λ -directed colimit of its subobjects from \mathcal{A} . The fact that these two definitions are equivalent follows from [1] 1.70 and [12] 9.2.

A further step was made in [2] where a cocomplete category \mathcal{K} with a proper factorization system $(\mathcal{E}, \mathcal{M})$ was called *\mathcal{M} -locally λ -generated* if it has a set \mathcal{A} of λ -generated objects with respect to \mathcal{M} such that every object is a λ -directed colimit of \mathcal{A} -objects and \mathcal{M} -morphisms. Here, an object A is *λ -generated with respect to \mathcal{M}* if its hom-functor $\mathcal{K}(A, -) : \mathcal{K} \rightarrow \mathbf{Set}$ preserves λ -directed colimits of \mathcal{M} -morphisms. Again, [2] proves that a category is \mathcal{M} -locally generated for some proper factorization system $(\mathcal{E}, \mathcal{M})$ if and only if it is locally presentable. Moreover, locally λ -generated categories are \mathcal{M} -locally λ -generated ones for the factorization system (strong epimorphisms, monomorphisms). We extend this definition a little bit.

2.1. DEFINITION. Let \mathcal{K} be a category with a factorization system $(\mathcal{E}, \mathcal{M})$ and λ a regular cardinal. We say that an object A is *λ -generated with respect to \mathcal{M}* if its hom-functor $\mathcal{K}(A, -) : \mathcal{K} \rightarrow \mathbf{Set}$ preserves λ -directed colimits of \mathcal{M} -morphisms.

In what follows, \mathcal{K}^\rightarrow will denote the class of all morphisms of \mathcal{K} .

2.2. EXAMPLES. (1) For the factorization system $(\text{Iso}, \mathcal{K}^\rightarrow)$, an object is λ -generated with respect to \mathcal{K}^\rightarrow if and only if it is λ -presentable.

(2) For the factorization system $(\mathcal{K}^\rightarrow, \text{Iso})$, every object is λ -generated with respect to Iso .

(3) For the factorization system $(\text{StrongEpi}, \text{Mono})$, an object is λ -generated with respect to Mono if and only if it is λ -generated.

2.3. DEFINITION. Let λ be a regular cardinal. A factorization system $(\mathcal{E}, \mathcal{M})$ in a category \mathcal{K} will be called *λ -convenient* if

1. \mathcal{K} is \mathcal{E} -cowellpowered, i.e., if every object of \mathcal{K} has only a set of \mathcal{E} -quotients, and
2. \mathcal{M} is closed under λ -directed colimits, i.e., every λ -directed colimit of \mathcal{M} -morphisms has the property that

- (a) a colimit cocone consists of \mathcal{M} -morphisms, and
- (b) for every cocone of \mathcal{M} -morphisms, the factorizing morphism is in \mathcal{M} .

$(\mathcal{E}, \mathcal{M})$ is *convenient* if it is λ -convenient for some λ .

2.4. REMARK. If $(\mathcal{E}, \mathcal{M})$ is λ -convenient and $\lambda' \geq \lambda$ is regular then $(\mathcal{E}, \mathcal{M})$ is λ' -convenient.

2.5. DEFINITION. Let \mathcal{K} be a cocomplete category with a λ -convenient factorization system $(\mathcal{E}, \mathcal{M})$ where λ is a regular cardinal. We say that \mathcal{K} is *\mathcal{M} -locally λ -generated* if it has a set \mathcal{A} of λ -generated objects with respect to \mathcal{M} such that every object is a λ -directed colimit of objects from \mathcal{A} and morphisms from \mathcal{M} .

\mathcal{K} is called *\mathcal{M} -locally generated* if it is \mathcal{M} -locally λ -generated for some regular cardinal λ .

2.6. REMARK. In the same way as in II. (i) of the proof of [1] 1.70, we show that \mathcal{A} is dense in \mathcal{K} . Hence the canonical functor $E : \mathcal{K} \rightarrow \mathbf{Set}^{\mathcal{A}^{\text{op}}}$ (sending K to $\mathcal{K}(-, K)$ restricted on \mathcal{A}^{op}) is fully faithful. Since \mathcal{K} is cocomplete, E is a right adjoint. Moreover, E preserves λ -directed colimits of \mathcal{M} -morphisms. Consequently, in a \mathcal{M} -locally λ -generated category, λ -small limits commute with λ -directed colimits of \mathcal{M} -morphisms.

2.7. NOTATION. In what follows, \mathcal{K}_λ will denote the full subcategory of \mathcal{K} consisting of λ -generated objects with respect to \mathcal{M} and by \mathcal{E}_λ the set $\mathcal{E} \cap \mathcal{K}_\lambda^{\rightarrow}$.

2.8. EXAMPLES. (1) The factorization system $(\text{Iso}, \mathcal{K}^{\rightarrow})$ is convenient and \mathcal{K} is locally λ -presentable if and only if it is $\mathcal{K}^{\rightarrow}$ -locally λ -generated.

(2) The category \mathcal{K} is Iso-locally generated if and only if it is small. In fact, every object K of \mathcal{K} has a morphism $\coprod \mathcal{A} \rightarrow K$ where the domain is the coproduct of all objects A from \mathcal{A} . Thus $(\mathcal{K}^{\rightarrow}, \text{Iso})$ is convenient if and only if \mathcal{K} is small.

(3) A Mono-locally λ -generated category \mathcal{K} in the sense of [12] and [1] satisfies 2.5 for the factorization system (strong epi, mono). Then this factorization system is λ -convenient. Indeed, following 2.6 the canonical functor $E : \mathcal{K} \rightarrow \mathbf{Set}^{\mathcal{A}^{\text{op}}}$ is fully faithful and preserves λ -directed colimits of monomorphisms. This implies that monomorphisms are closed in \mathcal{K} under λ -directed colimits (in the sense of 2.3) because this holds in $\mathbf{Set}^{\mathcal{A}^{\text{op}}}$ and E preserves and reflects monomorphisms. Now, we can show that \mathcal{A} is closed under strong quotients. Let $e : A \rightarrow B$ be a strong epimorphism with $A \in \mathcal{A}$, $m_i : K_i \rightarrow K$ a λ -directed colimit of monomorphisms and $f : B \rightarrow K$. Since A is λ -generated, there is $g : A \rightarrow K_i$ such that $m_i g = fe$. Since m_i is a monomorphism, the diagonalization property yields $h : B \rightarrow K_i$ such that $m_i h = f$. Hence B is λ -generated. Then II. in the proof of [1] 1.70 yields that \mathcal{K} is locally presentable, hence cowellpowered. This verifies 2.3 (1), hence the (strong epi, mono) factorization system is λ -convenient. We gave this argument in detail because [1] is not accurate at this point.

(4) For a factorization system $(\text{Epi}, \text{StrongMono})$, we get StrongMono-locally λ -generated categories. Here, we have to assume that this factorization system is λ -convenient because it does not seem that we could get it for free like in (3).

(5) Any full reflective subcategory \mathcal{L} of \mathcal{K} determines a factorization system $(\mathcal{E}, \mathcal{M})$ where \mathcal{E} consists of morphisms sent by the reflector to an isomorphism (see [23]). Both (1) and (2) are special cases for $\mathcal{L} = \mathcal{K}$ or \mathcal{L} consisting just from a terminal objects.

2.9. EXAMPLE. The category **CGWH** of compactly generated weakly Hausdorff spaces and continuous maps, has a factorization system (dense maps, closed embeddings). It is \aleph_0 -convenient and [25, 3.8] shows that compact spaces are \aleph_0 -generated with respect to this factorization system. [25, 1.11] can be taken as a reformulation of the fact that compact spaces form a dense subcategory of **CGWH**. As a result, **CGWH** would be a class-locally generated category with respect to closed embeddings in the spirit of [9], if only we had defined such notion. Yet, its behavior is incredibly similar to that of a locally generated category, as the only obstruction is that \aleph_0 -generated objects form a proper class. This example finally clarifies the sense in which compact spaces are *compact* objects, as the classical terminology of algebraic topology addresses finitely presentable objects.

2.10. EXAMPLE. A similar example is given by Δ -generated spaces [11], for the same choice of a factorization system. In this case though, the category is indeed locally finitely generated with respect to closed embeddings.

2.11. NOTATION. We say that a category is *locally λ -generated* if it is \mathcal{M} -locally λ -generated for some \mathcal{M} . Locally λ -generated categories in the sense of [12] are then Mono-locally λ -generated.

2.12. LEMMA. *λ -generated objects with respect to \mathcal{M} are closed under \mathcal{E} -quotients.*

PROOF. Let $e : A \rightarrow B$ is in \mathcal{E} and $A \in \mathcal{A}$. Consider a λ -directed colimit $m_i : K_i \rightarrow K$, $i \in I$ of \mathcal{M} -morphisms and $f : B \rightarrow K$. Since A is λ -generated with respect to \mathcal{M} , there is $g : A \rightarrow K_i$ such that $m_i g = fe$. Since m_i is in \mathcal{M} , the diagonalization property yields $h : B \rightarrow K_i$ such that $m_i h = f$. Assume that $m_i h' = f$ for $h' : B \rightarrow K_j$. Then $m_i h e = m_i h' e$ and, since A is λ -generated with respect to \mathcal{M} , there is $i \leq j \in I$ such that $m_{ij} h e = m_{ij} h' e$. Hence both $m_{ij} h$ and $m_{ij} h'$ are diagonals in the commutative square

$$\begin{array}{ccc}
 A & \xrightarrow{e} & B \\
 m_{ij} h e \downarrow & & \downarrow f \\
 K_j & \xrightarrow{m_j} & K
 \end{array}$$

and thus they are equal. Hence B is λ -generated. ■

2.13. LEMMA. *In an \mathcal{M} -locally λ -generated category \mathcal{K} , λ -generated objects with respect to \mathcal{M} are closed under λ -small colimits and, up to isomorphism, form a set.*

PROOF. Let $\text{colim } A_j$ be a λ -small colimit of λ -generated objects with respect to \mathcal{M} and $\text{colim } K_i$ a λ -directed colimit of \mathcal{M} -morphisms. Since λ -directed colimits commute in **Set** with λ -small limits, we have

$$\begin{aligned} \mathcal{K}(\text{colim } A_j, \text{colim } K_i) &\cong \lim_j \mathcal{K}(A_j, \text{colim } K_i) \\ &\cong \lim_j \text{colim}_i \mathcal{K}(A_j, K_i) \\ &\cong \text{colim}_i \lim_j \mathcal{K}(A_j, K_i) \\ &\cong \text{colim}_i \mathcal{K}(\text{colim}_j A, K_i) \end{aligned}$$

Hence $\text{colim } A_j$ is λ -generated with respect to \mathcal{M} .

The second claim follows from the fact that every λ -generated object with respect to \mathcal{M} is a λ -directed colimit of \mathcal{A} -objects and \mathcal{M} -morphisms, which makes it a retract of an \mathcal{A} -object, i.e., a finite colimit of \mathcal{A} -objects. Indeed, given a retract $u : B \rightarrow A$ with a section $s : A \rightarrow B$, then s is a coequalizer of us and id_A . ■

2.14. REMARK. For the first claim, we do not need to assume that \mathcal{K} is \mathcal{M} -locally λ -generated.

2.15. NOTATION. Given a family of arrows \mathcal{E} , we call \mathcal{E}^\perp the family of arrows that are orthogonal to \mathcal{E} . This means that an arrow m belongs to \mathcal{E}^\perp if it has the unique right lifting property with respect to every arrow in \mathcal{E} . An object A is *orthogonal* to \mathcal{E} if the unique morphism from A to the terminal object is in \mathcal{E}^\perp .

2.16. LEMMA. *If \mathcal{K} is \mathcal{M} -locally λ -generated then $\mathcal{M} = (\mathcal{E}_\lambda)^\perp$.*

PROOF. Every $e : A \rightarrow B$ in \mathcal{E} is a λ -directed colimit $(a_i, b_i) : f_i \rightarrow e$ in $\mathcal{K}_\lambda^\rightarrow$. Every $f_i : A_i \rightarrow B_i$ has the factorization $f_i = m_i e_i$ with $e_i : A_i \rightarrow C_i$ in \mathcal{E} and $m_i : C_i \rightarrow B_i$ in \mathcal{M} . Then $e = \text{colim } m_i \cdot \text{colim } e_i$ where $\text{colim } e_i : A \rightarrow C$ and $\text{colim } m_i : C \rightarrow B$. Since $C = \text{colim } C_i$ is a λ -directed colimit of \mathcal{M} -morphisms and $(\mathcal{E}, \mathcal{M})$ is λ -convenient, $\text{colim } m_i$ is in \mathcal{M} and thus it is an isomorphism. Hence $e = \text{colim } e_i$. Consequently $\mathcal{M} = \mathcal{E}_\lambda^\perp$. ■

2.17. REMARK. We have shown that, in a \mathcal{M} -locally λ -generated category \mathcal{K} , \mathcal{E} is the closure $\text{colim}(\mathcal{E}_\lambda)$ of \mathcal{E}_λ under all colimits in \mathcal{K}^\rightarrow .

Conversely, if $\mathcal{M} = \mathcal{E}_\lambda^\perp$ then \mathcal{M} satisfies 2.3(2).

The proof of the next theorem follows [2] and, like [2], is based on [12]. However, [2] lacks the assumption that $(\mathcal{E}, \mathcal{M})$ is convenient.

2.18. THEOREM. *For every category \mathcal{K} equivalent are:*

1. \mathcal{K} is locally presentable,
2. \mathcal{K} is \mathcal{M} -locally generated for some convenient factorization system $(\mathcal{E}, \mathcal{M})$, and
3. \mathcal{K} is \mathcal{M} -locally generated for every convenient factorization system $(\mathcal{E}, \mathcal{M})$.

PROOF. Clearly, (3) \Rightarrow (1) \Rightarrow (2). The implication (1) \Rightarrow (3) is analogous to I. of the proof [1] 1.70.

(2) \Rightarrow (1) follows [2]. In fact, let \mathcal{K} be \mathcal{M} -locally λ -generated. Then $E : \mathcal{K} \rightarrow \mathbf{Set}^{\mathcal{A}^{\text{op}}}$ makes \mathcal{K} equivalent to a full reflective subcategory of $\mathcal{L} = \mathbf{Set}^{\mathcal{A}^{\text{op}}}$ closed under λ -directed colimits of \mathcal{M} -morphisms (see 2.6). Let \mathcal{P} consist of reflections of \mathcal{L} -objects in \mathcal{K} which are either λ -small colimits in \mathcal{L} of diagrams in \mathcal{A} or codomains of multiple pushouts in \mathcal{L} of \mathcal{E} -morphisms with a domain in \mathcal{A} . Observe that \mathcal{P} is small: the case of λ -small colimits is clear, for the multiple pushout use 2.12 and the fact that every λ -generated object with respect to \mathcal{M} is a retract of an \mathcal{A} -object. Thus, the class $\text{Ort}(\mathcal{P})$ of all \mathcal{L} -objects orthogonal to \mathcal{P} is a locally presentable category. Moreover, \mathcal{K} is closed in $\text{Ort}(\mathcal{P})$ under

1. λ -small colimits of diagrams in \mathcal{A} ,
2. multiple pushouts of \mathcal{E} -morphisms with a domain in \mathcal{A} , and
3. λ -directed colimits of \mathcal{M} -morphisms.

Consider a morphism $f : A \rightarrow L$ with A in \mathcal{A} and L in $\text{Ort}(\mathcal{P})$. Let $e_f : A \rightarrow K_f$ be the cointersection in \mathcal{K} of all \mathcal{E} -morphisms through which f factors. Following (2) above, there is $m_f : K_f \rightarrow L$ such that $f = m_f e_f$. Clearly, e_f is in \mathcal{E} and m_f is \mathcal{E} -extremal in the sense that any \mathcal{E} -morphism $K_f \rightarrow K$ through which m_f factors is an isomorphism. Moreover, this factorization is "functorial": given $f' : A' \rightarrow L$ and $h : A \rightarrow A'$ with $f = f'h$ there exists $h^* : K_f \rightarrow K_{f'}$ in \mathcal{M} with $m_f = m_{f'} h^*$. In fact, form a pushout

$$\begin{array}{ccc}
 A & \xrightarrow{e_{f'} h} & K_{f'} \\
 e_f \downarrow & & \downarrow \tilde{e}_f \\
 K_f & \xrightarrow{h^*} & K
 \end{array}$$

There is $g : K \rightarrow L$ such that $gh^* = m_f$ and $g\tilde{e}_f = m_{f'}$. Since \tilde{e}_f is in \mathcal{E} and $m_{f'}$ is \mathcal{E} -extremal, \tilde{e}_f is an isomorphism.

Every L in $\text{Ort}(\mathcal{P})$ is a λ -directed colimit $d_i : D_i \rightarrow L$ where D_i are λ -small colimits of diagrams in \mathcal{A} . Form reflections $r_i : D_i \rightarrow D'_i$ in \mathcal{K} . Since $r_i \in \mathcal{P}$, d_i factors through r_i , say, $d_i = d'_i r_i$. Since D'_i is in \mathcal{A} , we can take the factorization $d'_i = m_i e_i$ above with $m_i : D''_i \rightarrow L$. Since m_i is \mathcal{E} -extremal, d''_i is in \mathcal{M} . Thus $m_i : D''_i \rightarrow L$ is a λ -directed colimit of \mathcal{M} -morphisms, which implies that L is in \mathcal{K} .

We have proved that $\mathcal{K} = \text{Ort}(\mathcal{P})$, hence \mathcal{K} is locally presentable. ■

2.19. REMARK. (1) All implications, except (2) \Rightarrow (1), are valid for a given λ . In (2) \Rightarrow (1), λ can increase.

(2) The previous theorem sheds a light on the notion of locally generated category. Indeed, the framework of locally generated categories is not more expressive than that of locally presentable ones. Yet, as pointed out (1), it might happen that a category

is locally finitely generated without being locally finitely presentable. From the point of view of essentially algebraic theories (see 3.D in [1]), this means that some λ -ary essentially algebraic theories can be axiomatized by the data of a finite limit theory and a factorization system. Thus, the complexity of infinitary (partial) operations can be hidden under the carpet of the factorization system. From a less technical perspective, the factorization system that makes the category locally generated is very often an extremely natural one to consider. A sketch-oriented (see 2.F in [1]) interpretation of this discussion is the core motivation for the forthcoming notion of λ -nest.

2.20. REMARK. We do not know whether, given a locally λ -presentable category \mathcal{K} , one can find an \aleph_0 -convenient factorization system $(\mathcal{E}, \mathcal{M})$ such that \mathcal{K} is \mathcal{M} -locally finitely generated. Following 2.6, finite limits should commute in \mathcal{K} with directed colimits of \mathcal{M} -morphisms.

2.21. NOTATION. $\text{Gen}_\lambda(\mathcal{K})$ will denote the full subcategory of λ -generated objects with respect to \mathcal{M} in a \mathcal{M} -locally λ -generated category \mathcal{K} .

2.22. REMARK. A multiple pullback P is a limit of a diagram consisting of morphisms $f_i : A_i \rightarrow A$, $i \in I$. We can well-order I as $\{i_0, i_1, \dots, i_j \dots\}$ and form pullbacks P_j as follows: P_0 is the pullback

$$\begin{array}{ccc} P_0 & \xrightarrow{\bar{f}_0} & A_0 \\ \bar{f}_1 \downarrow & & \downarrow f_0 \\ A_1 & \xrightarrow{f_1} & A \end{array}$$

Then P_1 is the pullback

$$\begin{array}{ccc} P_1 & \xrightarrow{p_{01}} & P_0 \\ \downarrow & & \downarrow f_0 \bar{f}_1 \\ A_2 & \xrightarrow{f_2} & A \end{array}$$

We proceed by recursion and in limit steps we take limits. In this way, we transform multiple pullbacks to limits of smooth well-ordered chains (*smooth* means that in limit steps we have limits).

Conversely, a limit of a smooth chain $p_{ij} : P_j \rightarrow P_i$, $i \leq j < \alpha$ is a multiple pullback of p_{0j} , $j \in I$. Moreover, if \mathcal{K} is equipped with a factorization system $(\mathcal{E}, \mathcal{M})$ then multiple pullbacks of \mathcal{M} -morphisms correspond to limits of smooth well-ordered chains of \mathcal{M} -morphisms. Indeed, \mathcal{M} is stable under pullbacks.

2.23. COROLLARY. *Let \mathcal{K} be an \mathcal{M} -locally λ -generated category. Then \mathcal{K} is equivalent to the full subcategory of $\mathbf{Set}^{\text{Gen}_\lambda(\mathcal{K})^{\text{op}}}$ consisting of functors preserving λ -small limits and sending multiple pushouts of \mathcal{E} -morphisms to multiple pullbacks.*

PROOF. Following [1] 1.33(8), this describes $\text{Ort}(\mathcal{P})$ from the proof of 2.18. ■

The following theorem makes easier to identify \mathcal{M} -locally generated categories among cocomplete categories with a factorization system and goes back to [12].

2.24. THEOREM. *A cocomplete category \mathcal{K} equipped with a λ -convenient factorization system $(\mathcal{E}, \mathcal{M})$ is \mathcal{M} -locally λ -generated if and only if it has a strong generator formed by λ -generated objects with respect to \mathcal{M} .*

PROOF. Using 2.13 and 2.12, the closure \mathcal{A} of our strong generator under λ -small colimits and \mathcal{E} -quotients consists of λ -generated objects with respect to \mathcal{M} . Following the proof of [1] 1.11, we show that every object in \mathcal{K} is a λ -filtered colimit of \mathcal{A} -objects. Using $(\mathcal{E}, \mathcal{M})$ -factorizations, we get that every object of \mathcal{K} is a λ -directed colimit of \mathcal{A} -objects and \mathcal{M} -morphisms. ■

2.25. REMARK. If $\lambda' \geq \lambda$ and \mathcal{K} is \mathcal{M} -locally λ -generated then \mathcal{K} is \mathcal{M} -locally λ' -generated.

2.26. THEOREM. *Let \mathcal{K} be a \mathcal{M} -locally λ -generated category where $(\mathcal{E}, \mathcal{M})$ is a proper λ -convenient factorization system and let T be a monad preserving \mathcal{M} -morphisms and λ -directed colimits of \mathcal{M} -morphisms. Then, assuming Vopěnka’s principle, the category of algebras $\text{Alg}(T)$ is locally λ -generated.*

PROOF. Following [4] 3.3, $\text{Alg}(T)$ is cocomplete. Consider the adjunction $F : \mathcal{K} \rightleftarrows \text{Alg}(T) : U$ and put $\mathcal{M}' = U^{-1}(\mathcal{M})$. Using [5] 4.3.2, we get that \mathcal{M}' is closed under λ -directed colimits. Following [21] 11.1.5, $\mathcal{M}' = F(\mathcal{E})^\perp$. Since $\mathcal{E} = \text{colim } \mathcal{E}_\lambda$ (see 2.17), we have $\text{colim } F(\mathcal{E}) = \text{colim}(F(\mathcal{E}_\lambda))$. U is clearly conservative (and faithful), thus F maps the dense generator $\text{Gen}_\lambda(\mathcal{K})$ to a strong generator \mathcal{G} in $\text{Alg}(T)$. Let \mathcal{A} be the closure of \mathcal{G} under λ -small colimits. Following 2.14, \mathcal{A} is dense in \mathcal{K} . Assuming Vopěnka’s principle, $\text{Alg}(T)$ is locally presentable (see [1] 6.14). Hence, following [11] 2.2, $(\text{colim } F(\mathcal{E}), \mathcal{M}')$ is a factorization system on $\text{Alg}(T)$. Since $\text{Alg}(T)$ is cowellpowered (see [1] 1.58) and $\text{colim } F(\mathcal{E}) \subseteq \text{Epi}$, this factorization system $(\text{colim } F(\mathcal{E}), \mathcal{M}')$ is λ -convenient. We conclude the proof by the previous theorem and 3.11. ■

2.27. REMARK. We do not know whether Vopěnka’s principle is really needed. This is related to the Open Problem 3 in [1]. In fact, let \mathcal{L} be a full reflective subcategory of a locally λ -presentable category \mathcal{K} closed under λ -directed colimits of monomorphisms. Then the monad $T = FG$, where $G : \mathcal{L} \rightarrow \mathcal{K}$ is the inclusion and F its left adjoint, preserves λ -directed colimits of monomorphisms. Since $\mathcal{L} \cong \text{Alg}(T)$, 2.26 without Vopěnka’s principle would yield a positive solution of the Open Problem 3.

3. Extended Gabriel-Ulmer duality

The Gabriel-Ulmer duality is a contravariant biequivalence between categories with λ -small limits and locally λ -presentable categories (see [12] 7.11 or [1] 1.45). We are going to extend this duality to \mathcal{M} -locally λ -generated categories. It will also cover the

Gabriel-Ulmer duality for Mono-locally λ -generated categories. In order to do so, we will introduce the notion of a nest.

3.1. DEFINITION. A λ -nest is a small category \mathcal{A} equipped with a factorization system $(\mathcal{E}_{\mathcal{A}}, \mathcal{M}_{\mathcal{A}})$ and having λ -small limits and multiple pullbacks of $\mathcal{M}_{\mathcal{A}}$ -morphisms.

3.2. REMARK. λ -nests \mathcal{A} with $(\text{Iso}, \mathcal{A}^{\rightarrow})$ are precisely small categories with λ -small limits. λ -nests with $(\text{StrongEpi}, \text{Mono})$ are precisely small "echt" λ -complete categories of [12].

3.3. EXAMPLE. $\text{Gen}_{\lambda}(\mathcal{K})^{\text{op}}$ is a λ -nest for every \mathcal{M} -locally λ -generated category \mathcal{K} . This follows from 2.13, 2.12 and the fact that $(\mathcal{M}_{\mathcal{A}} \cap \mathcal{A}^{\rightarrow}, \mathcal{E}_{\mathcal{A}} \cap \mathcal{A}^{\rightarrow})$ is a factorization system on $\text{Gen}_{\lambda}(\mathcal{K})^{\text{op}}$.

3.4. NOTATION. For a λ -nest \mathcal{A} , $\text{Mod}_{\lambda}(\mathcal{A})$ denotes the category of all *models*, i.e., of functors $\mathcal{A} \rightarrow \mathbf{Set}$ preserving λ -small limits and multiple pullbacks of \mathcal{M} -morphisms.

3.5. LEMMA. $\text{Mod}_{\lambda}(\mathcal{A})$ is a locally presentable category equipped with a factorization system $(\mathcal{E}, \mathcal{M})$.

PROOF. $\text{Mod}_{\lambda}(\mathcal{A})$ is a full subcategory of $\mathbf{Set}^{\mathcal{A}}$ and the codomain restriction of the Yoneda embedding $Y : \mathcal{A}^{\text{op}} \rightarrow \mathbf{Set}^{\mathcal{A}}$ is a full embedding of \mathcal{A}^{op} to $\mathbf{Set}^{\mathcal{A}}$. Following [1] 1.51, $\text{Mod}_{\lambda}(\mathcal{A})$ is locally presentable. The factorization system on \mathcal{A}^{op} is $(\mathcal{M}_{\mathcal{A}}, \mathcal{E}_{\mathcal{A}})$ and the factorization system $(\mathcal{E}, \mathcal{M})$ is given, following [11] 2.2, as: $\mathcal{E} = \text{colim}(\mathcal{M}_{\mathcal{A}})$ and $\mathcal{M} = \mathcal{M}_{\mathcal{A}}^{\perp}$. ■

3.6. THEOREM. $\text{Mod}_{\lambda}(\mathcal{A})$ is a \mathcal{M} -locally λ -generated category for every λ -nest \mathcal{A} .

PROOF. I. At first, we will show that $\mathcal{K} = \text{Mod}_{\lambda}(\mathcal{A})$ is closed in $\mathcal{L} = \mathbf{Set}^{\mathcal{A}}$ under λ -directed colimits of \mathcal{M} -morphisms. Following [1] 1.33(8), $\mathcal{K} = \text{Ort}(\mathcal{P})$ where \mathcal{P} is defined as in the proof of 2.18. This means that \mathcal{P} consists of reflections of \mathcal{L} -objects in \mathcal{K} which are either λ -small colimits of diagrams in \mathcal{A}^{op} or codomains of multiple pushouts of $\mathcal{M}_{\mathcal{A}}$ -morphisms in \mathcal{A}^{op} .

Let $k_i : K_i \rightarrow K$ be a colimit in \mathcal{L} of a λ -directed diagram of \mathcal{M} -morphisms in \mathcal{K} . Let $r : \text{colim } A_j \rightarrow A$ be a reflection of a λ -small colimit in \mathcal{L} of a diagram in \mathcal{A}^{op} ; its reflection in \mathcal{K} lies in \mathcal{A}^{op} . It is easy to see that objects orthogonal to r are closed under λ -directed colimits; in fact, these objects correspond to functors $\mathcal{A} \rightarrow \mathbf{Set}$ preserving λ -small limits.

Let $r : P \rightarrow A$ be a reflection of a multiple pushout of $\mathcal{M}_{\mathcal{A}}$ -morphisms. Alternatively, P can be seen as a colimit of a well ordered smooth chain

$$P_0 \xrightarrow{p_{01}} P_1 \xrightarrow{p_{12}} P_2 \xrightarrow{p_{23}} \dots$$

of $\mathcal{M}_{\mathcal{A}}$ -morphisms. Let $f : P \rightarrow K$. There exists i_0 and $g_0 : P_0 \rightarrow K_{i_0}$ such that $k_{i_0}g_0 = fp_0$. There exists $i_1 > i_0$ and $g'_1 : P_1 \rightarrow K_{i_1}$ such that $k_{i_0i_1}g_0 = g'_1p_{01}$. Since $k_{i_0i_1} \in \mathcal{M}_{\mathcal{A}}^{\perp}$ and $p_{01} \in \mathcal{M}_{\mathcal{A}}$, there is $g_1 : P_1 \rightarrow K_{i_0}$ such that $g_1p_{01} = g_0$ and $k_{i_0i_1}g_1 = g'_1$. Continuing this procedure by taking colimits in limit steps, we get a cocone $g_j : P_j \rightarrow K_{i_0}$ inducing $g : P \rightarrow K_{i_0}$ such that $k_{i_0}g = f$. There is $h : A \rightarrow K_{i_0}$ with $hr = g$. Hence $k_{i_0}hr = f$. The uniqueness of this extension follows from A being in \mathcal{A}^{op} .

II. From I., it follows that every object of \mathcal{A}^{op} is λ -generated with respect to \mathcal{M} in \mathcal{K} . Following 3.5 and 2.17, \mathcal{M} satisfies 2.3(2). In the same way as in 2.12, we show that every \mathcal{E} -quotient of an \mathcal{A}^{op} -object is λ -generated with respect to \mathcal{M} . Since every λ -generated object with respect to \mathcal{M} is a retract of an \mathcal{A}^{op} -object (cf. 2.13) and \mathcal{A}^{op} is closed under retracts, $\mathcal{A}^{\text{op}} = \text{Gen}_\lambda(\mathcal{K})$. Using $(\mathcal{E}, \mathcal{M})$ factorizations, we show that every object in \mathcal{K} is a λ -directed colimit of \mathcal{A}^{op} -objects.

It remains to show that \mathcal{K} is \mathcal{E} -cowellpowered. Let $e : K \rightarrow L$ be an \mathcal{E} -quotient of K . Then e is a λ -directed colimit of \mathcal{E} -morphisms $e_i : K_i \rightarrow L_i$ where K_i and L_i are in \mathcal{A}^{op} . Since there is only set of expressions of K as a λ -directed colimit of \mathcal{A}^{op} -objects, there is only a set of \mathcal{E} -quotients of K . ■

3.7. REMARK. We have

$$\mathcal{A} \simeq (\text{Gen}_\lambda(\text{Mod}_\lambda(\mathcal{A})))^{\text{op}}$$

for every λ -nest \mathcal{A} , and

$$\mathcal{K} \simeq \text{Mod}_\lambda(\text{Gen}_\lambda(\mathcal{K})^{\text{op}})$$

for every \mathcal{M} -locally λ -generated category \mathcal{K} .

Note that these equivalences also include the corresponding factorization systems. In the first case, $(\text{colim}(\mathcal{M}_\mathcal{A}), \mathcal{M}_\mathcal{A}^\perp)$ restricts to $(\mathcal{M}_\mathcal{A}, \mathcal{E}_\mathcal{A})$ (see 3.5) and, in the second case, it follows from 2.17.

3.8. REMARK. For a λ -nest \mathcal{A} , $\text{Mod}_\lambda(\mathcal{A})$ is a full subcategory of the free completion $\text{Ind}_\lambda(\mathcal{A})$ of \mathcal{A} under λ -directed colimits. Indeed, the latter category consists of functors $\mathcal{A}^{\text{op}} \rightarrow \mathbf{Set}$ preserving λ -small limits.

3.9. DEFINITION. A *morphism of locally λ -generated categories* $R : \mathcal{K} \rightarrow \mathcal{L}$ is a right adjoint preserving \mathcal{M} -morphisms and λ -directed colimits of them.

3.10. REMARK. R preserves \mathcal{M} if and only if its left adjoint $L : \mathcal{L} \rightarrow \mathcal{K}$ preserves \mathcal{E} (see [21] 11.1.5).

3.11. LEMMA. *Let $R : \mathcal{K} \rightarrow \mathcal{L}$ be a morphism of locally λ -generated categories. Then its left adjoint L preserves λ -generated objects with respect to \mathcal{M} .*

PROOF. Let A be λ -generated with respect to \mathcal{M} in \mathcal{L} . Consider a directed colimit $\text{colim } K_i$ of \mathcal{M} -morphisms in \mathcal{K} . Then

$$\begin{aligned} \mathcal{K}(LA, \text{colim } K_i) &\cong \mathcal{L}(A, R \text{colim } K_i) \\ &\cong \mathcal{L}(A, \text{colim } RK_i) \\ &\cong \text{colim } \mathcal{L}(A, RK_i) \\ &\cong \text{colim } \mathcal{K}(LA, K_i). \end{aligned}$$

■

3.12. DEFINITION. A *morphism of λ -nests* $F : \mathcal{A} \rightarrow \mathcal{B}$ is a functor preserving λ -small limits, \mathcal{M} -morphisms and multiple pullbacks of them.

3.13. NOTATION. Let \mathbf{LG}_λ be the 2-category of locally λ -generated categories and \mathbf{N}_λ be the 2-category of λ -nests, in the both cases 2-cells are natural transformations.

3.14. CONSTRUCTION. [The functor Gen_λ] Given a locally λ -generated category \mathcal{K} , we have defined $\text{Gen}_\lambda(\mathcal{K})^{\text{op}}$ to be the opposite category of its λ -generated objects. It is easy to see that this construction is (contravariantly) functorial. Indeed, given a morphism of locally λ -generated categories $R : \mathcal{K} \rightarrow \mathcal{L}$, its left adjoint L restricts to λ -generated objects

$$L : \text{Gen}_\lambda(\mathcal{L}) \rightarrow \text{Gen}_\lambda(\mathcal{K})$$

by 3.11, and passing to the opposite category, is a morphism of λ -nests because of 2.12 and 2.13.

3.15. CONSTRUCTION. [The functor Mod_λ] Given a λ -nest \mathcal{A} , we have seen that the category $\text{Mod}_\lambda(\mathcal{A})$ is locally λ -generated. We will extend this construction to a (contravariant) functor. Given a morphism of λ -nests $F : \mathcal{A} \rightarrow \mathcal{B}$, the functor $\text{Mod}_\lambda(F)$ sends H from $\text{Mod}_\lambda(\mathcal{B})$ to HF . This functor is the domain restriction of the functor $\text{Ind}_\lambda(\mathcal{B}) \rightarrow \text{Ind}_\lambda(\mathcal{A})$ given, again, by precompositions with F . The latter functor has the left adjoint $\text{Ind}_\lambda(F) : \text{Ind}_\lambda(\mathcal{A}) \rightarrow \text{Ind}_\lambda(\mathcal{B})$. The domain restriction $L(F)$ of this left adjoint is a left adjoint to $\text{Mod}_\lambda(F)$. The functor $L(F)$ preserves \mathcal{E} -morphisms because

$$L(F)(\mathcal{E}_{\text{Mod}_\lambda(\mathcal{A})}) = L(F)(\text{colim } \mathcal{M}_\mathcal{A}) \subseteq \text{colim}(F(\mathcal{M}_\mathcal{A})) \subseteq \mathcal{E}_{\text{Mod}_\lambda(\mathcal{B})}.$$

Thus $\text{Mod}_\lambda(F)$ preserves \mathcal{M} -morphisms and, therefore, it is a morphism of λ -generated categories.

3.16. THEOREM.

$$\text{Gen}_\lambda : \mathbf{LG}_\lambda \rightleftarrows \mathbf{N}_\lambda^{\text{op}} : \text{Mod}_\lambda$$

is a dual biequivalence between locally λ -generated categories and λ -nests.

PROOF. It follows from 3.7 and 3.15. ■

3.17. REMARK. (1) Our duality restricts to the standard Gabriel-Ulmer duality between locally λ -presentable categories and small categories with λ -small limits (for $\mathcal{M} = \text{Iso}$) and, a little bit forgotten, Gabriel-Ulmer duality for locally λ -generated categories (see [12] 9.8). In the second case $\mathcal{M} = \text{Mono}$ and the dual is formed by small categories with λ -small limits and pullbacks of strong monomorphisms.

On the other hand, our extended Gabriel-Ulmer duality is a restriction of the standard one (see 3.15).

(2) Like the Gabriel-Ulmer duality (see [19]), our duality is given by the category \mathbf{Set} being both a large λ -nest and a locally λ -generated category. Clearly, models of a λ -nest \mathcal{A} are λ -nest morphisms $\mathcal{A} \rightarrow \mathbf{Set}$. Conversely, a morphism $U : \mathcal{K} \rightarrow \mathbf{Set}$ of locally

λ -generated categories is uniquely determined by its left adjoint $F : \mathbf{Set} \rightarrow \mathcal{K}$, these restrictions uniquely correspond to objects in $\text{Gen}_\lambda(\mathcal{K})$.

(3) [8] generalized the Gabriel-Ulmer duality to certain limit doctrines. This is based on the commutation of certain limits and colimits in \mathbf{Set} . But our duality (even that for Mono-locally λ -generated categories) does not fall under this scope.

4. Enriched locally generated categories

In what follows, \mathcal{V} will be a complete and cocomplete symmetric monoidal closed category. We will work with \mathcal{V} -categories and under a λ -directed colimit we will mean a conical λ -directed one. \mathcal{V} -factorization systems were introduced in [10] and their theory was later developed by Lucyshyn-Wright in [18]. We refer to [18] for the main definitions and notations.

4.1. DEFINITION. Let \mathcal{K} be a \mathcal{V} -category with a \mathcal{V} -factorization system $(\mathcal{E}, \mathcal{M})$ and λ a regular cardinal. We say that an object A is λ -generated with respect to \mathcal{M} if its hom-functor $\mathcal{K}(A, -) : \mathcal{K} \rightarrow \mathcal{V}$ preserves λ -directed colimits of \mathcal{M} -morphisms.

4.2. REMARK. In a tensored \mathcal{V} -category \mathcal{K} , \mathcal{V} -factorization systems are precisely factorization systems $(\mathcal{E}, \mathcal{M})$ such that \mathcal{E} is closed under tensors (see [18] 5.7).

4.3. DEFINITION. A \mathcal{V} -factorization system $(\mathcal{E}, \mathcal{M})$ in a tensored \mathcal{V} -category \mathcal{K} is called λ -convenient if it is λ -convenient as an (ordinary) factorization system.

4.4. DEFINITION. Let \mathcal{K} be a cocomplete \mathcal{V} -category with a λ -convenient \mathcal{V} -factorization system $(\mathcal{E}, \mathcal{M})$ where λ is a regular cardinal. We say that \mathcal{K} is \mathcal{M} -locally λ -generated if it has a set \mathcal{A} of λ -generated objects with respect to \mathcal{M} such that every object is a λ -directed colimit of objects from \mathcal{A} and morphisms from \mathcal{M} .

\mathcal{K} is called \mathcal{M} -locally generated if it is \mathcal{M} -locally λ -generated for some regular cardinal λ .

4.5. EXAMPLE. For a cocomplete \mathcal{V} -category \mathcal{K} , $(\text{Iso}, \mathcal{K}^\rightarrow)$ is a convenient \mathcal{V} -factorization system. \mathcal{K}^\rightarrow -locally λ -generated categories are locally presentable \mathcal{V} -categories in the sense of [17].

4.6. REMARK. In 4.4, \mathcal{A} is dense in \mathcal{K} , i.e., the canonical \mathcal{V} -functor $E : \mathcal{K} \rightarrow \mathcal{V}^{\mathcal{A}^{\text{op}}}$ is fully faithful. In fact, E preserves λ -directed colimits of \mathcal{M} -morphisms and, then, the result follows from [16], 5.19.

4.7. REMARK. For a \mathcal{V} -category \mathcal{K} , the arrow category \mathcal{K}^\rightarrow is a \mathcal{V} -category with $\mathcal{K}^\rightarrow(f, g)$ defined by the following pullback in \mathcal{V}

$$\begin{array}{ccc} \mathcal{K}^\rightarrow(f, g) & \longrightarrow & \mathcal{K}(A, C) \\ \downarrow & & \downarrow \kappa(A, g) \\ \mathcal{K}(B, D) & \xrightarrow{\kappa(f, D)} & \mathcal{K}(A, D) \end{array}$$

where $f : A \rightarrow B$ and $g : C \rightarrow D$.

4.8. PROPOSITION. *Assume that pullbacks commute with λ -directed colimits in \mathcal{V} . Then, for a \mathcal{M} -locally λ -generated \mathcal{V} -category \mathcal{K} , the \mathcal{V} -category \mathcal{K}^\rightarrow is \mathcal{M}_\rightarrow -locally λ -generated.*

PROOF. Following [21] 13.1, \mathcal{K}^\rightarrow is tensored and cotensored. Since \mathcal{K}^\rightarrow has limits and colimits (calculated pointwise), it is complete and cocomplete (see [5] 6.6.16). The factorization system $(\mathcal{E}_\rightarrow, \mathcal{M}_\rightarrow)$ on \mathcal{K}^\rightarrow is defined pointwise from that on \mathcal{K} and is clearly λ -convenient. We will show that any morphism $f : A \rightarrow B$ with A and B λ -generated with respect to \mathcal{M} is λ -generated with respect to \mathcal{M}_\rightarrow .

Consider a λ -directed colimit $g_i \rightarrow g$ on \mathcal{M}_\rightarrow -morphisms in \mathcal{K}^\rightarrow . We have pullbacks

$$\begin{array}{ccc} \mathcal{K}^\rightarrow(f, g_{g_i}) & \longrightarrow & \mathcal{K}(A, C_i) \\ \downarrow & & \downarrow \kappa(A, g_i) \\ \mathcal{K}(B, D_i) & \xrightarrow{\kappa(f, D_i)} & \mathcal{K}(A, D_i) \end{array}$$

Since pullbacks commute with λ -directed colimits in \mathcal{V} ,

$$\mathcal{K}^\rightarrow(f, g) \cong \operatorname{colim} \mathcal{K}^\rightarrow(f, g_i).$$

Clearly, every h in \mathcal{K}^\rightarrow is a λ -directed colimit of λ -generated objects with respect to \mathcal{M}_\rightarrow and \mathcal{M}_\rightarrow -morphisms. ■

4.9. LEMMA. *Assume that λ -small (conical) limits commute in \mathcal{V} with λ -directed colimits. Then, in an \mathcal{M} -locally λ -generated \mathcal{V} -category, λ -generated objects with respect to \mathcal{M} are closed under λ -small (conical) colimits.*

PROOF. The proof for λ -small conical colimits is the same as that of 2.13 and the proof for λ -small weighted limits is analogous (cf. [6] 3.2). ■

4.10. REMARK. For the first claim, we do not need to assume that \mathcal{K} is \mathcal{M} -locally λ -generated.

4.11. LEMMA. *Assume that finite conical limits commute with λ -directed colimits in \mathcal{V} . Then, in an \mathcal{M} -locally λ -generated \mathcal{V} -category, λ -generated objects with respect to \mathcal{M} are closed under \mathcal{E} -quotients.*

PROOF. Let $e : A \rightarrow B$ is in \mathcal{E} and $A \in \mathcal{A}$. Express B as a λ -directed colimit $m_i : B_i \rightarrow B$, $i \in I$, of \mathcal{A} -objects and \mathcal{M} -morphisms. Form pullbacks

$$\begin{array}{ccc} P_i & \xrightarrow{\bar{e}} & B_i \\ \bar{m}_i \downarrow & & \downarrow m_i \\ A & \xrightarrow{e} & B \end{array}$$

Since pullbacks commute with λ -directed colimits in \mathcal{V} , they do it in $\mathcal{V}^{\mathcal{A}^{\text{op}}}$ as well. Since E preserves pullbacks and λ -directed colimits of \mathcal{M} -morphisms, pullbacks commute with λ -directed colimits of \mathcal{M} -morphisms in \mathcal{K} . Hence $\bar{m}_i : P_i \rightarrow A$ is a λ -directed colimit. Since \mathcal{M} is λ -convenient, $m_i \in \mathcal{M}$ for every $i \in I$. Hence $\bar{m}_i \in \mathcal{M}$ for every $i \in I$ and, thus, $\bar{m}_{ij} : P_i \rightarrow P_j$ are in \mathcal{M} for every $i < j \in I$. Since A is λ -generated with respect to \mathcal{M} , m_{i_0} splits for some $i_0 \in I$. Thus there exists $s : A \rightarrow P_{i_0}$ such that $\bar{m}_{i_0}s = \text{id}_A$. We have

$$m_{i_0}\bar{e}s = e\bar{m}_{i_0}s = e.$$

The commutative square

$$\begin{array}{ccc} A & \xrightarrow{e} & B \\ \bar{e}s \downarrow & & \downarrow \text{id}_B \\ B_{i_0} & \xrightarrow{m_{i_0}} & B \end{array}$$

has the diagonal $t : B \rightarrow B_{i_0}$ making B a retract of B_{i_0} . Since t is a coequalizer of $\text{id}_{B_{i_0}}$ and tm_{i_0} , 4.9 implies that B is λ -generated with respect to \mathcal{M} . ■

4.12. LEMMA. *If \mathcal{K} is \mathcal{M} -locally λ -generated \mathcal{V} -category then $\mathcal{M} = (\mathcal{E}_\lambda)^{\perp \mathcal{V}}$.*

PROOF. It follows from 4.2, 2.16 and [18] 5.4. ■

4.13. REMARK. (1) Given a small \mathcal{V} -category, every \mathcal{V} -functor $H : \mathcal{A} \rightarrow \mathcal{V}$ is a weighted colimit of representable \mathcal{V} -functors. If \mathcal{V} is locally λ -presentable as a closed category (see [17] 5.5) then, for a small category \mathcal{A} with λ -small limits, any functor $H : \mathcal{A} \rightarrow \mathcal{V}$ preserving λ -small limits is a λ -filtered (and thus λ -directed) conical colimit of representable functors (see [6] 4.5). In such a \mathcal{V} , λ -small limits commute with λ -directed colimits (see [6] 2.4).

(2) For a general \mathcal{V} , let $H : \mathcal{A} \rightarrow \mathcal{V}$ preserve λ -small conical limits and consider the category $Y(\mathcal{A}^{\text{op}}) \downarrow H$ of representable functors over H . This category is λ -filtered and let H_* be its colimit in $\mathcal{V}^{\mathcal{A}}$ and $\gamma : H_* \rightarrow H$ be the comparison morphism. If \mathcal{A} is a λ -nest then, due to $(\mathcal{E}, \mathcal{M})$ factorizations in \mathcal{A} , the category $Y(\mathcal{A}^{\text{op}}) \downarrow H$ has a cofinal subcategory \mathcal{D} whose morphisms are $Y(m) : Y(A) \rightarrow Y(B)$ where m are in \mathcal{M} . \mathcal{D} is λ -directed and H_* is its colimit of the projection $D : \mathcal{D} \rightarrow \mathcal{V}^{\mathcal{A}}$ with a cocone $\varphi_d : Dd \rightarrow H_*$.

Assume that \mathcal{V} is equipped with a λ -convenient factorization system $(\mathcal{E}_\mathcal{V}, \mathcal{M}_\mathcal{V})$ and that hom-functors $\mathcal{A}(A, -) : \mathcal{A} \rightarrow \mathcal{V}$ send \mathcal{M} -morphisms to $\mathcal{M}_\mathcal{V}$ -morphisms. Then, for

every A in \mathcal{A} , we have a λ -directed colimit $(\varphi_d)_A : Dd(A) \rightarrow H_*(A)$ of $\mathcal{M}_{\mathcal{V}}$ -morphisms in \mathcal{V} . Since the factorization system $(\mathcal{E}, \mathcal{M})$ is λ -convenient, $(\varphi_d)_A$ are in \mathcal{M} .

Assume that \mathcal{V} is $\mathcal{M}_{\mathcal{V}}$ -locally λ -generated and that \mathcal{A} has cotensors with λ -generated objects V with respect to $\mathcal{M}_{\mathcal{V}}$. If H preserves these cotensors then the argument from [6] 4.5 yields that γ is an isomorphism.

4.14. THEOREM. *Let \mathcal{V} be locally λ -presentable as a closed category. Then, for every \mathcal{V} -category \mathcal{K} equivalent are:*

1. \mathcal{K} is locally presentable,
2. \mathcal{K} is \mathcal{M} -locally generated for some convenient \mathcal{V} -factorization system $(\mathcal{E}, \mathcal{M})$, and
3. \mathcal{K} is \mathcal{M} -locally generated for every convenient \mathcal{V} -factorization system $(\mathcal{E}, \mathcal{M})$.

PROOF. We proceed like in 2.18. In the implication (2) \Rightarrow (1), we take $\text{Gen}_{\lambda}(\mathcal{K})$ for \mathcal{A} . Following 4.9, \mathcal{P} is closed under finite tensors. Analogously as in 4.12, we get that $\mathcal{P}^{\perp} = \mathcal{P}^{\perp \mathcal{V}}$. Hence $\text{Ort}(\mathcal{P})$ consists of objects \mathcal{V} -orthogonal to \mathcal{P} . Finally, following 4.13(1), every L in $\text{Ort}(\mathcal{P})$ is a λ -directed colimit of objects from $\text{Gen}_{\lambda}(\mathcal{K})$. ■

4.15. COROLLARY. *Let \mathcal{V} be locally λ -presentable as a closed category and \mathcal{K} be an \mathcal{M} -locally λ -generated \mathcal{V} -category. Then \mathcal{K} is equivalent to the full subcategory of $\mathcal{V}^{\text{Gen}_{\lambda}(\mathcal{K})^{\text{op}}}$ consisting of \mathcal{V} -functors preserving λ -small limits and sending multiple pushouts of \mathcal{E} -morphisms to multiple pullbacks.*

PROOF. We proceed analogously as in 2.23. Note that [1] 1.33(8) is true in the enriched setting too but m_i there should be

$$m_i : \text{colim } \text{hom}(Dd, -) \rightarrow \text{hom}(\text{lim } Dd, -).$$

■

4.16. NOTATION. Again, we say that a \mathcal{V} -category is *locally λ -generated* if it is \mathcal{M} -locally λ -generated for some \mathcal{M} .

4.17. THEOREM. *Assume that λ -small conical limits commute in \mathcal{V} with λ -directed colimits. Then a cocomplete \mathcal{V} -category \mathcal{K} equipped with a λ -convenient \mathcal{V} -factorization system $(\mathcal{E}, \mathcal{M})$ is \mathcal{M} -locally λ -generated if and only if its underlying category \mathcal{K}_0 has a strong generator formed by λ -generated objects with respect to \mathcal{M} .*

PROOF. Using 4.9 and 4.11, the closure \mathcal{A} of our strong generator under λ -small conical colimits and \mathcal{E} -quotients consists of λ -generated objects with respect to \mathcal{M} . Then we follow the proof of 2.24. ■

5. Enriched Gabriel-Ulmer duality

5.1. DEFINITION. A (λ, \mathcal{V}) -nest is a small \mathcal{V} -category \mathcal{A} equipped with a \mathcal{V} -factorization system $(\mathcal{E}_{\mathcal{A}}, \mathcal{M}_{\mathcal{A}})$ and having λ -small limits and multiple pullbacks of \mathcal{M} -morphisms.

5.2. EXAMPLE. $\text{Gen}_{\lambda}(\mathcal{K})^{\text{op}}$ is a (λ, \mathcal{V}) -nest for every \mathcal{M} -locally λ -generated \mathcal{V} -category \mathcal{K} . This follows from 4.9, 4.11 and the fact that $(\mathcal{M}_{\mathcal{A}} \cap \mathcal{A}^{\rightarrow}, \mathcal{E}_{\mathcal{A}} \cap \mathcal{A}^{\rightarrow})$ is a \mathcal{V} -factorization system on $\text{Gen}_{\lambda}(\mathcal{K})^{\text{op}}$.

5.3. REMARK. Let \mathcal{V} be locally λ -presentable as a closed category. Then a small \mathcal{V} -category \mathcal{A} is a (λ, \mathcal{V}) -nest if and only if it is a λ -nest and $\mathcal{M}_{\mathcal{A}}$ is closed under λ -small cotensors.

We get this analogously to 4.2 applied to \mathcal{A}^{op} .

5.4. NOTATION. For a (λ, \mathcal{V}) -nest \mathcal{A} , $\text{Mod}_{\lambda}(\mathcal{A})$ denotes the category of all \mathcal{V} -functors $\mathcal{A} \rightarrow \mathcal{V}$ preserving λ -small limits and multiple pullbacks of \mathcal{M} -morphisms.

5.5. LEMMA. *Let \mathcal{V} be locally λ -presentable as a closed category. Then $\text{Mod}_{\lambda}(\mathcal{A})$ is a locally presentable \mathcal{V} -category equipped with a \mathcal{V} -factorization system $(\mathcal{E}, \mathcal{M})$.*

PROOF. We proceed like in 3.5 with [1] 1.51 replaced by [6] 7.3. We also use 4.2. ■

5.6. THEOREM. *Let \mathcal{V} be locally λ -presentable as a closed category. Then $\text{Mod}_{\lambda}(\mathcal{A})$ is a \mathcal{M} -locally λ -generated \mathcal{V} -category for every (λ, \mathcal{V}) -nest \mathcal{A} .*

PROOF. It follows from 5.5, 3.6 and 4.2. ■

5.7. DEFINITION. A *morphism of λ -generated \mathcal{V} -categories* $R : \mathcal{K} \rightarrow \mathcal{L}$ is a right \mathcal{V} -adjoint preserving \mathcal{M} -morphisms and λ -directed colimits of them.

A *morphism of (λ, \mathcal{V}) -nests* $F : \mathcal{A} \rightarrow \mathcal{B}$ is a \mathcal{V} -functor preserving λ -small limits, \mathcal{M} -morphisms and multiple pullbacks of them.

5.8. THEOREM. *Let \mathcal{V} be locally λ -presentable as a closed category. Then*

$$\text{Gen}_{\lambda} : \mathbf{LG}_{\lambda} \rightleftarrows \mathbf{N}_{\lambda}^{\text{op}} : \text{Mod}_{\lambda}$$

is a dual biequivalence between locally λ -generated \mathcal{V} -categories and (λ, \mathcal{V}) -nests.

PROOF. It follows from 3.16, 5.2 and 5.6. ■

5.9. REMARK. (1) Our duality restricts to the enriched Gabriel-Ulmer duality between locally λ -presentable \mathcal{V} -categories and small \mathcal{V} -categories with λ -small limits given in [17].

(2) 3.11 and 3.17 (2) apply to the enriched case as well.

6. Banach spaces

Let **CMet** be the category of (generalized) metric spaces and nonexpanding maps. This category is symmetric monoidal closed and locally \aleph_1 -presentable as a closed category (see [3] 2.3(2) and 4.5(2)). But it is not locally \aleph_0 -presentable. In fact, only the empty space is \aleph_0 -presentable in **CMet** (see [3] 2.7 (1)). **CMet** has a strong generator consisting of a one-point space 1 and of two-point spaces 2_ε where the two points have the distance $\varepsilon > 0$.

The category **Ban** of Banach spaces and linear maps of norm ≤ 1 is enriched over **CMet**. Moreover, it is locally \aleph_1 -presentable **CMet**-category (see [3] 6.3).

6.1. REMARK. (1) Epimorphisms in **CMet** or **Ban** coincide with dense maps. See [20] 1.15 for **Ban** and the argument for **CMet** is analogous. Both in **CMet** and **Ban**, there is a factorization system $(\mathcal{E}, \mathcal{M})$ where \mathcal{E} consists of dense maps and \mathcal{M} of isometries (see [3] 3.16(2)). Hence isometries coincide, both in **CMet** and in **Ban** with strong monomorphisms. Both **CMet** and **Ban** are \mathcal{E} -cowellpowered and, from the description of directed colimits (see [3] 2.5), it follows that $(\mathcal{E}, \mathcal{M})$ is, in the both cases, \aleph_0 -convenient.

(2) Both in **CMet** and **Ban**, \aleph_0 -generated objects with respect to \mathcal{M} coincide with approximately \aleph_0 -generated objects in the sense of [3] (see 5.11(3)). In **CMet**, approximately \aleph_0 -generated finite metric spaces are precisely discrete ones (see [3] 5.18 and 5.19).

(3) **Ban** is \mathcal{M} -locally \aleph_0 -generated (see [3] 7.8). Every finite-dimensional Banach space is approximately \aleph_0 -generated ([3] 7.6).

(4) The concept of a finite weight does not make sense in **CMet** because only \emptyset is \aleph_0 -presentable. Thus, under finite limits in **CMet**, we understand finite conical limits, cotensors with finite metric spaces, and their combinations. Here, cotensors with finite metric spaces can be replaced by ε -pullbacks (see [3] 4.6). But in **CMet**, \aleph_0 -generated objects with respect to \mathcal{M} are not closed under these finite colimits (see [3] 5.20). Hence, following 4.9 and [3] 5.20 and 4.1(4), finite limits do not commute with directed colimits in **CMet**.

6.2. REMARK. A metric space is called *convex* if for every points x and y there is a point z such that $d(x, z) + d(z, y) = d(x, y)$. A subset S of a metric space A is called a *metric segment* if for every two distinct points $a \neq b$ there is an isometry $f : [0, d(a, b)] \rightarrow A$ from a closed interval on \mathbb{R} such that $f(0) = a$, $f(d(a, b)) = b$ and $f([0, d(a, b)]) = S$. A complete metric space is convex if and only if every distinct points are connected by a metric segment (see [7]). The proof consists in creating a dense set of points between a and b and taking its completion. Hence it also applies to $d(a, b) = \infty$ where the interval is $[0, \infty]$ in \mathbb{R} with ∞ added.

6.3. LEMMA. *For every $\delta > 0$, $\mathbf{CMet}(2_\delta, -) : \mathbf{CMet} \rightarrow \mathbf{CMet}$ preserves directed colimits of convex spaces and isometries.*

PROOF. Let $k_i : K_i \rightarrow K$, $i \in I$ be a directed colimit of convex complete metric spaces and isometries. Consider $f : 2_\delta \rightarrow K$ and choose $\varepsilon > 0$. Denote the two points of 2_δ as x

and y . There is $i \in I$ and $a, b \in K_i$ such that $d(a, fx), d(b, fy) \leq \frac{\varepsilon}{2}$. Then

$$d(a, b) \leq d(a, fx) + d(fx, fy) + d(fy, b) \leq \varepsilon + \delta.$$

Let S be the metric segment in K_i connecting a and b . Choose $a', b' \in S$ such that $d(a, a'), d(b, b') \leq \frac{\varepsilon}{2}$. Then $d(a', b') \leq \delta$. Let $g : 2\delta \rightarrow K_i$ be given by $fx = a'$ and $fy = b'$. Then $d(f, k_i g)$ because $d(a', fx) \leq d(a', a) + d(a, fx) \leq \varepsilon$ and similarly $d(b', fy) \leq \varepsilon$. ■

6.4. EXAMPLE. There is an approximately \aleph_0 -generated Banach spaces which is not finite-dimensional. Consider the complete metric space $A = \{0\} \cup \{\frac{1}{n}; n = 1, 2, \dots\}$. Let $A_m = \{\frac{1}{n}; n = 1, 2, \dots, m\}$ and $r_m : A_m \rightarrow A$ be the inclusion. Let $u_m : A \rightarrow A_m$ be the identity on A_m and send $A \setminus A_m$ to $\frac{1}{m}$. We have $r_m u_m \sim \frac{1}{m} \text{id}_{A_m}$.

Let $F : \mathbf{CMet} \rightarrow \mathbf{Ban}$ be the left adjoint to the unit ball functor $U : \mathbf{Ban} \rightarrow \mathbf{CMet}$ (see [3] 4.5(3)). This adjunction is enriched and thus $F(r_m u_m) \sim \frac{1}{m} \text{id}_{F(A_m)}$. Since $F(1) = \mathbb{C}$, F sends finite discrete metric spaces to finite-dimensional Banach spaces. We have surjective maps $D_m \rightarrow A_m$ where D_m is discrete space of m points. Since U preserves isometries, F preserves \mathcal{E} -maps. Thus $F(A_m)$ are finite-dimensional Banach spaces. Following [3] 7.7(1), $F(A)$ is approximately \aleph_0 -generated.

Since u_m are surjective, $F(u_m)$ are dense. Thus $F(A)$ cannot be finite-dimensional because it has dense maps to m -dimensional Banach spaces for every m .

6.5. LEMMA. An ε -pushout in \mathbf{Ban} is

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow g & & \downarrow \bar{g} \\ C & \xrightarrow{\bar{f}} & B \oplus_{f,g,\varepsilon} C \end{array}$$

where $B \oplus_{f,g,\varepsilon} C$ is the coproduct $B \oplus C$ endowed with the norm

$$\|(x, y)\| = \inf\{\|b\| + \|c\| + \varepsilon\|a\| \mid x = b + f(a), y = c - g(a)\}.$$

PROOF. In the special case of $B \oplus_{f, \text{id}_A, \varepsilon} C$, it is [13] 2.1. The general case is analogous. ■

6.6. COROLLARY. The dual of the full subcategory \mathbf{Ban}_{fd} of \mathbf{Ban} consisting of finite-dimensional Banach spaces is a $(\aleph_0, \mathbf{CMet})$ -nest.

PROOF. Finite-dimensional Banach spaces are closed under dense quotients. Following [3] 4.6 and 6.5, they are closed under finite colimits. ■

6.7. THEOREM. \mathbf{Ban} is equivalent to $\text{Mod}_{\aleph_0}(\mathbf{Ban}_{fd}^{\text{op}})$.

PROOF. Following 4.15, \mathbf{Ban} is equivalent to the full subcategory of $\text{Mod}_{\aleph_0} \mathbf{Ban}_{fd}^{\text{op}}$. Consider H in $\text{Mod}_{\aleph_0} \mathbf{Ban}_{fd}^{\text{op}}$ and follow 4.13(2). For every Banach space X , $H(X)$ is the directed colimit $\text{colim}_d Dd(X)$ of complete metric spaces $Dd(X)$ and isometries. Every $Dd(X)$ is $\mathbf{Ban}(A, X)$ for some finite-dimensional Banach space A . Since these complete metric spaces are convex, we have

$$\mathbf{CMet}(2_\delta, \text{colim}_d Dd(X)) \cong \text{colim}_d \mathbf{CMet}(2_\delta, Dd(X)).$$

Since 1 and $2_\delta, \delta > 0$ form a strong generator in \mathbf{CMet} , $\gamma : H_* \rightarrow H$ is an isomorphism (following the argument from [6] 4.5). ■

6.8. REMARK. The category \mathbf{Ban}^\rightarrow does not seem to be \mathcal{M} -locally \aleph_0 -generated – 4.8 does not apply to it. Hence [3] cannot be immediately applicable to the construction of approximately \aleph_0 -saturated objects in \mathbf{Ban}^\rightarrow like in \mathbf{Ban} . The existence of such objects in \mathbf{Ban}^\rightarrow was proved in [14].

6.9. REMARK. Let \mathbf{CCMet} be the category of convex complete metric spaces and non-expansive maps. \mathbf{CCMet} is an injectivity class in \mathbf{CMet} given by $2_\delta \rightarrow [0, \delta]$, $\delta > 0$, sending the two points of 2_δ to the end-points of $[0, \delta]$. Thus \mathbf{CCMet} is a full weakly reflective subcategory of \mathbf{CMet} . A weak reflection of a complete metric space A to \mathbf{CCMet} is constructed as follows. If $a, b \in A$ are not connected by a metric segment of length $d(a, b)$, we glue $[0, d(a, b)]$ to A by identifying 0 with a and $d(a, b)$ with b . The distances of points of distinct added segments are ∞ . We repeat the procedure and add metric segments $[0, \infty]$. We proceed by induction and the union $\cup_n A_n, n = 1, 2, \dots$ is a desired weak reflection. But \mathbf{CCMet} is not reflective in \mathbf{CMet} because it is not closed under equalizers. Indeed, let C be the circle of radius 1 . Then the equalizer of the identity and the axial symmetry on C is 2_2 . Even, this equalizer does not exist in \mathbf{CCMet} at all. \mathbf{CCMet} does not have coproducts - for instance $1 + 1$ does not exist in \mathbf{CCMet} .

\mathbf{CCMet} is closed in \mathbf{CMet} under \aleph_1 -directed colimits. Indeed, let $k_i : K_i \rightarrow K, i \in I$ be an \aleph_1 -directed colimit of convex complete metric spaces in \mathbf{CMet} . Let $a, b \in K$. Following [3] 2.5(1), there is $i \in I$ and $a', b' \in K_i$ such that $d(a', b') = d(a, b), k_i a' = a$ and $k_i b' = b$. There is $c \in K_i$ such that $d(a', c) + d(c, b') = d(a', b')$. Since $d(a, fc) + d(fc, b) \leq d(a', c) + d(c, b') = d(a, b)$, we have $d(a, fc) + d(fc, b) = d(a, b)$. Thus K is convex. Analogously, using [3] 2.5(2), we prove that \mathbf{CCMet} is closed under directed colimits of isometries in \mathbf{CMet} . We do not know whether the segments $[0, \delta]$ are \aleph_0 -generated with respect to isometries in \mathbf{CCMet} .

The tensor product $A \otimes B$ of convex complete metric spaces A and B is convex and complete. Recall that $A \otimes B$ is $A \times B$ with the $+$ -metric

$$d(a, b), (a', b') = d(a, a') + d(b, b').$$

Indeed, let $a'' \in A$ and $b'' \in B$ satisfy $d(a, a'') + d(a'', a') = d(a, a')$ and $d(b, b'') + d(b'', b') =$

$d(b, b')$. Then

$$\begin{aligned} d((a, b), (a'', b'')) + d((a'', b''), (a', b')) &= d(a, a'') + d(b, b'') + d(a'', a') + d(b'', b') \\ &= d(a, a') + d(b, b') = d((a, b), (a', b')). \end{aligned}$$

We do not know whether **CCMet** is symmetric monoidal closed.

6.10. **REMARK.** Let **CAlg** be the category of C^* -algebras and **CCAlg** the category of commutative C^* -algebras. The forgetful functor $U : \mathbf{CAlg} \rightarrow \mathbf{Ban}$ preserves limits, isometries and \aleph_1 -directed colimits. Thus it has a left adjoint F . The same holds for $U : \mathbf{CCAlg} \rightarrow \mathbf{Ban}$. The unit $\eta_B : B \rightarrow UFB$ is a linear isometry. Thus F is faithful. In the commutative case, this left adjoint was described in [24] and called the Banach-Mazur functor.

The forgetful functor $U : \mathbf{CAlg} \rightarrow \mathbf{Ban}$ even preserves directed colimits. In fact, directed colimits in **Ban** are calculated like in **CMet**, i.e., as a completion of a directed colimit in **Met**. However, the same holds in **CAlg** because one completes the directed colimit of $*$ -algebras. Indeed, if $x = \lim_n x_n$ and $y = \lim_m y_m$ are in this completion then both $x \cdot y = \lim_{n,m} (x_n \cdot y_m)$ and $x^* = \lim_n x_n^*$ are there.

The category of (commutative) C^* -algebras is locally \aleph_1 -presentable and monadic over **Set** (see [22]). Since epimorphisms are surjective in **CAlg** (see [15]), (epi, strong mono)-factorization system on **CAlg** is (surjective, dense) one and it is \aleph_0 -convenient. We expect that **CAlg** is not isometry-locally \aleph_0 -generated.

References

- [1] J. Adámek and J. Rosický, *Locally Presentable and Accessible Categories*, Cambridge University Press 1994.
- [2] J. Adámek and J. Rosický, *What are locally generated categories?*, Proc. Categ. Conf. Como 1990, Lect. Notes in Math. 1488 (1991), 14-19.
- [3] J. Adámek and J. Rosický, *Approximate injectivity and smallness in metric-enriched categories*, J. Pure Appl. Alg. 226 (2022), 106974.
- [4] M. Barr, *Coequalizers and free triples*, Math. Z. (1970), 307-322.
- [5] F. Borceux, *Handbook of Categorical Algebra*, Vol. 2, Cambridge University Press 1994.
- [6] F. Borceux, C. Quinteiro and J. Rosický, *A theory of enriched sketches*, Th. Appl. Categ. 4 (1998), 47-72.
- [7] V. W. Bryant, *The convexity of the subset of a metric space*, Comp. Math. 22 (1970), 383-385.
- [8] C. Centazzo and E. M. Vitale, *A duality relative to a limit doctrine*, Th. Appl. Categ. 10 (2002), 486-497.

- [9] B. Chorny and J. Rosický, *Class-locally presentable and class-accessible categories*, J. Pure Appl. Alg. 216 (2012), 2113-2125.
- [10] B. Day, *On adjoint-functor factorisations*, Lect. Notes in Math. 145 (1970), 1-19.
- [11] L. Fajstrup and J. Rosický, *A convenient category for directed homotopy*, Th. Appl. Cat. 21 (2008), 7-20.
- [12] P. Gabriel and F. Ulmer, *Lokal präsentierbare Kategorien*, Lect. Notes in Math. 221, Springer 1971.
- [13] J. Garbulińska-Węgrzyn, *Isometric uniqueness of a complementably universal Banach space for Schauder decompositions*, Banach J. Math. Anal. 8 (2014), 211-220.
- [14] J. Garbulińska-Węgrzyn and W. Kubiś, *A universal operator on the Gurarii space*, J. Operator Theory 73 (2015), 143-158.
- [15] K. H. Hofmann and K.-H. Neeb, *Epimorphisms of C^* -algebras are surjective*, arXiv:funct-an/9405003.
- [16] G. M. Kelly, *Basic Concepts of Enriched Category Theory*, Cambridge University Press 1982.
- [17] G. M. Kelly, *Structures defined by finite limits in the enriched context*, Cah. Top. Géom. Diff. XXIII (1982), 3-42.
- [18] R. B. B. Lucyshyn-Wright, *Enriched factorization systems*, Th. Appl. Cat. 29 (2014), 475-495.
- [19] M. Makkai and A. Pitts, *Some Results on Locally Finitely Presentable Categories*, Trans. Amer. Math. Soc. 299 (1987), 473-496.
- [20] K. L. Pothoven, *A category of Banach spaces*, Master thesis, Western Michigan Univ. 1968.
- [21] E. Riehl, *Categorical Homotopy Theory*, Cambridge University Press 2014.
- [22] J. W. Pelletier and J. Rosický, *On the equational theory of C^* -algebras*, Alg. Univ. 30 (1993), 275-284.
- [23] J. Rosický and W. Tholen, *Factorizations, fibrations and torsion*, J. Hom. Rel. Str. 2 (2007), 295-314.
- [24] Z. Semadeni, *Some categorical characterizations of algebras of continuous functions*, Symp. Math. XVII (1978), 97-112.
- [25] N. P. Strickland, *The category of CGWH spaces*, preprint 2009.

Ivan Di Liberti
Institute of Mathematics
Czech Academy of Sciences
Žitná 25, Prague, Czech Republic
diliberti.math@gmail.com

Jiří Rosický
Department of Mathematics and Statistics
Masaryk University, Faculty of Sciences
Kotlářská 2, 611 37 Brno, Czech Republic
rosicky@math.muni.cz

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

INFORMATION FOR AUTHORS \LaTeX 2e is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

MANAGING EDITOR. Geoff Cruttwell, Mount Allison University: gcruttwell@mta.ca

TEXNICAL EDITOR. Michael Barr, McGill University: michael.barr@mcgill.ca

ASSISTANT TEX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Julie Bergner, University of Virginia: jeb2md@virginia.edu

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Maria Manuel Clementino, Universidade de Coimbra: mmc@mat.uc.pt

Valeria de Paiva, Nuance Communications Inc: valeria.depaiva@gmail.com

Richard Garner, Macquarie University: richard.garner@mq.edu.au

Ezra Getzler, Northwestern University: getzler@northwestern.edu

Dirk Hofmann, Universidade de Aveiro: dirk@ua.pt

Pieter Hofstra, Université d' Ottawa: phofstra@uottawa.ca

Anders Kock, University of Aarhus: kock@math.au.dk

Joachim Kock, Universitat Autònoma de Barcelona: kock@mat.uab.cat

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: matias.menni@gmail.com

Susan Niefield, Union College: niefiels@union.edu

Kate Ponto, University of Kentucky: kate.ponto@uky.edu

Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

Jiří Rosický, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Michael Shulman, University of San Diego: shulman@sandiego.edu

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Tim Van der Linden, Université catholique de Louvain: tim.vanderlinden@uclouvain.be